UNIVERSIDAD COMPLUTENSE DE MADRID

FACULTAD DE CIENCIAS BIOLÓGICAS



## ESTUDIO HODOLÓGICO E INMUNOHISTOQUÍMICO DE LAS PROYECCIONES ASCENDENTES SOMATOSENSORIALES EN EL SISTEMA NERVIOSO CENTRAL DE LOS ANFIBIOS

V<sup>0</sup> B<sup>0</sup> de los Directores

20.402

Fdo. Dra. Dña. Margarita Muñoz Martín

Fdo. Dr. Hans J. ten Donkelaar

Tesis Doctoral presentada por D. Alberto Muñoz Céspedes para optar al Grado de Doctor en Biología.

Fdo. **D. Alberto Muñoz Céspedes** Madrid, 4 de diciembre de 1995

A Mónica

A mis padres y hermanos

#### AGRADECIMIENTOS

Quiero agradecer a todas las personas que de una forma u otra han colaborado en la realización del presente estudio. Aunque es imposible reflejar todos sus nombres, quisiera mencionar los que de manera más cercana me han brindado su apoyo.

En primer lugar, agradecer a los Directores de este trabajo: a la Dra. Dña Margarita Muñoz Martín, del Departamento de Biología Celular de la Facultad de Biología de la Universidad Complutense de Madrid; y al Dr. Hans J. ten Donkelaar, del Departamento de Anatomía y Embriología de la Facultad de Medicina de la Universidad de Nimega, la confianza que han depositado en mi, los medios que pusieron a mi alcanze, y sus consejos y ayuda incondicional en todo momento.

Hace unos años tuve la fortuna de empezar a formar parte del grupo de investigación dirigido por Margarita y Agustín, en el cual he tenido la oportunidad de iniciarme en el estudio de la Neuroanatomía Comparada. Durante este tiempo, he contado con el privilegio de sus consejos y su amistad. Muchas gracias a los dos por vuestro esfuerzo y por tantos buenos momentos.

Agradecer a todos mis compañeros del Cuarto 12: Ana, Begoña, Paco y Oscar, ya que su paso por este Departamento ha conducido a una amistad que perdura gracias a las experiencias compartidas. En recuerdo de Almudena que, de encontrarse con nosotros, provablemente estaría en la misma situación en la que yo me encuentro actualmente. Gracias a Mónica por su amistad, cariño y ayuda incondicional que me han posibilitado llegar hasta el final.

Gracias a todos los miembros del Departamento de Biología Celular y especialmente a Mar Sánchez y a Ana Fernández por darme ánimos durante todo este tiempo. Quiero agradecer igualmente la ayuda y amistad de todas las personas del Departamento de Anatomía y Embriología de la Facultad de Medicina de la Universidad de Nimega, que han hecho más llevaderas las estancias fuera de casa.

Quisiera mostrar mi gratitud a la Dra. Estrella Rausell del Dpto. de Morfología de la Universidad Autónoma de Madrid, y al grupo del Dr. Jose Ramón Alonso de la Facultad de Biología de la Universidad de Salamanca por su colaboración en las técnicas de tinción inmunohistoquímica de las proteínas ligantes de calcio, e histoquímica para la NADPH diaforasa, respectivamente.

Gracias a los Doctores Wolfgang Walkowiak y Harald Luksch del Instituto de Zoología de la Universidad de Colonia, por la colaboración en el desarrollo de las preparaciones *in vitro* y a Roelie de Boer-van Huizen, Ine Bergeboet-Vernooy y a Jos Dederen de la Universidad de Nimega por su valiosa ayuda técnica.

Por último quisiera agradecer a mi familia la comprensión, el cariño y el apoyo que me han brindado durante todo este tiempo.

## ÍNDICE

CA	PÍTULO 1 INTRODUCCIÓN	7
Introducción histórica		
Consideraciones generales sobre la información somatosensorial		
1.1 MAMÍFEROS		
I.	Receptores periféricos en mamíferos	11
II.	Generalidades sobre la médula espinal de mamíferos: Citoarquitectura	
	y patrón de terminación de las aferencias primarias	12
III.	Proyecciones espinales ascendentes en mamíferos	13
	- Funículo dorsal:	13
	Sistema columna dorsal-lemnisco medial	
	Otras proyecciones	
	- Funículo dorsolateral:	15
	Sistema espino-cervico-talámico	
	Tracto espinocerebeloso dorsal	
	Sistema espino-rombencefalico-talámico	
	Otras proyecciones	
	- Funículos ventral y ventrolateral:	16
	Tracto espinoreticular	
	Tracto espinocerebeloso ventral	
	Tracto espinomesencefálico	
	Tracto espinohipotalámico	
	Tracto espinotalámico	
IV.	Principales modalidades sensoriales y canales funcionales	18
	- Tacto-presión	18
	- Temblor-vibración	19
	- Propiocepción	20
	- Dolor	21
	- Temperatura	22
	- Sensaciones viscerales	23

1.2 VEKTEBRADUS NO	) MAMÍFEROS	24
I. Receptores periféri	cos somatosensoriales en los anfibios	25
II. La médula espinal	de los anfibios	26
- Citoarquitectura		26
- Patrón de termina	ción espinal de las proyecciones de las raíces dorsales	28
III. Proyecciones espir	ales ascendentes	30
- Funículo dorsal		30
Sistema c	olumna dorsal-lemnisco medial, y otras proyecciones incluídas	
en el funí	culo dorsal	
- Funículo dorsola	eral	33
Sistema e	spino-cervico-talámico, y otras proyecciones incluídas en el	
funículo d	lorsolateral	
- Funículos ventral	y ventrolateral (cuadrante ventral)	34
Sistema v	entrolateral	
1.3 BIBLIOGRAFÍA		37
CAPÍTULO 2	CONSIDERACIONES TÉCNICAS	47
CAPÍTULO 2	<b>CONSIDERACIONES TÉCNICAS</b> 2.1 The use of <u>in vitro</u> preparations of the isolated	47
CAPÍTULO 2	<b>CONSIDERACIONES TÉCNICAS</b> 2.1 The use of <u>in vitro</u> preparations of the isolated amphibian central nervous system in neuroanatomy	47
CAPÍTULO 2	<b>CONSIDERACIONES TÉCNICAS</b> 2.1 The use of <u>in vitro</u> preparations of the isolated amphibian central nervous system in neuroanatomy and electrophysiology	<b>47</b> 48
CAPÍTULO 2	<b>CONSIDERACIONES TÉCNICAS</b> 2.1 The use of <u>in vitro</u> preparations of the isolated amphibian central nervous system in neuroanatomy and electrophysiology 2.2 Comentarios	<b>47</b> 48 66
CAPÍTULO 2 CAPÍTULO 3	CONSIDERACIONES TÉCNICAS 2.1 The use of <u>in vitro</u> preparations of the isolated amphibian central nervous system in neuroanatomy and electrophysiology 2.2 Comentarios PROYECCIONES ESPINALES ASCENDENTES	<b>4</b> 7 48 66 <b>7</b> 1
CAPÍTULO 2 CAPÍTULO 3	CONSIDERACIONES TÉCNICAS 2.1 The use of <u>in vitro</u> preparations of the isolated amphibian central nervous system in neuroanatomy and electrophysiology 2.2 Comentarios PROYECCIONES ESPINALES ASCENDENTES 3.1 Spinothalamic projections in amphibians as revealed	47 48 66 71
CAPÍTULO 2 CAPÍTULO 3	CONSIDERACIONES TÉCNICAS 2.1 The use of <u>in vitro</u> preparations of the isolated amphibian central nervous system in neuroanatomy and electrophysiology 2.2 Comentarios PROYECCIONES ESPINALES ASCENDENTES 3.1 Spinothalamic projections in amphibians as revealed with anterograde tracing techniques	<b>47</b> 48 66 <b>71</b> 72
CAPÍTULO 2 CAPÍTULO 3	<ul> <li>CONSIDERACIONES TÉCNICAS</li> <li>2.1 The use of <u>in vitro</u> preparations of the isolated amphibian central nervous system in neuroanatomy and electrophysiology</li> <li>2.2 Comentarios</li> <li>PROYECCIONES ESPINALES ASCENDENTES</li> <li>3.1 Spinothalamic projections in amphibians as revealed with anterograde tracing techniques</li> <li>3.2 Spinal ascending pathways in amphibians: cells of</li> </ul>	<b>47</b> 48 66 <b>71</b> 72
CAPÍTULO 2 CAPÍTULO 3	<ul> <li>CONSIDERACIONES TÉCNICAS</li> <li>2.1 The use of <u>in vitro</u> preparations of the isolated amphibian central nervous system in neuroanatomy and electrophysiology</li> <li>2.2 Comentarios</li> <li>PROYECCIONES ESPINALES ASCENDENTES</li> <li>3.1 Spinothalamic projections in amphibians as revealed with anterograde tracing techniques</li> <li>3.2 Spinal ascending pathways in amphibians: cells of origin and main targets</li> </ul>	<ul> <li>47</li> <li>48</li> <li>66</li> <li>71</li> <li>72</li> <li>81</li> </ul>

CAPÍTULO 4	ONTOGENIA DEL SISTEMA COLUMNA DORSAL-		
	LEMNISCO MEDIAL	132	
	4.1 Early development of dorsal column-medial		
	lemniscal projections in the clawed toad, <u>Xenopus laevis</u>	133	
	4.2 Comentarios	140	
CAPÍTULO 5	EL LEMNISCO MEDIAL EN ANUROS	143	
	5.1 The dorsal column-medial lemniscal projection of		
	anuran amphibians	143	
	5.2 Anuran dorsal column nucleus: Organization,		
	immunohistochemical characterization, and fiber		
	connections in <u>Rana perezi</u> and <u>Xenopus laevis</u>	149	
	5.3 Evidence for an anuran homologue of the mammalian		
	spinocervicothalamic system: An i <u>n vitro</u> tract-tracing study		
	in <u>Xenopus laevis</u>	188	
	5.4 Comentarios	208	
CAPÍTULO 6	PLACA ALAR DEL ROMBENCÉFALO CAUDAL		
	EN URODELOS	218	
	6.1 Organization of the caudal rhombencephalic alar plate		
	of the ribbed newt <u>Pleurodeles waltl</u> : Evidence for the presence		
	of dorsal column and lateral cervical nuclei	219	
	6.2 Comentarios	249	
CAPÍTULO 7	CONCLUSIONES	256	

# CAPÍTULO 1

Introducción

7

Las funciones del sistema nervioso de los vertebrados incluyen la percepción de distintos estímulos del medio que les rodea, lo que permite la elaboración de respuestas adecuadas para cada tipo de situación, permitiendo así su relación con su entorno. Los estímulos ambientales son captados por los receptores sensoriales, encargados de transducirlos a un código bioeléctrico de potenciales de acción inteligibles para el sistema nervioso, y transportarlos por medio de los nervios sensitivos o aferentes al sistema nervioso central, el cual, después de procesar y analizar la información en diversos centros específicos, elabora respuestas que son conducidas a través de los nervios eferentes hasta los órganos efectores.

#### Introducción histórica

El filósofo y naturalista griego, Aristóteles (322 a. de C.), describió los sentidos en su obra *Historia de los animales*, como las vías o los canales a través de los cuales los animales perciben información de su entorno. Propuso la existencia de cinco sentidos comunes para la mayoría de los animales, entre los que se encontraba el tacto, actualmente considerado una parte integrante del concepto de somestesia o información somatosensorial.

El hecho de considerar las sensaciones cutáneas como una entidad independiente y específica se remonta a los estudios fisiológicos realizados a principios del siglo XIX, (ver Sinclair, 1981), y a las observaciones anatómicas realizadas por Bell (1811). Posteriormente, Magendie (1822) demostró la implicación de las raíces dorsales de los nervios

espinales en la transmisión de la información cutánea. En estudios sucesivos, Müller (1840-42), retomando la doctrina Aristotélica de los cinco sentidos, introdujo el término tacto como la agrupación de las diferentes sensaciones cutáneas. Sin embargo, durante la década de 1840 a 1850, de acuerdo con Sinclair (1981), varios autores comenzaron a establecer la teoría de la especificidad, al distinguir entre los diversos aspectos de la información cutánea, postulando la existencia de terminales nerviosos específicos para cada sensación. Describieron los criterios de irritabilidad selectiva, es decir, cada tipo de órgano sensorial podría ser excitado por un estímulo en particular, que a su vez sería conducido por un nervio concreto. Las citadas hipótesis fueron corroboradas fisiológicamente (Blix, 1884; Goldscheider, 1884, Donaldson, 1885). Blix (1884) demostró que la estimulación específica de distintos puntos con una localización concreta en la piel, producía sensaciones de presión, calor, frio o dolor; y confirmó la especificidad anteriormente propuesta, mediante la demostración de que cada punto responde a una modalidad concreta de estimulación y no a otras. Se localizaron puntos sensoriales que respondían específicamente a variaciones térmicas, a estímulos de presión o de dolor, y se observó que la densidad de la distribución de cada uno de ellos variaba en función de la región corporal analizada (Blix, 1884, Goldscheider, 1884; von Frey, 1896; Dallenbach, 1927). Completando estas valoraciones, von Frey (1906, 1910) postuló la posibilidad de que cada clase de punto sensitivo, podría estar asociado a un órgano sensorial en particular, que previamente había sido identificado histológicamente (Krause, 1859, Meissner, 1859, Ruffini, 1894), sin embargo, su correlación ha sido corregida en estudios morfofuncionales posteriores (ver Willis y Coggeshall, 1991)

Como consecuencia de la transmisión codificada de los estímulos en las fibras que inervan los órganos sensoriales Adrian (1946), Sinclair (1955) y Weddell (1955) propusieron la Teoría de los patrones que sugiere que no es necesaria la existencia de vías o canales específicos para cada modalidad de estimulación cutánea. Por el contrario, la especificidad de la transmisión de los distintos estímulos, vendría determinada por patrones espaciales y temporales de codificación diferencial de la información, empleando las mismas vías. Dicha teoría, sin embargo, ha sido parcialmente rechazada a la luz de observaciones que ofrecen evidencias de la existencia de fibras morfológica y funcionalmente distintas, por ejemplo: mielínicas o amielínicas; o con diámetros y velocidades de conducción desiguales, encargadas de la transmisión de los diferentes estímulos (ver Willis y Coggeshall, 1991). La especificidad presente en las vías anatómicas del sistema nervioso central, en relación a la naturaleza de la información que procesan (Noordenbos y Wall, 1976), apoya igualmente dicha teroría, al demostrar que la información nociceptiva, térmica, y algunos aspectos de la información tactil, son transmitidos a través del cuadrante anterolateral de la médula espinal, contrariamente a las sensaciones mecánicas que ascienden por sistemas espinales diferentes. Los resultados obtenidos en experimentos realizados en humanos con técnicas de microneurografía, consistentes en la evocación de sensaciones mediante la estimulación de una única fibra sensorial periférica, y la posterior identificación de su campo receptivo y del estímulo necesario para su excitación, apoyan igualmente la teroría de la

especificidad de inervación de los diversos órganos sensoriales por distintas fibras aferentes (Willis y Coggeshall, 1991).

## Consideraciones generales sobre la información somatosensorial

Existen diversas clasificaciones de la información somática, atendiendo a criterios como la naturaleza del estímulo, su procedencia y la posible evocación de sensaciones.

La sensibilidad somatosensorial se divide en información somática visceral e información somática no visceral. La primera produce sensaciones conscientes como el dolor visceral, la saciedad producida por estímulos de distensión, así como la percepción del grado de vaciamiento de las vísceras; e información no consciente que participa en la regulación del medio interno. La sensibilidad somática no visceral consiste en la percepción de estímulos básicos considerados como primarios, en los que se incluyen los que conducen a sensaciones de dolor, térmicas y los estímulos mecanorreceptivos, que a su vez se dividen en exteroceptivos -si provienen del exterior del organismo (tacto-presión y temblorvibración)-, y en propioceptivos, -si se originan en el interior del organismo como los responsables de las sensaciones de posición y de movimiento (cinestesia) de los miembros y de las articulaciones-. Asimismo, la sensibilidad somática no visceral incluye también la percepción de estímulos secundarios más complejos, como el sentido espacial y el reconocimiento de las formas, resultado de la combinación e integración de la percepción de

estímulos primarios (Mountcastle, 1973; Loewy y Spyer, 1990; Willis y Coggeshall, 1991).

En términos generales, la información somatosensorial puede clasificarse en *especial* y *general* dependiendo de que ésta proceda de la región cefálica o del tórax y las extremidades respectivamente. La información somática *especial* ingresa en el sistema nervioso central a través de las ramas sensitivas de algunos pares craneales (V, VII, IX, X y XII), a diferencia de la información *general* que lo hace a través de las raíces dorsales de los nervios espinales.

En 1920 Head distinguió entre sensibilidad epicrítica, cuando se trata de información inocua tactil o de pequeñas variaciones térmicas, y sensibilidad protopática para las aferencias nociceptivas o de sensaciones de temperaturas extremas. Dicha nomenclatura, actualmente en desuso (Willis y Coggeshall, 1991), ha sido empleada para calificar y diferenciar algunas vías centrales somestésicas de manera incorrecta, debido a que en estudios posteriores se ha demostrado que ambos tipos de información están entremezclados a nivel de sistema nervioso central.

Igualmente, Poggio y Mountcastle (1960; 1963) describieron los conceptos *lemniscal* y *no lemniscal* con el fin de clasificar patrones diferenciales de respuesta fisiológica en neuronas de los complejos ventrobasal y posterior talámicos respectivamente. Los autores supusieron que las neuronas del complejo ventrobasal eran activadas por el sistema de la columna dorsal-lemnisco medial, mientras que las del complejo porterior lo eran a través del tracto espinotalámico (no lemniscal). Sin embargo, en trabajos sucesivos (Boivie y Perl, 1975) se ha criticado dicha nomenclatura como consecuencia de que tanto algunas de las neuronas que originan el lemnisco medial, como las que proyectan a través del tracto espinotalámico, comparten características que fueron consideradas por Poggio y Mouncastle (1960) como distintivas de ambos sistemas, y a que las dos poblaciones neuronales inervan los complejos ventrobasal y posterior talámicos (Boivie y Perl, 1975).

El término de modalidad sensorial fue introducido por Helmholtz (ver Boring, 1942) como una clase de sensaciones cualitativamente continuas, entre las que únicamente existen diferencias cuantitativas como las que presentan dos sensaciones tactiles con intensidades desiguales. En contraposición, las diferencias presentes entre las sensaciones correspondientes a modalidades sensoriales distintas, por ejemplo el tacto y el oido, son cualitativas. Aunque la información procedente de la piel se consideró inicialmente de forma global (Müller, 1840-42), actualmente se distinguen diversas modalidades sensoriales cutáneas: tacto-presión, temblor-vibración, cosquilleo, calor, frio, dolor, picor; así como las subcutáneas propioceptivas que incluyen el sentido de la posición y la cinestesia (Willis y Coggeshall, 1991).

La transmisión de cada modalidad sensorial en la somestesia depende de la activación y el funcionamiento de una o varias vías anatómicas en el sistema nervioso central. Recientemente, Willis y Coggeshall (1991) han introducido el término de *canal sensorial* que engloba todos los mecanismos necesarios para la percepción de la información relacionada con una modalidad sensorial, con la posibilidad de incluir diversos tipos de receptores sensoriales, vías anatómicas de transmisión de información y regiones cerebrales encargadas de su procesamiento.

### MAMÍFEROS

#### Receptores periféricos en mamíferos.

Los receptores de las distintas modalidades sensoriales fueron inicialmente descritos por Bell (1811) y Magendie (1822) como las prolongaciones periféricas de las neuronas de los ganglios de las raíces dorsales espinales, cuyas prolongaciones centrales se ramifican en la médula espinal. Los receptores se han clasificado en función de la naturaleza y el umbral del estímulo o los estímulos necesarios para su activación, así como por la morfología de sus terminales en los tejidos periféricos, libres o encapsulados, (Perl, 1992) y por las diferencias en cuanto a su comportamiento electrofisiológico (Koerber y Mendell, 1992). Las células ganglionares llevan a cabo la recepción de los estímulos, su transducción en señales inteligibles, y la conducción de las mismas hasta el sistema nervioso central (Perl, 1922) y se clasifican atendiendo a numerosos parámetros.

Morfológicamente, mediante tinciones de Nissl, se distinguen dos grupos de células ganglionares, con la posibilidad de nuevas clasificaciones, atendiendo a observaciones con ténicas de microscopía electrónica. Ambas poblaciones neuronales se han descrito como células grandes que se tiñen débilmente y células de menor tamaño que se tiñen con mayor intensidad (Andres, 1961; Lieberman, 1976; Duce y Keen, 1977; Lawson, 1979; Rambourg y cols., 1983), debido a su diferente contenido en orgánulos citoplásmicos (Yamadori, 1970; Duce y Keen, 1977). Las células grandes y claras se caracterizan por su alto contenido en neurofilamentos y por dar origen a fibras de tipo A. Las células pequeñas y oscuras, por el contrario, tienen un escaso contenido en neurofilamentos y originan fibras de tipo C (ver Lawson, 1992).

Mediante técnicas inmunocitoquímicas y de hibridación *in situ*, se han identificado una gran variedad de neuronas ganglionares en cuanto a su patrón de expresión de uno o varios péptidos, enzimas, filamentos intermedios, neurohormonas, monoaminas, proteínas ligantes de calcio, receptores y oligosacáridos de superficie. Sin embargo, todavía no existe una correlacción clara entre estas subpoblaciones, definidas por su contenido neuroquímico, y los distintos tipos funcionales de células ganglionares, identificados según el destino de su inervación periférica, o por su patrón de terminación en la sustancia gris espinal (Rustioni y Weimberg, 1989; Lawson, 1992; Hunt y cols., 1992).

La velocidad de conducción del estímulo en las fibras aferentes varia en relacción a su diámetro y a su grado de mielinización. Según estos criterios, existen fibras de tipo A, mielinizadas y de alta velocidad de conducción, entre las que se distinguen tres subgrupos (alfa, beta y delta) ordenadas de mayor a menor según la velocidad de conducción y grado de mielinización (Perl, 1992). En contraposición, las fibras C tienen menor diámetro, son amielínicas y tienen una menor velocidad de conducción (Gasser, 1950, 1955). Las aferencias musculares, sin embargo, se clasifican en los grupos I, II III y IV ordenadas de mayor a menor diámetro (Chang, 1948).

## Generalidades sobre la médula espinal de mamíferos: citoarquitectura y patrón de terminación de las aferencias primarias.

La sustancia gris de la médula espinal de mamíferos ha sido dividida en diez regiones denominadas láminas y numeradas del I al X (Rexed, 1952, 1954) atendiendo a criterios citoarquitectónicos como el tamaño, el agrupamiento y la morfología celular. Sin embargo, dicha subdivisión morfológica no refleja una subdivisión funcional (Fyffe, 1992).

A lo largo de todo el espesor del asta dorsal, que incluye las láminas I-VI de Rexed (1952) se ha descrito en cada segmento espinal, un mapa somatotópico de representación de la superficie corporal, correspondiente a un dermatomo (Sherrington, 1898; Brown y cols., 1992), de forma que cada raíz dorsal inerva una región corporal o dermatomo diferente (Brown y Fuchs, 1975; Brown y cols., 1980; Light y Durkovick, 1984; Wilson y cols., 1986). Dentro de este esquema de organización, los nervios que inervan regiones advacentes de la piel, tienen proyecciones adyacentes en el asta dorsal (Koerber y Brown, 1980, 1982). Según la hipótesis de la somatotopía presináptica (Brown y Finch, 1975), debe existir una correlación entre la extensión de los mapas de representación de cada región cutánea, observados electrofisiológicamente, y la extensión de

las arborizaciones, en la médula espinal, de las aferencias primarias procedentes de dicha región corporal. Esta hipótesis parece cumplirse al menos en el caso de las aferencias cutáneas (Brown y cols., 1992), aunque podría ser igualmente válida en el caso de las aferencias procedentes de otros tejidos.

El asta dorsal de la médula espinal se puede dividir en tres regiones diferentes, en cuanto a la naturaleza de sus aferencias, lo que podría reflejar su organización funcional. En general, las aferencias mielínicas, A delta, procedentes de los nociceptores cutáneos terminan en las láminas I, V y en la parte externa de la lámina II. Las aferencias A beta, procedentes de los mecanorreceptores cutáneos, se arborizan en las regiones profundas del asta dorsal, que incluyen las láminas III, IV y la zona interna de la lámina II, Por último, las aferencias musculares de tipo I y II, alcanzan las láminas VI, VII y IX y ocasionalmente, la III y la IV (Brown y cols., 1977, 1978, 1980, 1981; Hamano y cols., 1978; Hongo y cols., 1978; Isizuka y cols., 1979; Matsushita y Tanami, 1983; ver Fyffe, 1992 para una revisión). Las fibras C amielínicas terminan mayoritariamente en la lámina II en el caso de los mecanoreceptores cutáneos, mientras que las fibras C nociceptivas, procedentes de receptores polimodales y de frio, se arborizan principalmente en las láminas I y II y en menor medida en la III y la IV. Por último, las aferencias C de origen visceral terminan separadamente en las láminas I, II, V y X (Sugiura 1986, Sugiura y cols., 1989). La región superficial del asta dorsal fue identificada por Rolando (1824) a consecuencia de su aspecto gelatinoso, incluye las láminas I y II de Rexed (1952) y representa un centro de procesamiento inicial de la información nociceptiva

(Cerveró, 1989). En esta región superficial se diferencian la zona marginal y la sustancia gelatinosa correspondientes a las láminas I y II respectivamente (Cerveró e Iggo, 1980). En esta zona fundamentalmente convergen aferencias nociceptivas y fibras amielínicas de tipo C. El asta dorsal intermedia formada por las láminas III y IV, recibe mayoritariamente aferencias mielínicas procedentes de mecanoreceptores cutáneos, mientras que las regiones más profundas del asta dorsal están inervadas por aferencias musculares (Fyffe, 1992).

## Proyecciones espinales ascendentes en mamíferos.

En vertebrados terrestres se ha descrito la existencia de dos sistemas básicos de proyecciones espinales ascendentes (Willis y Coggeshall, 1991). 1) El Sistema columna dorsal-lemnisco medial, constituído por axones primarios procedentes de neuronas ganglionares espinales, así como por fibras no primarias que forman el sistema postsináptico de la columna dorsal. Ambos tipos de fibras ascienden en el funículo dorsal hasta los núcleos de la columna dorsal, situados en la placa alar del rombencéfalo caudal, que a su vez originan el lemnisco medial que asciende contralateralmente para alcanzar diversas dianas rombencefálicas, mesencefálicas y talámicas. 2) El sistema ventrolateral, formado por fibras espinales no primarias que ascienden a través del cuadrante ventrolateral (funículos ventral y ventrolateral), hasta la formación reticular, el cerebelo (tracto espinocerebeloso ventral, ver Ito (1984)), el mesencéfalo (Yezierski, 1988), el hipotálamo (Burstein y cols, 1990; Dado y cols., 1994) y el tálamo (Willis y Coggeshall, 1991).

Además, en mamíferos se ha descrito un tercer sistema denominado *espino-cervico-talámico* compuesto por fibras no primarias espinales que ascienden a través del **funículo dorsolateral**, formando el tracto espinocervical que termina en el núcleo cervical lateral. Dicho núcleo origina proyecciones contralaterales que incluyen los tractos cervicomesencefálico y cervicotalámico, que ascienden a través del lemnisco medial hasta el mesencéfalo y el tálamo respectivamente (Willis y Coggeshall, 1991). Asimismo, en el funículo dorsolateral ascienden otras proyecciones como los tractos espinocerebeloso dorsal (Ito, 1984), espinohipotalámico y espinotalámico, y el sistema espino-rombencefálico-talámico (Willis y Coggeshall, 1991).

Los tres sistemas ascendentes están sujetos a un control descendente, a través de proyecciones que se originan en distintas regiones corticales y talámicas, en el núcleo rojo, cerebelo, formación reticular, complejo nuclear del rafe y diversos núcleos rombencefálicos cocleares y vestibulares (ver Willis y Coggeshall, 1991).

#### Funículo dorsal:

### - Sistema Columna Dorsal-Lemnisco Medial.

Dicho sistema está formado por las ramas ascendentes de las aferencias primarias de todos los niveles espinales, organizadas somatotópicamente; es decir, la parte medial del funículo dorsal o fascículo gracilis contiene las fibras originadas en niveles espinales caudales a los niveles medios torácicos, que transmiten información procedente de las extremidades inferiores y de las regiones caudales del torax, y terminan en el núcleo gracilis, que ocupa las posiciones más mediales dentro del complejo de los núcleos de la columna dorsal. Únicamente un 20% de las fibras primarias que entran en el fascículo gracilis, ascienden hasta el rombencéfalo, mientras que el resto terminan en el asta dorsal, la región intermedia y el asta ventral de la sustancia gris espinal (Glees y Soler, 1951). Las fibras primarias que terminan en la médula espinal, antes de penetrar en la sustancia gris, ascienden diferentes distancias en el funículo dorsal, dependiendo del tipo de receptor periférico y de la naturaleza de la información transmitida (ver Willis y Coggeshall, 1991). Las aferencias procedentes de segmentos torácicos superiores y cervicales, que transmiten información de las extremidades superiores y de la región superior del tórax, ascienden en el fascículo cuneatus, situado lateralmente en el funículo dorsal, para terminar en el núcleo *cuneatus* localizado en la región lateral del complejo nuclear de la columna dorsal. En algunas especies de vertebrados se ha descrito el núcleo de Bischoff en la línea media de los núcleos de la columna dorsal que recibe que información de la cola (Kappers y cols., 1936).

Las aferencias primarias espinales emiten colaterales a distintos niveles rostrocaudales dentro del mismo plano parasagital de los núcleos de la columna dorsal, lo que permite establecer un mapa de representación somatotópica de la superficie corporal, que se mantiene constante en toda la extensión de los núcleos (Willis y Coggeshall, 1991).

El funículo dorsal contiene, además de las aferencias primarias, axones de neuronas situadas

fundamentalmente en las láminas espinales III, IV, V y VI y en menor grado en las I, III, y VII, que proyectan a los núcleos de la columna dorsal, y constituyen el denominado sistema postsináptico de la columna dorsal. Éste participa en la transmisión de distintas modalidades de sensaciones y presenta un ordenamiento somatotópico similar al de las aferencias primarias espinales, es decir, las neuronas localizadas en los niveles cervical y lumbar proyectan a los núcleos *cuneatus* y gracilis respectivamente (ver Willis y Coggeshall, 1991).

En los núcleos de la columna dorsal se establece un mapa de representación somatotópico que incluye el procesamiento de información propioceptiva y exteroceptiva. En mamíferos los núcleos de la columna dorsal presentan distintas regiones citoarquitectónicas desde las que se organizan sus diferentes eferencias. Éstas incluyen una proyección descendente a la médula espinal, así como un conjunto de proyecciones ascendentes, fundamentalmente contralaterales, que constituyen el lemnisco medial. Entre las dianas diencefálicas del lemnisco medial se encuentran: la zona incerta, la región existente entre los núcleos ventrolateral y ventral posterior que recibe una proyección fundamentalmente propioceptiva, la parte medial del complejo posterior talámico, así como el núcleo ventral posterolateral del complejo ventrobasal talámico. Éste último recibe la proyección cuantitativamente más importante, relacionada fundamentalmente con la información tactil y en la cual se produce una inversión espacial del mapa de representación somática, presente en los núcleos de la columna dorsal, de manera que las proyecciones del núcleo gracilis terminan lateralmente a las procedentes

del núcleo *cuneatus* (ver Berkley y cols., 1986; y Willis y Coggeshall., 1991). Otras proyecciones del lemnisco medial alcanzan la región pretectal así como el núcleo rojo, colículo superior y núcleo intercolicular en el mesencéfalo; el cerebelo; y diferentes núcleos pontinos y de la oliva inferior en el rombencéfalo (ver Willis y Coggeshall, 1991).

#### - Otras proyecciones.

Además de las proyecciones anteriormente descritas, en relación con los núcleos de la columna dorsal; Holbrook y Wilcox (1965) observaron en el funículo dorsal de la cabra, mediante técnicas de degeneración, aferencias primarias espinales que sobrepasan rostralmente los núcleos de la columna dorsal y ascienden a través del rombencéfalo hasta el cerebelo, aunque su presencia no ha sido confirmada en trabajos posteriores en otras especies de mamíferos.

Igualmente se ha demostrado la existencia en el funículo dorsal del componente dorsal del tracto espinoreticular, que consiste en una proyección bilateral, originada principalmente en neuronas situadas en las láminas I y X y en menor medida en las láminas II-IV y VII, y que termina en el núcleo reticular dorsal (McMahon y Wall, 1985; Lima, 1990; Lima y Coimbra, 1990; Bing y cols., 1990).

#### Funículo dorsolateral:

- Sistema espino-cervico-talámico.

En mamíferos se ha descrito la existencia de una vía bisináptica formada por los tractos espinocervical y cervicotalámico (Morin, 1955; Nijensohn y Kerr, 1975; Boivie, 1983). El tracto espinocervical consiste en una proyección, fundamentalmente ipsilateral, que se origina en neuronas del asta dorsal situadas a lo largo de la médula espinal en el núcleo propio, láminas II, IV y V, y en niveles cervicales además en las láminas I, VI y VII. El tracto espinocervical asciende en el funículo dorsolateral y termina en el núcleo cervical lateral (Willis y Coggeshall, 1991). A lo largo de su recorrido el tracto espinocervical emite colaterales que alcanzan distintas dianas intraespinales, estableciendo conexiones intersegmentarias (Snow y cols., 1976; Rastad y cols., 1977; Jankowska y cols., 1979; Maxwell y Koerber, 1986; Cao y cols., 1993). El núcleo cervical lateral es una población neuronal propia de segmentos cervicales superiores (C1-C3), ventrolateral al asta dorsal, y segregada dentro del funículo dorsolateral. Dicho núcleo organiza proyecciones ascendentes contralaterales que alcanzan las regiones somatosensoriales mesencefálicas y talámicas, a través de los tractos cervicomesencefálico y cervicotalámico respectivamente, los cuales forman parte del lemnisco medial (ver Willis y Coggeshall, 1991). El tracto cervicomesencefálico inerva principalmente el núcleo intercolicular y las capas profundas del colículo superior, mientras que el cervicotalámico termina mayoritariamente en el núcleo ventral posterolateral y en la parte medial del complejo posterior talámico. El núcleo lateral cervical origina también proyecciones descendentes a la médula espinal (Svensson y cols., 1985b).

#### - Tracto espinocerebeloso dorsal.

En mamíferos el tracto espinocerebeloso dorsal o tracto de Flechsig está formado por fibras mielínicas que transmiten información sensorial propioceptiva, procedente de husos muculares y receptores tendinosos de Golgi de las extremidades posteriores, asciende en la parte dorsal del funículo dorsolateral y penetra en el cerebelo a través del cuerpo restiforme (Yoss, 1952a), para terminar en distintas regiones cerebelosas (Grant, 1962b; Manns, 1973; Matsushita y Okado, 1981; Ito, 1984). Dicha proyección se origina en neuronas de la columna de Clarke, situada en la base del asta dorsal en niveles torácicos y lumbares (hasta L4), que reciben aferencias desde los funículos dorsal y lateral así como de neuronas del asta dorsal. Sin embargo, se han descrito neuronas en niveles cervicales, que proyectan al cerebelo, cuyos axones podrían ascender a través del funículo dorsolateral (Ito, 84).

#### - Sistema espino-rombencefalico-talámico.

El sistema espino-rombencefálico-talámico está formado por las proyecciones de las neuronas espinales de la columna de Clarke, las cuales reciben aferencias propioceptivas, procedentes de receptores musculares de tipo Ia de los miembros posteriores. Los axones de las neuronas de la columna de Clarke ascienden en el funículo dorsolateral formando el tracto espinocerebeloso dorsal, que emite colaterales hacia la sustancia gris para sinaptar en el núcleo Z del rombencéfalo. Dicho núcleo establece proyecciones contralaterales, a través del lemnisco medial, que terminan en el núcleo ventral posterolateral así como en la región de éste que limita con el núcleo ventral lateral (Landgren y Silfvenius, 1969; 1971; Low y cols., 1986; Willis y Coggeshall, 1991).

#### - Otras proyecciones.

Otros tractos espinales ascendentes en el funículo dorsolateral incluyen las proyecciones espinomesencefálicas (Zelman y col., 1978; McMahon y Wall, 1983, 1985; Baker y Giesler, 1984; Swett y cols., 1985), espinohipotalámicas (Dado y cols., 1994c) y espinotalámicas (Jones y cols., 1985, 1987; Apkarian y Hodge, 1989a,b; Dado y cols., 1994c), descritas en algunos mamíferos, y que se originan en neuronas situadas en diversas regiones espinales mayoritariamente en el asta dorsal.

## Funículos ventral y ventrolateral:

En mamíferos el sistema ventrolateral que ocupa los funículos ventral y ventrolateral, está formado principalmente por los tractos espinoreticular, espinocerebeloso ventral, espinomesencefálico, espinohipotalámico y espinotalámico.

#### - El tracto espinoreticular.

En el funículo ventrolateral existen dos tipos de proyecciones espinoreticulares. La primera se origina en neuronas de las láminas V, VII y VIII, y de manera muy escasa en la lámina X así como en el asta dorsal superficial de todos los niveles espinales, aunque más densamente desde niveles cervicales superiores (ver Willis y Coggeshall, 1991) y alcanza fundamentalmente el núcleo reticular lateral, que a su vez proyecta al cerebelo (Oscarsson, 1973). La segunda consiste en el componente medial del tracto espinoreticular que asciende hasta diversos núcleos del tronco cerebral que proyectan a la médula espinal, al mesencéfalo o a los núcleos intralaminares del tálamo (Peschanskiy Besson, 1984; Willis y Coggeshall, 1991).

#### - Tracto espinocerebeloso ventral.

En mamíferos la proyección espinocerebelosa ventral o tracto de Grower está formada por fibras mielínicas que ascienden en la superficie ventrolateral de la médula espinal para entrar en el cerebelo, a través del pedúnculo cerebeloso superior (Yoss, 1952). Dicha proyección se origina en neuronas de gran tamaño, localizadas fundamentalmente en la sustancia gris dorsomedial y ventrolateral del asta ventral espinal en niveles lumbares (Cooper y Sherrington, 1940; Hubbard y Coscarsson, 1962; Lundberg y Weight, 1971; Burke y cols., 1971), que reciben información propioceptiva procedente de husos musculares y de receptores tendinosos de Golgi de las extremidades posteriores, así como de diversas proyecciones descendentes (Ito, 1984). Además, algunas fibras de las neuronas espinales que proyectan al cerebelo desde niveles cervicales, podrían igualmente ascender a través del funículo ventrolateral (Ito, 1984).

#### - Tracto espinomesencefálico.

Algunas proyecciones espinomesencefálicas ascienden a través del funículo dorsolateral como se ha expuesto anteriormente, sin embargo otras, esencialmente contralaterales, ascienden por el sistema ventrolateral, e incluyen proyecciones que se originan en diversas poblaciones neuronales de la médula espinal y terminan en tres regiones mesencefálicas diferenciables: 1) una zona de transición entre el rombencéfalo y el mesencéfalo, que incluye el área parabraquial, el núcleo cuneiforme y la sustancia gris periacueductal más caudal; 2) la región intermedia, que comprende el núcleo intercolicular, el núcleo cuneiforme, la sustancia gris periacueductal correspondiente a dicha zona y las capas profundas del colículo superior; y 3) niveles mesencefálicos rostrales que incluyen los núcleos de Darkschewitsch, pretectales anterior y posterior, rojo, de Edinger-Westphal e intersticial de Cajal (Yezierski, 1988). Las proyecciones espinomesencefálicas se originan en las láminas I, III, V, VII, VIII, X y en el núcleo espinal lateral, en toda la extensión rostrocaudal de la médula, así como en el núcleo cervical lateral (ver Willis y Coggeshall, 1991).

#### - El tracto espinohipotalámico.

En mamíferos el tracto espinohipotalámico es una proyección bilateral, aunque con predemonio contralateral, que se origina mayoritariamente en neuronas de las regiones profundas del asta dorsal, en el núcleo espinal lateral y en menor medida en el asta dorsal superficial, la sustancia intermedia y la región que rodea al canal central (Burstein y cols., 1990). En diversos trabajos se ha descrito que las proyecciones espinales alcanzan regiones mediales, laterales, ventrales y posteriores hipotalámicas, así como la región de la decusación supraóptica (Anderson y Berry, 1959; Minderhoud, 1967; Yamada y Otani, 1978; Kerr, 1975; Ju, 1984; Burstein y cols., 1987, 1990; Dado y cols., 1984a,b,c). La proyección espinohipotalámica asciende a través de los funículos ventrolateral y en menor grado dorsolateral (Dado y cols., 1994c).

- El tracto espinotalámico.

El tracto espinotalámico de mamíferos es una proyección principalmente contralateral, que se origina en neuronas situadas en todos los niveles espinales, aunque su máxima densidad corresponde a niveles cervicales superiores, donde se observa además la mayor concentración de células de proyección ipsilateral. En el resto de la médula espinal se origina mayoritariamente en los niveles de las intumescencias, cervical y en menor número de la lumbar. Se han descrito tres componentes en la proyección espinotalámica, en función de la localización de las células de origen, y de los centros de destino. 1) Constituído a patir de neuronas situadas mayoritariamente en la lámina I, aunque también en las láminas II y III, y que proyectan principalmente al núcleo ventral posterolateral y en menor grado al núcleo submedio y a los núcleos intralaminares. 2) Se organiza a partir de neuronas de las regiones profundas del asta dorsal (láminas IV-VI) y termina en los núcleos ventral posterolateral y centrolateral. 3) formados por axones de neuronas de las láminas VII-X que proyectan al núcleo centrolateral. El complejo posterior recibe igualmente proyecciones espinotalámicas (Willis Coggeshall, 1991).

## Principales modalidades sensoriales y canales funcionales

#### Tacto-presión

El tacto y la presión constituyen dos componentes de una modalidad sensorial cuyas diferencias residen en la intensidad del estímulo. Ambos son percibidos a través de dos tipos de mecanorreceptores de adaptación lenta, SAI y SAII, que corresponden a los receptores de Merkel y de Ruffini respectivamente. Los receptores de Merkel (SAI) desempeñan un papel más importante que los de Ruffini (SAII), en esta modalidad sensorial, ya que intervienen en la transmisión de tacto y de presión, mientras que los receptores SAII producen sensaciones de presión mantenida, y están más implicados en la transmisión de información propioceptiva de sentido espacial (Harrington y Merzenich, 1970, Knibestöl y Vallvo, 1970; Willis y Coggeshall, 1991).

Las aferencias procedentes de los receptores SAI y SAII terminan en el asta dorsal en el núcleo propio y las láminas III, IV y V (Brown, 1977, Brown y cols., 1978). La información de tacto y presión es transmitida a niveles supraespinales a través de los funículos dorsal y dorsolateral.

En el funículo dorsal la información tactil, procedente de las extremidades anteriores, es transmitida por fibras primarias espinales, mientras que la de las extremidades posteriores asciende a través del sistema postsináptico de la columna dorsal. La información alcanza, por ambos sistemas, los núcleos de la columna dorsal que posteriormente la transmiten, a través del lemnisco medial, a diversas áreas mesencefálicas y talámicas. Las fibras tactiles en el funículo dorsolateral están en relación con aferencias a los núcleos de la columna dorsal, si bien las del tracto espinocervical parecen no estar implicadas (ver Willis y Coggeshall, 1991). Willis y cols. (1974) demostraron cierto grado de implicación del cuadrante ventrolateral espinal, en la transmisión de información tactil.

#### **Temblor-Vibración**

Esta modalidad sensorial incluye varios tipos de sensaciones, principalmente el temblor y la vibración, percibidas al aplicar en la superficie corporal estímulos oscilatorios de distintas frecuencias, entre 5 y 40 Herz o superior a 60 Herz respectivamente (Willis and Coggeshall, 1991). Su percepción en mamíferos parece depender de la activación de mecanorreceptores de bajo umbral y adaptación rápida, FAI (temblor) y FAII (vibración), localizados en la piel y en tejidos subyacentes respectivamente (Limdblom y Lund, 1966), y que se distinguen entre morfológica sí y electrofisiológicamente (Willis and Coggeshall, 1991). Los receptores FAI incluyen a los copúsculos de Meissner en las regiones de piel sin pelo, y a los receptores de campo y de folículo piloso, en las regiones de piel dotadas de pelo. Los receptores FAII corresponden a los corpúsculos de Paccini (Lindblom y Lund, 1966; Knibestöl, 1973).

En los primeros experimentos, basados en técnicas microneurográficas, se sugirió la implicación de los mecanorreceptores de adaptación rápida (FAI y FAII) en la evocación de sensaciones descritas como tacto, presión, cosquilleo y vibración (Vallvo, 1981), aunque en estudios posteriores (Schady y cols., 1983; Vallvo y cols., 1984) se han relacionado únicamente con la percepción de temblor, vibración y de golpes ligeros, dependiendo de la frecuencia de la estimulación.

Igualmente se ha determinado que los mecanorreceptores tipo C en regiones restringidas de la piel (Kumazawa y Perl, 1977; Nordin, 1990), así como los receptores pacciniformes de las articulaciones y los receptores musculares (Willis y Coggeshall, 1991) pueden estar implicados en la percepción de sensaciones de cosquilleo, temblor y vibración.

Las aferencias primarias procedentes de los mecanorreceptores FAI, que evocan sensaciones de temblor, terminan en las capas profundas del asta dorsal (láminas III y IV) (Brown y cols., 1977) y algunas ascienden en el funículo dorsal hasta los núcleos de la columna dorsal (Willis y Coggeshall, 1991). Sin embargo, las aferencias procedentes de los receptores FAII terminan en las regiones mediales de las láminas III y VI (Brown y cols., 1980).

Las proyecciones ascendentes a través de las cuales los estímulos de temblor (mecanorreceptores FAI) son conducidos al complejo ventrobasal del tálamo, incluyen el sistema columna dorsal (aferencias primarias y no primarias)-lemnisco medial en el funículo dorsal, el sistema espino-cervicotalámico en el funículo dorsolateral (Brown y Franz, 1969; Bryan y cols., 1974; Willis y cols., 1974; Brown, 1981; Brown y Fyffe, 1981), y el tracto espinotalámico en el funículo ventrolateral (Willis y Coggeshall, 1991). Los estímulos vibratorios procedentes de los receptores de Paccini (FAII), (Brown y Fyffe, 1981; Brown y cols., 1983) alcanzan el complejo ventrobasal talámico únicamente a través del sistema columna dorsal (aferencias primarias y no primarias)lemnisco medial (Willis y Coggeshall, 1991).

#### Propiocepción

La propiocepción incluye el sentido de la posición y localización de los miembros en el espacio y su cinestesia, es decir, la noción de su movimiento. Dicha modalidad sensorial permite diferenciar entre movimientos activos y pasivos, así como la intensidad de la fuerza aplicada en las contracciones voluntarias (Googwin y cols., 1972; McCloskey, 1973; Horch y cols., 1975; Roland y Ladegaard-Pedersen, 1977). Existen ciertas controversias en la bibliografía acerca de los receptores encargados de la percepción de la posición (ver Willis y Coggeshall, 1991), aunque actualmente se acepta que los husos musculares están implicados en dicha función, así como en la distinción entre movimientos activos y pasivos. En mamíferos, los husos musculares poseen dos tipos de fibras aferentes: primaria y secundaria, que transmiten distintos parámetros de la contracción y el estiramiento muscular.

Los receptores de adaptación lenta de las articulaciones (terminales de Ruffini), responden ante estímulos extremos de flexión y extensión de las mismas, y están relacionados con la detección de los límites de sus movimientos, más que con la transmisión de información acerca de la posición de los miembros (Willis y Coggeshall, 1991). Los mecanorreceptores de adaptación lenta (SAII), además de estar implicados en la trasmisión de información tactil y de presión, responden a estímulos de estiramiento de la piel y participan en la evocación de sensaciones de posición y de movimiento (Macefield y cols., 1990; Willis y Coggeshall, 1991).

Por último, los órganos tendinosos de Golgi responden ante estímulos de estiramiento y contracción muscular proporcionalmente a la fuerza de la contracción (Willis y Coggeshall, 1991).

Las aferencias musculares propioceptivas terminan principalmente en la región profunda del asta dorsal, donde se han descrito neuronas que intervienen tanto en la elaboración de reflejos motores, como en procesos sensoriales propioceptivos (Willis y Coggeshall, 1991). Las aferencias musculares propioceptivas de tipo Ia, terminan en las láminas VI, VII y IX (Brown y Fyffe, 1978, Brown, 1981), y las de tipo II en las láminas IV, VII y IX (Fyffe, 1979; Brown, 1981). Las aferencias de tipo Ib, procedentes de los órganos tendinosos de Golgi, se arborizan en las láminas VI y VII. Otras aferencias musculares que parecen ser de naturaleza nociceptiva, concluyen en las láminas I y V, (Mense y Prabhakar, 1986), en la columna de Clarke y algunas ascienden en el funículo dorsal hasta el núcleo cuneatus externo (Craig y Mense, 1983; Nyberg y Blomqvist, 1984; Hongo y cols., 1987).

La información propioceptiva, procedente de los mecanorreceptores del músculo y de las articulaciones de las extremidades anteriores, asciende a través del fascículo *cuneatus* de la columna dorsal hasta el núcleo *cuneatus* (Willis y Coggeshall, 1991). Sin embargo, las aferencias propioceptivas de las extremidades posteriores (Lloyd y McIntyre, 1950; Burguess y Clark, 1969), ascienden a través del sistema espino-rombencefalico-talámico, primero en el funículo dorsolateral hasta el núcleo Z rombencefálico, el cual proyecta al tálamo contralateral (Willis y Coggeshall, 1991). Aunque no existen evidencias definitivas, se ha sugerido la participación del tracto espinotalámico en la propiocepción (Willis, 1974; Appelbaum y cols., 1975; Milne y cols, 1982).

#### Dolor

La percepción del dolor incluye dos tipos fundamentales de sensación de acuerdo con Lewis (1942): superficial y profundo. El dolor superficial es una sensación localizada, resultado de la estimulación de la piel, que puede ser dividido en primario y secundario, transmitidos a través de fibras A delta y C respectivamente, que evocan dos sensaciones dolorosas sucedidas en el tiempo, debido a la velocidad diferencial de transmisión de ambos tipos de fibras (Lewis y Pochin 1938; Lewis, 1942; Sinclair y Strokes, 1964; Price y cols., 1977; Campbell y LaMotte, 1983). El dolor profundo es una sensación con una localización imprecisa que se origina en tejidos subcutáneos como el músculo esquelético, tendones, periostio, y articulaciones. Un tercer tipo de dolor es el visceral, que comparte diversas características con el dolor profundo. Una de las consecuencias de las sensaciones dolorosas consiste en provocar respuestas en el sistema nervioso autónomo, cambios endocrinos y respuestas motivacionales y

afectivas (ver Willis y Coggeshall, 1991). Por último se ha descrito una modalidad de dolor neurogénico que se produce sin una necesaria activación de los nociceptores y está asiociado con lesiones del sistema nervioso central (ver Willis y Coggeshall, 1991).

Los nociceptores, receptores encargados de percibir los estímulos nociceptivos, consisten en fibras con terminaciones libres no encapsuladas, escasamente mielinizadas (A delta) o amielínicas (C), que transmiten el estímulo con una mayor o menor velocidad respectivamente (Burgess y Perl, 1967; Georgopoulos, 1976; Willer y Albe-Fessarrd, 1983). No existen diferencias estructurales, entre los que inervan distintos órganos o tejidos (piel, músculo, articulaciones o visceras). Se han desrito, sin embargo, varios tipos funcionales de nociceptotes, dependiendo de parámetros fisiológicos como el umbral de activación, o el estímulo o estímulos necesarios para su excitación. En la piel, por ejemplo, están presentes diversos nociceptores A delta que se activan ante estímulos mecánicos, de frio, o a ambos; nociceptores C que responden igualmente a estímulos de frio, mecánicos y térmicos, y receptores polimodales que se excitan frente a estimulación mecánica, térmica y química.

Las aferencias primarias de los nociceptores A delta terminan en las láminas I y V (Light y Perl, 1979), mientras que las procedentes de los nociceptores C se arborizan en las láminas I y II (Sugiura y cols., 1986). En las láminas I, II y V se han descrito neuronas que responden a estimulación nociceptiva, así como en los núcleos de la columna dorsal que proyectan contralateralmente al tálamo. La información nociceptiva que llega a los núcleos de la columna dorsal, asciende a través del sistema postsináptico de la columna dorsal (Angaut Petit 1975a,b; Giesler y Cliffer, 1985; Brown y cols., 1983; Bennett y cols., 1984; Kamogawa y Bennet, 1986) y en menor medida a través de las aferencias primarias.

En el funículo dorsolateral, algunas fibras de los tractos espinocervical (Brown y Franz, 1969; Bryan y cols., 1973, 1974; Cerveró y cols., 1977) y espinotalámico dorsal (Apkarian y Hodge a,b, 1989) responden a estimulación nociceptiva.

La mayoría de las neuronas de la proyección espinotalámica, en el funículo ventrolateral, responden a estimulación nociceptiva (Willis y cols., 1974; Price y cols., 1978; Chung y cols., 1979; Kenshalo y cols., 1979; Surmeier y cols., 1988) aunque en ocasiones integran los estímulos dolorosos con otros tipos de información (Willis y Coggeshall, 1991). Igualmente, se ha descrito la transmisión de estímulos nociceptivos a través de los tractos espinoreticular (Fields y cols., 1975, 1977b), espinomesencefálico (Menetrey y cols., 1980) y espinohipotalámico (Willis y Coggeshall, 1991), si bien dichas proyecciones podrían estar implicadas más en los componentes afectivos y motivacionales del dolor, que en el sensorial (Willis y Coggeshall, 1991).

#### Temperatura

Según los resultados de los trabajos de Hensel (1950, 1973a, 1974), las sensaciones térmicas de frio y de calor se producen ante la variación ascendente o descendente de la temperatura neutra de la piel, estimada en 32-37°C. Los extremos en la sensación térmica, tanto de frio (Wolf y Hardy, 1941) como de calor (Hardy y cols., 1951; Neisser, 1959; LaMotte y Campbell, 1978) evocan sensaciones dolorosas.

La sensación térmica depende de la activación de los receptores específicos de frio y de calor. Los primeros están innervados por fibras A delta y C, mientras que los de calor, únicamente reciben fibras C (ver Willis y Coggeshall, 1991) y responden a variaciones de temperatura de la piel, y a determinados agentes químicos (Hensel y Zotterman, 1951c; Dodt y cols., 1953; Hensel, 1973a) que provocan sensaciones térmicas. Las sensaciones de frio pueden además ser producidas por la disminución en la actividad de los receptores de calor, causada por un descenso de la temperatura, aunque no existen evidencias del fenómeno contrario. Las variaciones térmicas afectan a la actividad de algunos mecanorreceptores, si bien este fenómeno no parece inetervenir significativamente en el origen de sensaciones térmicas (Willis y Coggeshall., 1991).

Las aferencias cutáneas A delta y C, que probablemente incluyen las procedentes de receptores térmicos, terminan en las láminas I, II y V (Cerveró, 1989) en las que se han descrito neuronas que responden a la estimulación térmica (Willis y Coggeshall, 1991). Hasta el momento, únicamente se ha demostrado la implicación de las proyecciones espinotalámicas, originadas en neuronas de lámina I y que ascienden en el sistema ventrolateral, en la transmisión de información térmica a centros supraespinales (Craig y Kniffki, 1985; Ferrington y

cols., 1987, Kanui, 1988). Algunas de las neuronas que originan esta proyección, responden exclusivamente a información térmica (Craig y Kniffki, 1985), aunque en otras se produce una integración con otras estimulaciones generalmente nociceptivas (Craig y Kniffki, 1985; White y Sweet, 1969). Foerster y Gagel (1932) describieron que los axones de las neuronas espinotalámicas termorreceptivas parecen ascender en el funículo ventrolateral, segregados dorsalmente respecto a las proyecciones nociceptivas. En 1985 Craig y Kniffki demostraron que la proyección espinotalámica termoreceptiva alcanza la parte medial del complejo ventrobasal talámico, donde se han descrito neuronas que responden a variaciones térmicas de la piel de distintas regiones corporales (ver Willis y Coggeshall, 1991).

#### Sensaciones viscerales

Las percepciones viscerales incluyen la activación de receptores que transmiten la información necesaria para la regulación del medio interno, y no provocan sensaciones conscientes, como la información procedente de los barorreceptores arteriales, o de los quimiorreceptores de los cuerpo carotídeos (Cerveró y Foreman, 1990). Por el contrario, existen receptores que transmiten información visceral que sí evoca sensaciones conscientes. La sensación visceral mayoritaria es el dolor, si bien existen otras, como la de saciedad o el sentido de vaciamiento de los órganos (Leek, 1972). Aunque se ha descrito la existencia de termorreceptores abdominales (Riedel, 1976), estos deben intervenir en la termorregulación corporal, ya que aparentemente no se producen sensaciones térmicas viscerales.

La inervación sensorial de las vísceras es cuantitativamente menor que la de la piel (Cerveró y Foreman, 1990), sin embargo, presenta una dualidad ya que los órganos internos están inervados por el sistema nervioso simpático y el parasimpático (ver Cerveró y Foreman, 1990). En general la sensación visceral de dolor es percibida a través del sistema nervioso simpático el cual proyecta al sistema nervioso central a través de las raíces dorsales espinales. Por el contrario, la entrada de otros tipos de información viceral, requeridos para la elaboración de reflejos y para la regulación visceral, es percibida a través del sistema nervioso parasimpático, y alcanza el sistema nervioso central a través de los pares craneales VII, IX y X (Cerveró y Foreman, 1990).

El 90% de los receptores viscerales son fibras amielínicas, o fibras que pierden la mielina al penetrar en la víscera (Kuo y cols., 1982). Se han descrito nociceptores viscerales en el corazón, los pulmones, la vejiga urinaria, los conductos biliares, los testículos y en el útero (Cerveró, 1985). Igualmente se han observado receptores en el mesenterio y en la vaina serosa de diversos órganos, que se activan ante estímulos de movimiento y distensión de los mismos, y por lo tanto podrían contribuir a las sensaciones de saciedad y de dolor en aquellos casos de estiramientos excesivos (Willis y Coggeshall, 1991). Asimismo, se han descrito receptores en el músculo liso del tracto gastrointestinal y de la vejiga, que parecen intervenir en la sensación de vaciamiento de los distintos órganos, así como en el dolor (Leek, 1972), ya que

responden a estímulos de distensión y de contración (Winter, 1971, Leek, 1972).

Las aferencias viscerales primarias mielínicas terminan en las láminas I, II, V-VII y X espinales sacras. A nivel torácico terminan en las láminas I y V y con menor densidad en la parte externa de la lámina II. Sugiura y cols. (1989) han descrito que las aferencias de las fibras amielínicas terminan en las láminas I, II, IV-V y X (Cerveró y Foreman, 1990; Willis y Coggeshall, 1991). Algunas de las aferencias viscerales ascienden y descienden a niveles espinales adyacentes a través del tracto de Lissauer en el funículo dorsolateral (Cerveró y Foreman, 1990).

La información visceral asciende a niveles supraespinales por el funículo dorsal hasta los núcleos de la columna dorsal (Amassian, 1951; Aidar y cols., 1952; Perl y cols., 1962). Asimismo se ha descrito la existencia de fibras ascendentes viscerales en el funículo dorsolateral, procedentes de neuronas del asta dorsal superficial (Cerveró y Foreman, 1990). Sin embargo, las proyecciones espinoreticular y espinotalámica del sistema ventrolateral constituyen cuantitativamente los sustratos anatómicos más importantes, por los cuales asciende la información visceral (Cerveró y Foreman, 1990; Willis y Coggeshall, 1991). Se ha sugerido que las proyecciones espinosolitarias, que se originan en neuronas de las láminas I, V, X y en el núcleo espinal lateral (Menetrey y Basbaum, 1987; Leha y cols., 1988); y las proyecciones espinomesencefálicas que terminan en la sustancia gris central y en el área parabraquial, podrían también estar implicadas en procesos de integración viscerosomáticos y visceroviscerales (Cerveró y Foreman, 1990).

### **VERTEBRADOS NO MAMÍFEROS**

Dentro de los vertebrados no mamíferos el estudio del sistema nervioso de los anfibios resulta de gran interés desde el punto de vista evolutivo, por representar éstos un grupo taxonómico intermedio entre los vertebrados acuáticos y los terrestres.

Los anfibios surgieron en el periodo Devónico superior, a partir de los peces Crosopterigios, y sufrieron numerosas adaptaciones motivadas por la transición que experimentaron al cambiar del hábitat acuático al terrestre. Algunas de estas transformaciones se producen durante la ontogenia de los anfibios actuales, debido a que a lo largo de su desarrollo embrionario y larvario experimentan un cambio en el modo de vida, que en estadios tempranos se desarrolla estrictamente en el agua y pasa posteriormente a depender de forma relativa de los hábitats acuáticos.

Actualmente los anfibios están clasificados en tres órdenes: Anura, Urodela y Apoda. Cada uno de ellos incluye géneros con formas de vida que se desarrollan con un grado variable de dependencia del agua. Todos ellos presentan características comunes con los vertebrados acuáticos, así como con los terrestres.

Las transformaciones producidas a lo largo de la ontogenia de los anfibios, incluyen entre otras, la regresión de las branquias externas, la adquisición de pulmones saculares que permiten el aprovechamiento del oxígeno atmosférico, las variaciones tegumentarias que impiden parcialmente la desecación, el desarrollo de las extremidades y las variaciones en el aparato locomotor como los cambios del esqueleto y de las masas musculares necesarios para la adquisición de los distintos patrones de locomoción propios de cada grupo. De forma paralela a dichas transformaciones, los anfibios sufren variaciones en los sistemas sensoriales y en el sistema nervioso central. Se produce una regresión o involución del sistema de la línea lateral, presente en los anamniotas acuáticos, en los géneros cuyos individuos adultos abandonan el medio acuático (Wahnschafre y cols., 1987). Una transformación sorprendente durante la ontogenia de algunos géneros de anfibios es la producida al cambiar de un modelo de locomoción natatoria, mediante movimientos corporales cuvo control nervioso reside básicamente en la médula espinal (Hughes, 1957; Roberts y cols., 1983), a un modelo de locomoción tetrápodo, coordinado por un control nervioso supraespinal. Paralelamente se producen variaciones en los sistemas sensoriales espinales en los que el sistema de Rohon-Beard es progresivamente reemplazado por el desarrollo del sistema constituído por las células ganglionares de las raíces dorsales (Forehand y Farel, 1982; ten Donkelaar y de Boer-van Huizen, 1991).

### Receptores periféricos somatosensoriales en los anfibios

En los anfibios se han descrito distintos tipos de receptores cutáneos responsables de la recepción de estímulos somatosensoriales, que incluyen tacto, presión, temperatura y dolor. Se diferencian cuatro tipos de receptores cutáneos, atendiendo a diversos parámetros morfológicos y fisiológicos como el diámetro de la fibra, el modo de terminación libre o encapsulado, la distribución dérmica o epidérmica, el tipo de estímulo percibido, así como la velocidad de transmisión del estímulo y la adaptación rápida o lenta del receptor (Marushashi y cols., 1952; Lindblon, 1962; Spray, 1976). En general, los mecanorreceptores consisten en terminaciones relativamente gruesas, libres o encapsuladas que se distribuyen en la epidermis, o en la zona de transición con la dermis y tienen una alta velocidad de conducción. Los receptores que responden a estímulos térmicos, y los que lo hacen frente a estímulos nociceptivos corresponden a fibras de menor diámetro, de terminación libre, que generalmente se distribuyen en las regiones profundas de la dermis, con una menor velocidad de conducción que en el caso de los mecanorreceptores. Dentro de los mecanorreceptores, Catton (1958, 1976) diferenció entre los que responden a estímulos tactiles o a presión, así como cuatro tipos diferentes de mecanorreceptores en base a parámetros fisiológicos carentes de diferencias morfológicas aparentes.

Las aferencias propioceptivas musculares provienen en los anfibios de los husos musculares, órganos encapsulados que responden a estímulos provocados por variaciones de tensión y de estiramiento de su fibra muscular (Katz, 1961; Ottoson, 1976). Los husos musculares de la rana tienen un único tipo de terminal sensitivo, homólogo del primario de los mamíferos, consistente en una fibra mielínica que penetra en la cápsula y progresivamente va perdiendo la mielina al ramificarse sobre la fibra intrafusal (Barker, 1974; Ottoson, 1976).

#### La médula espinal de los anfibios

La médula espinal de los anuros únicamente posee 11 segmentos y presenta dos intumescencias, una cervical (o braquial) y otra lumbar que corresponden a los niveles de inervación de las extremidades anteriores y posteriores respectivamente. Después de la metamorfosis, se observan diez nervios espinales que se numeran del 2 al 11, excluyendo el primer nervio ya que está muy poco desarrollado (Gaupp, 1899; Hughes y Tschumi, 1958; Deuchar, 1975; Thors, 1980). Dicha médula espinal está conectada con cada uno de los diez de pares de nervios espinales por medio de una raíz dorsal y otra ventral. En el género Rana la raíz ventral del segundo nervio espinal constituye el nervio hipogloso que, en parte, se incorpora al plexo braquial. En Xenopus laevis la raíz ventral del segundo nervio espinal únicamente contribuye a la formación del plexo braquial (Deuchar, 1975), mientras que dicho nervio carece del ganglio de la raíz dorsal. Los nervios 3 y 4 forman el plexo braquial e inervan las extremidades anteriores, los nervios espinales 5, 6 y 7 inervan el tronco, mientras que los 8, 9 y 10 forman el plexo lumbosacro que inerva las extremidades posteriores. El polo caudal de la médula o filum terminale, se reduce en los anuros a un cono relativamente largo, compuesto por un canal central, rodeado por una población celular mayoritariamente glial, que representan los restos de la involución sufrida por esta región durante la metamorfosis, al desaparecer la musculatura de la cola a la cual inervaba. No parecen existir excesivas diferencias interespecíficas en anuros en cuanto a la organización básica de la médula espinal, excepto en la variación de su longitud (Ebbesson, 1976).

En los anfibios urodelos la médula espinal se extiende a lo largo de toda la longitud del canal vertebral. Como en el caso de los anuros, se distinguen dos intumescencias, cervical o braquial y lumbar, que representan las regiones de inervación de las extremidades. Nuestro conocimiento de la médula espinal de los urodelos es incompleto, debido a la escasez de trabajos experimentales de trazado neuronal, que se limitan a estudios sobre las proyecciones de las raíces dorsales, algunas vías espinales ascendentes, conexiones propioespinales, así como sobre el recorrido y el lugar de terminación de algunas vías supraespinales descendentes.

Las secciones transversales de la médula espinal en anuros, a diferencia de la de urodelos que presenta una morfología ovalada, muestran la típica forma de H de la sustancia gris, presente en la mayoría de los grupos de vertebrados terrestres. Las astas dorsales, relativamente pequeñas en anuros, están separadas mutuamente por los funículos dorsales. Las astas ventrales poseen grupos de motoneuronas organizadas de manera relativamente simple, (ver Cruce, 1974; Frank y Westerfield, 1982; Hulshof y cols., 1987) que dan origen a las raíces ventrales espinales.

#### - Citoarquitectura

Debido a la escasa diferenciación de sus diversos grupos celulares, la sustancia gris espinal de anfibios se ha subdividido, en base a los resultados obtenidos en estudios con técnicas de Golgi, en una serie de campos espinales, coincidiendo con los lugares de terminación de los sistemas de fibras aferentes (Ebbesson, 1976), que contrasta con la subdivisión en láminas, citoarquitectónicamente diferenciables, establecida en mamíferos (Rexed 1952). Ebbesson (1976) en los anuros Ranacatesbeiana y R. pipiens, ha definido los campos dorsal, lateral, central, ventrolateral y ventromedial, así como los campos motor lateral, que inerva la musculatura de las extremidades, y motor medial que inerva la musculatura axial. Esta subdivisión es extensible a otras especies de anuros como R. esculenta o Xenopus laevis (Nikundiwe y cols. 1982).

El campo dorsal en anuros ocupa la mayor parte del asta dorsal, y es comparable a las láminas I a IV de Rexed (1964) de mamíferos, aunque sus límites son difíciles de establecer (Ebbesson, 1976). No obstante, las fibras de las raíces dorsales forman un plexo terminal característico sobre la superficie del asta dorsal, que fue descrito por Silver (1942) como el neuropilo dorsal, formado principalmente por fibras aferentes de las raíces dorsales y, en los primeros segmentos espinales, por aferencias primarias del nervio trigémino (Fuller y Ebbesson, 1973; Matesz y Székely, 1978; González y cols., 1993). En el campo dorsal se han descrito distintos tipos neuronales con prolongaciones dirigidas al neuropilo dorsal, a los funículos dorsal y dorsolateral, así como al neuropilo lateral (Ebbesson, 1976).

El campo lateral recibe fibras de las raíces dorsales (neuropilo ventral), que determinan sus límites medial y ventral (Joseph y Whitlock 1968b; Ebbesson, 1976). Además le llegan dorsalmente fibras del tracto reticuloespinal (Mensah, 1974), así como fibras del tracto rubroespinal (Corvaja y Grofová, 1972; ten Donkelaar 1982, 1992; Larson-Prior y Cruce, 1992); y carece de aferencias trigeminales (Fuller y Ebbesson, 1973; Matesz y Székely, 1978; González y cols., 1993). En este campo espinal existen dos tipos celulares, en la región medial se han descrito neuronas pequeñas con dendritas cortas y restringidas a los límites del campo, mientras que en las regiones laterales existen neuronas de mayor tamaño con dendritas dirigidas hacia todos los funículos. Debido al patrón de sus aferencias, Ebbesson (1976) comparó el campo lateral de los anuros con las láminas V-VI de Rexed (1964) en mamíferos.

El campo motor lateral, presente en los niveles de las intumescencias espinales, está formado por motoneuronas que se organizan en un grupo o columna lateral, el cual inerva la musculatura de las extremidades. Mientras que el campo motor medial, observable en todos los niveles espinales, está constituído por una columna delgada de motoneuronas más mediales que inervan la musculatura axial (Cruce, 1974; Frank y Westerfield, 1982; Hulshof y cols., 1987).

Los campos ventrolateral y ventromedial se caracterizan por sus aferencias supraspinales (Corvaja y Grofová, 1972; Corvaja y cols., 1973; Fuller, 1974; Mensah, 1974; ten Donkelaar, 1982). Ebbesson (1976) comparó los campos ventrolateral y ventromedial con las láminas VII y VIII de Rexed (1964) respectivamente. El campo ventrolateral está probablemente inervado por fibras reticulospinales, pero no recibe aferencias vestibulospinales y tectospinales (Ebbesson, 1976). Las neuronas de dicho campo son pequeñas o medianas y existen al menos tres tipos de acuerdo con su morfología dendrítica (Ebbesson, 1976). El campo ventromedial es la zona de terminación prioritaria para las proyecciones tectoespinales (Rubinson, 1968) y vestibuloespinales (Corvaja y Grofová, 1972; Corvaja y cols., 1973; Fuller, 1974), si bien también culminan en él algunas fibras reticuloespinales (Mensah, 74). Sus neuronas poseen árboles dendríticos reducidos y dirigidos bilateralmente al funículo ventral (Ebbesson, 1976).

El campo central, localizado alrededor el canal central, es comparable a la lámina X de Rexed (1964), y está compuesto por dos tipos neuronales (Ebbesson, 1976) según dirijan sus dendritas a todos los campos adyacentes o a los campos central y dorsal.

En urodelos los datos existentes sobre la organización citoarquitectónica de la médula espinal, se basan únicamente en la escasa información resultante de estudios pioneros (Burckhardt, 1889; Studnicka, 1895; Van Gehuchten, 1897; Nieuwenhuys, 1964). Las astas dorsales están formadas fundamentalmente por células pequeñas que constituyen la sustancia gelatinosa (Nieuwenhuys, 1964), aunque Burckhardt (1889) y Studnicka (1895) describieron algunas neuronas más grandes. La zona intermedia de la sustancia gris, situada ventralmente a la sustancia gelatinosa y dorsalmente a las motoneuronas, está formada por células pequeñas y medianas que extienden sus dendritas casi exclusivamente a la sustancia blanca (Van Gehuchten, 1897). Las astas ventrales son una masa compacta que contiene numerosas neuronas grandes que proyectan ventrolateralmente (Van Gehuchten, 1897).

- Patrón de terminación espinal de las proyecciones de las raíces dorsales

Estudios basados en técnicas de degeneración anterógrada (Liu y Chambers 1957; Joseph y Whitlock, 1968a,b; Liu, 1969) sugirieron que las fibras de las raíces dorsales terminan fundamentalmente en el asta dorsal, y no en el asta ventral. Igualmente se demostró que ascienden hasta niveles del tronco cerebral, y que poseen ramas descendentes que se extienden hasta cinco y seis segmentos caudalmente al nivel de la raíz lesionada. Joseph y Whitlock (1968a,b) propusieron que las neuronas del asta ventral, probablemente recibían proyecciones de fibras de las raíces dorsales a través de sus dendritas que invaden el asta dorsal. Sin embargo, la aplicación de técnicas de marcaje con cobalto en Rana esculenta (Székely, 1976; Antal y cols., 1980; Székely y cols., 1982; Székely y Antal, 1984) y con HRP en Rana catesbeiana (Frank y Westerfield, 1982; Jhaveri y Frank, 1983; Liuzzi y cols., 1984; Lichtman y cols., 1984; Smith y Frank, 1988a), Rana pipiens (Rosenthal y Cruce, 1985), Rana ridibunda (Motorina y cols., 1982a,b; Grantyn y cols. 1982, 1984a,b; Shapovalov y Shiriaev 1984a) y Xenopus laevis (Nikundiwe y cols., 1982; Shiriaev y Shupliakov, 1986) ha proporcionado más información sobre la organización de las aferencias primarias espinales en los anfibios anuros.

Las fibras primarias, al entrar en la médula espinal de anuros, se distribuyen segregándose en un *componente medial* de fibras de mayor diámetro, que ingresa en el funículo dorsal, para ascender hasta el núcleo de la columna dorsal y otras estructuras del tronco cerebral; y un *componente lateral* formado por axones de menor diámetro que se agrupan ventral y lateralmente a la entrada de la raíz para formar el tracto

de Lissauer, situado en el funículo dorsolateral (Székely, 1976; Rosenthal y Cruce, 1985), el cual se condidera implicado en la transmisión de información nociceptiva (Maruhashi y cols., 1952). Dichas aferencias primarias ocupan zonas espinales concretas de terminación dependiendo del origen y modalidad axonal (ver Székely y cols., 1982; Jhaveri y Frank, 1983; Grantyn y cols., 1984a,b; Székely y Antal, 1984). Posteriormente a su entrada en la médula espinal, a través de las distintas raíces, las fibras primarias del componente medial se bifurcan para dar ramas ascendentes y descendentes que discurren a través del funículo dorsal. A intervalos a lo largo de su recorrido espinal, las fibras emiten colaterales que se arborizan en la sustancia gris espinal, formando fundamentalmente dos neuropilos o campos de terminación: el neuropilo dorsal localizado en el asta dorsal, y el neuropilo ventral que ocupa el campo lateral (Székely, 1976; Frank y Westerfield, 1982; Jhaveri y Frank, 1983). Los axones procedentes de aferencias primarias cutáneas, proyectan únicamente al ncuropilo dorsal (Jhaveri y Frank, 1983; Székely y Antal, 1984), mientras que la gran mayoría de las aferencias sensoriales procedentes del músculo, proyectan al neuropilo ventral, en el que se distribuyen igualmente las dendritas de motoneuronas y de interneuronas.

En segmentos cervicales son relativamente pocos los axones que profundizan ventral y lateralmente para alcanzar los somas de las motoneuronas (Frank y Westerfield, 1982; Jhaveri y Frank 1983; Lichtman y cols. ,1984; Székely y Antal, 1984). Se ha demostrado en estudios fisiológicos (Frank y Westerfield, 1982; Lichtman y Frank, 1984), que las motoneuronas del triceps braquial reciben una activación monosináptica, desde las fibras primarias con información del propio músculo. Estos datos, conjuntamente con la escasez de contactos axosomáticos demostrados morfológicamente, indican que la mayoría de las conexiones monosinápticas en la médula espinal a nivel braquial, debe producirse a través de contactos axodendríticos en el neuropilo ventral.

En segmentos torácicos el neuropilo dorsal está muy desarrollado, mientras que el ventral contiene menor número de fibras (Székely, 1976; Jhaveri y Frank, 1983; Smith y Frank, 1988a,b). Esto coincide con los experimentos mediante estimulación de las raíces dorsales en *Rana ridibunda*, (Shapovalov y Shiriaev, 1984) y en *Rana pipiens* (Carlsen y Mendell, 1977) que demostraron que las interacciones sinápticas con las motoneuronas torácicas están mediadas por interneuronas, y no se producen sinapsis sensorimotoras directas.

En segmentos lumbares las aferencias primarias establecen diversas sinapsis de paso sobre las dendritas dorsales de las motoneuronas de las regiones dorsal (Corvaja y Pellegrini, 1975; Sotelo y Grofová, 1976; Adanina y Shapovalov, 1983) y dorsolateral (Székely, 1976; Liuzzi y cols., 1984) del campo motor lateral.

En urodelos las aferencias de las raíces dorsales presentan menor diferenciación en componentes *medial* y *lateral*. La mayoría de las fibras de las raíces dorsales se bifurcan, inmediatamente después de su entrada en la médula, en ramas ascendentes y descendentes (Ramón y Cajal, 1909; Roth y Wake, 1985). Las ramas ascendentes y

mielinizadas ocupan la mayoría del funículo dorsal (Ramón y Cajal, 1909; Herrick, 1948). Mientras que la división lateral, que incluye fibras mielínicas y amielínicas, forma un fascículo comparable al tracto de Lissauer de otros vertebrados (Ramón y Cajal, 1909; Ariëns Kappers y cols., 1936; Roth y Wake, 1985). En preparaciones basadas en técnicas de Golgi, Ramón y Cajal (1909) observó en Pleurodeles waltl que las proyecciones de las raíces dorsales en la médula de urodelos son, aunque mucho más simples. similares a las de los anuros, y en ocasiones alcanzan posiciones próximas a las motoneuronas. En urodelos, a excepción de lo anteriormente descrito, apenas existen datos experimentales disponibles sobre las proyecciones de las fibras de las raíces dorsales. Sin embargo, en un estudio más reciente (Holder y cols., 1991) en Ambystoma mexicanum se demostró, que las aferencias primarias espinales se entremezclan con las dendritas dorsales de las motoneuronas que inervan las extremidades, pero no con las que inervan la musculatura axial.

#### Provecciones espinales ascendentes

Estudios basados en técnicas de degeneración anterógrada (Kuru, 1956; Ebbesson, 1969, 1976, en *Rana catesbeiana*; Hayle, 1973a,b, en *Rana. temporaria*) han demostrado la existencia en anuros de dos sistemas de proyecciones espinales ascendentes: 1) sistema de aferencias *primarias* que asciende a través del funículo dorsal, hasta el núcleo de la columna dorsal, y 2) sistema de aferencias *secundarias* que asciende a través del funículo lateral, que fue descrita como el lemnisco espinal, y alcanza la formación reticular, y el techo mesencefálico. Mediante la utilización de marcaje con cobalto (Antal y cols., 1980; Urbán y Székely, 1982, Rana. esculenta) y de trazado neuronal con HRP (Nikundiwe y cols., 1982, *Xenopus laevis*), se ha demostrado la existencia de un sistema bien desarrollado de aferencias primarias espinales, ordenadas somatotópicamente, que ascienden, en el funículo dorsal hasta el núcleo de la columna dorsal. Esta proyección continúa rostralmente para inervar el complejo vestibular, así como la capa granular del cerebelo (tracto espinocerebeloso dorsal de Ebbesson) (Antal y cols., 1980; Székely y cols., 1980). Dichas fibras se originan en los ganglios espinales que inervan las extremidades (Antal y cols., 1980; González y cols., 1984).

#### Funículo dorsal:

 Sistema columna dorsal-lemnisco medial y otras proyecciones incluídas en el funículo dorsal.

La existencia del sistema columna dorsallemnisco medial es una característica común en el cerebro de vertebrados amniotas. Tanto en aves (Sinn, 1913; Craigie, 1928; Wild, 1985, 1989), como en reptiles (Kruger y Witkovsky, 1961; Goldby y Robinson, 1962; Ebbesson, 1967, 1969, 1978; Joseph y Withlock, 1968b, Künzle y Woodson, 1983; Pritz, 1983; Belekhova y cols., 1985; Pritz y Stritzel, 1986, 1989; Siemen y Künzle, 1994a,b), existe un patrón común de organización en cuanto a las aferencias primarias y no primarias (Funke, 1988 en aves; Pritz and Stritzel, 1994 en reptiles) de los núcleos de la columna dorsal, así como en sus proyecciones eferentes rombencefálicas,

mesencefálicas y talámicas. En anfibios exiten evidencias de una organización similar del sistema columna dorsal-lemnisco medial, lo que ha hecho sugerir a Willis y Coggeshall (1991) que su presencia es un caracter propio de vertebrados terrestres, al no haber sido descrito en los distintos grupos de peces (Zeehandelaar, 1921; Hayle, 1973), los cuales perciben gran parte de la información del medio que les rodea por el sistema de la línea lateral. No obstante, en diversas especies tanto en agnatos, petromizóntidos y mixinoideos (Northcutt y Ebbesson, 1980; Ronan y Northcutt, 1990) como en condrictios (Ebbesson y Hodde, 1981) y en osteictios (Finger, 1978; Oka et al., 1986; ver Ronan y Northcutt, 1990) se han descrito, en la región alar a nivel del óbex, terminales de proyecciones espinales que ascienden en el funículo dorsal. Sin embargo, no se han realizado hasta el momento estudios detallados de trazado neuronal anterógrado sobre las aferencias primarias espinales o de trazado retrógrado, con aplicaciones en el mesencéfalo y en el tálamo, que permitan establecer definitivamente la presencia del sistema columna dorsal-lemnisco medial en dichos vertebrados. Algunas especies de anfibios, en los que se ha sugerido la presencia del citado sistema, presentan modos de vida estrictamente acuáticos y mantienen el sistema de la línea lateral en estado adulto. Los estudios realizados hasta el momento en anfibios están mayoritariamente basados en técnicas anatómicas clásicas y degenerativas, lo que hace necesaria la aplicación de técnicas más recientes de trazado neuronal así como inmunocitoquímicas, con objeto de estudiar la organización detallada del sistema y establecer posibles homologías con otros grupos de vertebrados.

Tanto en agnatos (Northcutt y Ebbesson, 1980; Ronan y Northcutt 1990) como en condrictios (Ebbesson y Hodde, 1981), teleósteos (Ronan y Northcutt 1990) y reptiles (Ebbesson, 1967,1969; Jacobs, 1968; Pedersen, 1973; Künzle, 1982, Pritz y Stritzel, 1994), se ha descrito un componente de aferencias espinales primarias que asciende en el funículo dorsal, sobrepasa rostralmente los núcleos de la columna dorsal y llega hasta el cerebelo.

En anfibios anuros no se ha descrito el núcleo de columna dorsal (nucleus funiculis dorsalis) como entidad citoarquitectónica (Ariëns Kappers y Hammer, 1918; Zeehandelaar, 1921; Opdam y cols., 1976). Asimismo, Ariëns Kappers y cols. (1936), negaron la presencia en anfibios del sistema columna dorsal-lemnisco medial. Sin embargo, Woodburne (1939), en ránidos, mediante lesiones de los ganglios de las raíces lumbares, presentó evidencias de la presencia del núcleo de la columna dorsal, ya que observó la degeneración producida en una pequeña región rombencefálica situada dorsomedialmente a nivel del óbex, que definió como un primordio del núcleo gracilis. En trabajos basados en técnicas de degeneración anterógrada y de trazado neuronal con leucina tritiada (Liu y Chambers, 1957; Joseph y Whitlock, 1968; Ebbesson, 1969; Hayle, 1973b), se ha descrito el núcleo de la columna dorsal de los anuros como el lugar de terminación de fibras del funículo dorsal, pero sin considerarlo como una entidad citoarquitectónica. Sin embargo, Fernández de Molina y cols. (1966) y Silvey y cols. (1974) determinaron la presencia de un pequeño grupo de neuronas bien definido, ubicado en el polo rostral de la columna dorsal que responde a estimulación somatosensorial. La mayor parte de las aferencias

primarias espinales terminan en la región del núcleo de la columna dorsal, si bien un pequeño componente, al menos en el caso de las aferencias braquiales, continúa rostralmente a través del rombencéfalo para terminar en el cerebelo (Joseph y Witlock, 1968; Rushmer, 1970; Rushmer y Woodward, 1971; Székely y cols., 1980; Antal y cols., 1980; Nikundiwe y cols., 1982).

La existencia del lemnisco medial en anuros ha sido motivo de controversia hasta el principio de la década de los 80. Diversos trabajos (Vesselkin y cols., 1971; Vesselkin y Kovacevic, 1973; Silvey y cols., 1974; Neary y Wilczynski, 1977) propusieron la presencia de una proyección contralateral desde núcleo de la columna dorsal que concluía en el tálamo. Estudios más recientes de marcaje con cobalto en *Rana esculenta* (Antal y cols., 1980; Urbán y Székely, 1982), de trazado con HRP en *Xenopus laevis* (Nikundiwe y cols., 1982), o electrofisiológicos (Fernández de Molina. y cols., 1966; Urbán y Székely, 1982), han sugerido que el sistema columna dorsal-lemnisco medial en anfibios se asemeja al que existe en amniotas.

Mediante la utilización de técnicas electrofisiológicas, en experimentos con estimulación de la segunda raíz dorsal, del funículo dorsal o del núcleo de la columna dorsal, Urbán y Székely (1982) demostraron en *Rana esculenta* una proyección contralateral que alcanza el núcleo posterocentral talámico. Igualmente, los resultados procedentes de experimentos de trazado retrógrado han confirmado las proyecciones del lemnisco medial al tálamo en *Rana catesbeiana* (Neary y Wilczynski, 1977) y al torus semicircularis (Neary y Wilczynski, 1986; Neary, 1988).

En urodelos el núcleo de la columna dorsal (núcleo funiculi dorsalis) no ha sido descrito como entidad citoarquitectónica en el tronco cerebral (Herrick 1930, 1948; Kreht 1940a; Opdam y Nieuwenhuys, 1976), aunque, en experimentos de degeneración anterógrada, Nieuwenhuys y Cornelisz (1971) demostraron la presencia de una proyección organizada somatotópicamente, formada por las aferencias primarias espinales, que termina en la región caudal rombencefálica a nivel del óbex. Herrick (1948) postuló igualmente la terminación de fibras del funículo dorsal en dicha región, e indicó la posibilidad de que todas las fibras sensitivas somáticas que entran en el rombencéfalo, podrían terminar en un neuropilo común localizado a este nivel, en el que se combinan todas las modalidades sensoriales. Herrick (1948) sugirió que las fibras secundarias que se originan en dicha región rombencefálica no son estrictamente equipotenciales, y propuso una incipiente segregación de distintas funciones entre ellas. Del mismo modo, discutió la presencia del "lemnisco bulbar general" como un tracto mixto que asciende en la sustancia blanca ventrolateral del rombencéfalo, que recibe fibras contralaterales desde los núcleos del funículo dorsal y espinal del trigémino, y emite colaterales que alcanzan la formación reticular (Herrick, 1948; Herrick y Bishop, 1957). De acuerdo con estos trabajos, las fibras del lemnisco bulbar general se tuercen dorsalmente en la región del istmo y cruzan el mesencéfalo, en posición inmediatamente ventrolateral a las del lemnisco espinal, para terminar en el techo mesencefálico. Igualmente, postuló que algunas de ellas alcanzan la misma región talámica

que las del lemnisco espinal. Sin embargo, hasta el momento no existen apenas datos experimentales, disponibles en la bibliografía, basados en técnicas más modernas de trazado neuronal, sobre la presencia del lemnisco medial en urodelos, si bien, Wicht y Himstedt (1988) han descrito proyecciones rombencefalicotalámicas en *Triturus alpestris*, que alcanzan el tálamo dorsal.

La existencia del sistema columna dorsallemnisco medial, tanto en anuros como en urodelos, se ha descrito en base al patrón de terminación de las aferencias primarias espinales, así como en observaciones aisladas basadas en técnicas de trazado neuronal. En un estudio reciente con HRP sobre el desarrollo embrionario y larvario de las proyecciones ascendentes espinales en Xenopus laevis (ten Donkelaar y de Boer-van Huizen, 1991), se ha sugerido la existencia del sistema postináptico de la columna dorsal, si bien no se ha detallado, la localización de las neuronas que lo originan, el recorrido funicular de sus axones, ni su terminación en el DCN. Hasta el momento no se conoce con detalle la morfología o características de las células que dan origen al lemnisco medial ni sus principales destinos rombencefálicos, mesencefálicos y diencefálicos. Este hecho es consecuencia de la escasa diferenciación citoarquitectónica de los componentes de este sistema, que ha dificultado su estudio, así como por la falta de trabajos basados en técnicas de trazado e inmunohistoquímicas, que permitirian caracterizar su organización en los anfibios.

 Sistema espino-cervico-talámico y otras proyecciones incluídas en el funículo dorsolateral.

Hasta el momento no se ha descrito la existencia de los sistemas espino-cervico-talámico y espino-rombencefalico-talámico en vertebrados no mamíferos debido al escaso número de trabajos de investigación realizados sobre el tema. Sin embargo, existen algunos datos aislados en la bibliografía que apuntan la existencia de los tractos espinocervical y cervicotalámico, así como del núcleo cervical lateral (Herrick, 1948; van den Akker, 1970; Ebbesson, 1967; Finger, 1981; Forehand y Farel, 1982; Ito y cols., 1986; Necker, 1989; Ronan y Northcutt, 1990), con un modelo de organización similar al descrito en mamíferos.

Igualmente se han estudiado, en diversos vertebrados no mamíferos, las proyecciones espinales que ascienden en el funículo dorsolateral hasta el cerebelo, centros rombencefálicos como el núcleo del tracto solitario, núcleo del tracto descendente del trigémino, formación reticular lateral, y la región perifacial, y niveles de transición con el mesencéfalo como la región parabraquial (Pearson, 1936; Ebbesson, 1966, 1967, 1969; Karten, 1967; Jacobs, 1968; Pedersen 1973; Ebbesson y Hodde, 1981; Finger 1981; Funke y Necker, 1986; Funke, 1988), lo que sugiere una organización similar a la presente en mamíferos.

#### Funículo dorsolateral:

### Funículos ventral ventrolateral (cuadrante ventral):

y

#### - Sistema ventrolateral.

La disposición de proyecciones espinales que ascienden a través del cuadrante ventrolateral, para alcanzar diversas regiones rombencefálicas, mesencefálicas y el cerebelo, es un caracter común en el cerebro de todas las clases de vertebrados. Su presencia se ha descrito además de en los mamíferos (ver Willis y Coggeshall, 1991), en ciclóstomos (Tretjakoff, 1909; Pearson, 1936; Heier, 1948; Larsell, 1967; Northcutt y Ebbesson, 1980; Ronan y Northcutt, 1990), elasmobranquios (Ebbesson, 1972; Hayle, 1973; Ebbesson y Hodde, 1981; Smeets, 1982, Smmets y cols., 1983), teleósteos (Pearson, 1936; Hayle, 1973; Finger, 1978; Murakami y Ito, 1985; Ronan y Northcutt, 1990), anfibios (ver más adelante), reptiles (Huber y Grosby, 1926; Goldby y Robinson, 1967, 1969; Ebbesson, 1967; Pedersen, 1973; Northcutt y Pritz, 1978; ten Donkelaar y De Boer-van Huizen, 1978; Kusuma y cols., 1979; Ebbesson y Goodman, 1981; Künzle y Woodson, 1982; Pritz y Striztel, 1989) y aves (Karten, 1963; Karten y Rezvin, 1966; van den Akker, 1970; Vielvoye, 1977; Wild, 1983; Funke y Necker, 1986, Necker, 1989; 1990; Scheneider y Necker, 1989), en trabajos basados en técnicas degenerativas y de trazado neuronal.

La existencia de proyecciones espinotalámicas ha sido considerada como un caracter propio de vertebrados amniotas (ver Kevetter y Willis, 1984), debido a que su presencia fue únicamente descrita, además de en mamíferos, en el cerebro de aves (Karten, 1963; Karten y Rezvin, 1966, Wild, 1983; Necker, 1989; Scheneider y Necker, 1989, Funke, 1989b), y reptiles (Huber y Grosby, 1926; Ebbesson, 1967; Riss y cols., 1972, Kusuma, 1979; Pritz y Northcutt, 1980; Ebbesson y Goodman, 1981; Hoogland, 1981; Künzle y Woodson, 1982). En anamniotas, tanto en agnatos (Northcutt y Ebbesson, 1981, Ronan y Northcutt, 1990) como en elasmobranquios (Hayle, 1973) y teleósteos (Hayle, 1973), no se había descrito su existencia a excepción de en un trabajo en teleósteos, basado en técnicas anatómicas clásicas (Burr, 1928), y un estudio con técnicas de degeneración sobre el tiburón nodriza (Ebbesson y Hodde, 1981), Sin embargo, mediante trazado con HRP se ha confirmado la existencia de proyecciones espinotalámicas en teleósteos, que terminan mayoritariamente en el núcleo ventromedial (Murakami e Ito, 1985; Ito y cols., 1986). En anfibios, aunque su presencia fue sugerida por Herrick (1939), hasta el momento no existen evidencias que confirmen dicha proyección, si bien, no se han realizado hasta el momento experimentos con técnicas de trazado anterógrado.

En anfibios anuros (*Rana catesbeiana*) Ebbesson (1969, 1976) con técnicas de degeneración mediante hemisecciones espinales, describió que las fibras que ascienden a través del funículo lateral se agrupan fundamentalmente en dos tractos de fibras: un tracto medial, situado en los funículos ventral y ventrolateral que termina en niveles mesencefálicos; y un tracto lateral que se localiza en el funículo dorsolateral, en posición inmediatamente ventral al tracto descendente del trigémino. Éste último tracto asciende hasta el mesencéfalo, y en los casos en los que realizó lesiones inmediatemente caudales al óbex, hasta el diencéfalo caudal. La mayoría de las proyecciones de ambos tractos terminan en la formación reticular del tronco cerebral, mayoritariamente en niveles caudales, de forma que el medial proyecta a los núcleos reticulares inferior y medio, mientras que el lateral alcanza el núcleo reticular superior, localizado en niveles más rostrales (Ebbesson, 1976). Además, a lo largo de su recorrido a través del tronco cerebral, el lemnisco espinal envía fibras al núcleo motor dorsal del vago, así como al núcleo del tracto solitario, núcleo motor del facial, el posible equivalente en anuros de la oliva inferior, el cerebelo, y de manera más dispersa a la sustancia gris periventricular en niveles del istmo. En el mesencéfalo, dicho autor únicamente describió una proyección poco desarrollada que termina en una zona descrita por Potter (1965a) como los núcleos laminar y magnocelular del torus semicircularis. Igualmente sugirió la existencia de una proyección espinotalámica directa, en aquellos experimentos en que las hemisecciones espinales fueron inmediatamente caudales al óbex, en los que el lemnisco medial podría estar implicado. Dicha proyección no fue observada en los casos con lesiones en el segundo segmento espinal o en niveles más caudales.

En anuros, únicamente existen estudios de las proyecciones ascendetes espinales basados en técnicas degenerativas (Ebbesson, 1969; Hayle, 1973a). La ausencia de proyecciones espinotalámicas directas en los citados trabajos, contrasta con las respuestas electrofisiológicas descritas en el tálamo, y basadas en experimentos con estimulación somatosensorial en el nervio ciático (Vesselkin y cols., 1971). Igualmente, resulta llamativo el patrón restringido de fibras espinomesencefálicas (Ebbesson, 1969; 1976) frente al mapa somatosensorial de representación de la superficie corporal presente en el torus semicircularis (Comer y Grobstein, 1981). La escasa diferenciación de las proyecciones espinotalámicas y espinomesencefálicas descritas puede ser debida a las limitaciones en la sensibilidad y resolución de las técnicas degenerativas. La utilización de técnicas de trazado retrógrado con HRP en experimentos con aplicaciones en el tálamo (Neary y Wilczynski, 1977) y en el torus semicircularis (Wilczynski, 1981, Wilczynski y Neary, 1986), no han permitido identificar la población neuronal que pudiera originar las proyecciones espinales ascendentes. Resulta necesario, por lo tanto, la utilización de técnicas más sensibles de trazado neuronal bidireccional con los nuevos y más sensibles trazadores Leucoaglutinina de Phaseolus vulgaris o dextrano aminas, con objeto de estudiar detalladamente las posibles homologías y diferencias existentes entre las proyecciones espinales ascendentes de anfibios y las de otros vertebrados.

En los urodelos Ambystoma tigrinum y Ambystoma mexicanum, Herrick (1914, 1939a, 1948) y Herrick y Bishop (1957) observaron un sistema que denominaron lemnisco espinal, compuesto por fibras de segundo orden que se originan en neuronas de la sustancia gris espinal y en segmentos inferiores rombencefálicos, y que ascienden hasta niveles cerebrales más rostrales. De acuerdo con la descripción de Herrick (1914, 1939a, 1948), dicho sistema asciende a lo largo del rombencéfalo, en posición inmediatamente ventral al tracto decendente del trigémino, asociado al tracto espinocerebelar, dando muchos fibras terminales en la formación reticular rombencefálica (Herrick 1914) y en el cerebelo. En niveles más rostrales, la mayoría de las fibras terminan en el techo mesencefálico, aunque propone la posibilidad de que algunas de ellas continuen rostralmente para terminar en el tálamo dorsal (Herrick 1939b, 1948).

En un estudio posterior basado en técnicas de degeneración anterógrada en ajolotes juveniles, (Ambystoma mexicanum), mediante hemisecciones en diversos niveles espinales, y seciones de raíces dorsales braquiales y lumbares, Nieuwenhuys y Cornelisz (1971) describieron un tracto que recibe fibras desde todos los niveles espinales, y asciende por la parte dorsal del funículo lateral. Este sistema, interpretado como el lemnisco espinal de Herrick, emite fibras en el rombencéfalo y en el cerebelo, terminando principalmente en el techo mesencefálico (Nieuwenhuys y Cornelisz, 1971), si bien la presencia de una proyección espinotalámica no pudo demostrarse. Los citados autores describieron el sistema anterolateral formado por fibras que se originan en su mayor parte en niveles cervicobraquiales, y ascienden, a través del funículo ventrolateral, hacia el tronco cerebral para terminar en la placa basal del rombencéfalo, aunque un pequeño componente alcanza el tegmento mesencefálico. Este sistema comprende además un conjunto de fibras que rodea a las del lemnisco espinal y termina dorsalmente en el área de transición entre cerebelo y el techo mesencefálico (Nieuwenhuys y Cornelisz, 1971).

Como en el caso de los anuros, en urodelos no se han realizado, hasta el momento, trabajos de trazado anterógrado sobre las proyecciones ascendentes espinales, y en el único estudio de trazado retrógrado efectuado con aplicaciones talámicas de HRP (Wicht y Himstedt, 1988) no se ha logrado confirmar las sugerencias de Herrick en cuanto a la existencia de la proyección espinotalámica.
#### BIBLIOGRAFÍA

Adanina, A.A. and A.I. Shapovalov (1983) Mixed synapses of prymary afferents on frog motoneurons identified with neuroplasmic transport of HRP and electron microscopy. Dokl. Akan. Nauk. USSR. 269: 231-232.

Adrian, E.D. (1946). The Physical Background of Percepcion. In Oxford Univ. Press. Oxford:

Aidar, O., W.A. Geohegan and L.H. Ungewitter (1952) Splanchnic afferent pathways in the central nervous system. J. Neurophysiol. 15: 131-138.

Amassian, V.E. (1951) Fiber groups and spinal pathways of cortically represented visceral afferents. J. Neurophysiol. 14: 445-460.

Anderson, F.D. and C.M. Berry (1959) Degeneration studies of long ascending fiber systems in the cat brain stem. J. Comp. Neurol. 111: 195-229.

Andres, K.H. (1961) Untersuchungen uben den Feinbau von spinalganglien. Zeit. f. Zelforsch. 55: 1-48.

Angaut-Petit, D. (1975a) The dorsal column system: I Existence of long ascending postsynaptic fibers in the cat's fasciculus gracilis. Exp. Brain Res. 22: 457-470.

Angaut-Petit, D. (1975b) The dorsal column system: II Functional properties and bulbar relay of the postsynaptic fibers of the cat's fasciculus gracilis. Exp. Brain Res. 22: 471-493.

Antal, M., I. Tornai and G. Székely (1980) Longitudinal extent of dorsal root fibers in the spinal cord and brain stem of the frog. Neuroscience 5: 1311-1322.

Apkarian, A.V. and C.J. Hodge (1989a) A dorsolateral spinothalamic tract in macaque monkey. Pain 37: 323-333.

Apkarian, A.V. and H.C. Hodge (1989b) Primate spinothalamic pathway: II The cells of origin of the dorsolateral and ventral spinothalamic pathways. J. Comp. Neurol 288: 474-492.

Applebaum, A.E., J.E. Beall, R.D. Foreman and W.D. Willis (1975) Organization and receptive fields of primate spinothalamic tract neurons. J. Neurophysiol 38: 572-586.

Ariëns Kappers, C.U. and E. Hammer (1918) Das zentralnervensyten des Ochsenfrosches (*Rana catesbeiana*). Phychiat. Neurol. BI, Amsterdm 22 368-416.

Ariëns Kappers, C.U., G.C. Huber and E.C. Crosby (1936) The Comparative Anatomy of the Nervous System of Vertebrates, Including Man. Hafner, New York.

Aristoteles (322 a. de C.). Historia de los animales. In Akal (Ed.), Madrid; 1990.

Baker, M.L. and G.J. Giesler (1984) Anatomical studies of the spinocervical tract of the rat. Somatosensory Res. 2: 1-18.

Barker, D. (1974) The morphology of muscle receptors. In C. C. Hunt (eds): Handbook of sensory physiology. Berlin-Heidelberg New York: pp. 1-190.

Belekhova, M.G., V.D. Zharskaja, A.S. Khachunts, G.V. Gaidaenko and N.L. Tumanova (1985) Connections of the mesencephalic, thalamic and telencephalic auditory centers in turtles. Some structural bases for audiosomatic interrelations. J. Hirnforsch. 26: 127-152.

Bell (1811). Idea of a new anatomy of the brain. In Strahan and Preston (Ed.), Lodres; 1974.:

Bennett, G.J., N. Nishikawa, G.W. Lu, M.J. Hoffert and R. Dubner (1984) The morphology of dorsal column postsynaptic (DCPS) spino-medullary neurons in the cat. J. Comp. Neurol. 224: 568-578.

Berkley, K.J., R.J. Budell, A. Blomqvist and M. Bull (1986) Output systems of the dorsal column nuclei in the cat. Brain Res. Rev. 11: 199-225.

Bing, Z., L. Villanueva and D. Le Bars (1990) Ascending pathways in the spinal cord involved in the activation of subnucleus reticularis dorsalis neurons in the medulla of the rat. J. Neurophysiol. 63: 424-438.

Blix, M. (1884) Esperimentelle Beitraege zur losung der frage die specifische energie der hautnerven Z. Biol. 20: 141-156.

Boivie, J. and E.R. Perl (1975) Neural substrates of somatic sensation. MTP international Rew. of Science 3: 303-411.

Boivie, J. (1983) Anatomic and physiologyc features of the spino-cervico-thalamic pathway. In R. A. Macchi G Spresfico R (eds): Somatosensory integration in the thalamus. Elsevier, Amsterdam: pp. 63-106.

Boring, E.G. (1942). Sensation and perception in the history of experimental psychology. In Nueva York: Appleton-Century-Crofts.

Brown, A.G. and D.N. Franz (1969) Response of spinocervical tract neurones to natural stimulation of identified cutaneous receptors. Exp. Brain. Res. 7: 231-249.

Brown, P.B. and J.L. Fuchs (1975) Somatotopic representation of himdlimb skin in the cat dorsal horn. J. Neurophysiol. 38: 1-9.

Brown, A.G., P.K. Rose and P.J. Snow (1977) The morphology of spinocervical tract neurones revealed by intracellular injection of horseradish peroxidase. J. Physiol 270: 747-764.

Brown, A.G. and R.E.W. Fyffe (1978) The morphology of group Ia afferent fibre collaterals in the spinal cord of the cat. J. Physiol. 274: 111-127.

Brown, A.G. and R.E.W. Fyffe (1980) Projections from pacinian corpuscles and rapidly adapting mechanoreceptors of glabrous skin to the cat's spinal cord. J. Physiol. 307: 385-400.

Brown, A.G. and R.E.W. Fyffe (1981) Form and function of dorsal horn neurones with axons ascending the dorsal columns in the cat. J. Physiol. 321: 31-47.

Brown, A.G. (1981). Organization in the spinal cord: The anatomy and physiology of identified neurones. In Springer-Verlag (Ed.), Berlin:

Brown, A.G., P.B. Brown and R.E.W. Fyffe (1983) Receptive field organization and response properties of spinal neurones with axons ascending dorsal columns in the cat. J. Physiol. 337: 575-588.

Brown, P.B., H.R. Koerber and L.A. Ritz (1992) Laminar organization of primary afferent termination in the mammalian spinal cord. In S. A. Scott. (eds): Sensory neurons, diversity, development and plasticity. Oxford: Oxford University Press, pp. 116-130.

Bryan, R.N., D.L. Trevino, J.D. Coulter and W.D. Willis (1973) Location and somatotopic organization of the cells of origin of the spino-cervical tract. Exp. Brain Res. 17: 177-189.

Bryan, R.N., J.D. Coulter and W.D. Willis (1974) Cells of origin of the spinocervical tract in the monkey. Exp. Neurol. 42: 574-586.

Burckhardt, R. (1889) Histologische Untersuchungen am rückenmark der tritonen. Arch. Mikr. Anat. 34: 131-156.

Burgess, P.R. and E.R. Perl (1967) Myelinates afferent fibres responding specifically to noxius stimulation of the skin. J. Physiol 190: 541-562.

Burguess, P.R. and F.J. Clack (1969) Dorsal column projection of fibres from the cat Knee joint. J. Physiol 203: 301-315.

Burke, R., A. Lundberg and F. Weight (1971) Spinal border cell origin of the ventral spinocerebellar tract. Exp. Brain Res. 12: 283-294.

Burr, H.S. (1928) The central nervous system of Orthagoriscus mola. J. Comp. Neurol. 45: 33-127.

Burstein, R., K.D. Cliffer and G.J.J. Geisler (1987) Direct somatosensory projections from the spinal cord to the hypothalamus and telencephalon. J. Neurosci. 7(12): 4159-4164.

Burstein, R., K.D. Cliffer and G.J.J. Giesler (1990) Cells of origin of the spinohypothalamic tract in the rat. J. Comp. Neurol. 291: 329-344.

Campbell, J.N. and R.H. LaMotte (1983) Latency to detection of first pain. Brain Res. 266: 203-208.

Cao, C.Q., L. Djouhri and A.G. Brown (1993) Lumbosacral spinal neurons in the cat that are candidates for being activated by collaterals from the spinocervical tract. Neuroscience 57: 153-165.

Carlsen, R.C. and L.M. Mendell (1977) A comparison of the reflex organization of thoracic and lumbar segments in the frog spinal cord. Brain Res. 124: 415-426.

Catton, W.T. (1958) Some properties of frog skin mechanoreceptors. J. physiol. 141: 305-322.

Catton (1976) Cutaneous mechanoreceptors. In Frog Neurobiology (eds. Llinás R and Precht W.) 629-641. Springer-Verlag, Berlin-Heidelberg-New York.

Cervero, F., A. Iggo and V. Molony (1977) Response of spinocervical tract neurones to noxious stimulation of the skin. J. Physiol 267: 537-558.

Cervero, F. and A. Iggo (1980) The substantia gelatinosa of the spinal cord. A critical review. Brain. 103: 717-772.

Cervero, F. (1989) The superficial dorsal-horn. In F. B. Cervero G. J. and Headley, P. M. (eds): Processing of sensory information in the superficial dorsal horn of th spinal cord. New York and London: Plenum Press, pp. 1-12.

Cervero, F. and R.D. Foreman (1990) Sensory innervation of the viscera. In Arthur D. Loewy and K. Michael Spyer (eds): Central regulation of autonomic functions. Oxford: Oxford University Press, pp. 104-125.

Chahl, L.A., J. Szolcsányi and F. Lembeck (1984) Antidromic vasodilatation and neurogenic inflamation. Akadémiai Kiadó Budapest.

Chung, J.M., D.R. Kenshalo, K.D. Gerhart and W.D. Willis (1979) Excitation of primate spinothalamic neurons by cutaneus C fiber volleys. J. Neurophysiol 42: 1354-1369.

Comer, C. and P. Grobstein (1981) Organization of sensory imputs to the midbrain of the frog *Rana pipiens*. J. Comp. Physiol 142: 161-168.

Cooper, S. and C.S. Sherrington (1940) Grower's tract ans spinal border cells. Brain 63: 123-134.

Corvaja, N. and I. Grofova (1972) Vestibulospinal projection in the toad. Prog. Brain Res. 297-307.

Corvaja, N., I. Grofová and O. Pompeiano (1973) The origin, course and termination of vestibulospinal fibers in the toad. Brain Behav. Evol. 7: 401-423.

Corvaja, N. and M. Pellegrini (1975) Ultrastructure of dorsal root projections in the toad cord. Arch. Ital. Biol. 113: 122-149.

Corvaja, N. and I. Grofová (1978) Vestibulospinal projections in the toad. An experimental electron microscope study. Neurosci. 3: 619-628.

Craig, A.D. and S. Mense (1983) The distribution of afferents fibers from the gastrocnemius-soleus muscle in the dorsal horn of the cat as revealed by the transport of horseradish peroxidase. Neurosci. Lett. 41: 233-238.

Craig, A.D. and K.D. Kniffki (1985) Spinothalamic lumbosacral lamina I cells response to skin and muscle stimulation in the cat. J. Physiol. 365: 197-221.

Craigie, E.H. (1928) Observations on the brain if the humming bird (*Chrysolampis mosquitis* linn. and *Chlorostilbon caribaeus* Lawr.). J. Comp. Neurol. 45: 377-481.

Cruce, W.L.R. (1974) The anatomical organization of hindlimb motoneurons in the lumbar spinal cord of the frog. *Rana catesbeina*. J. Comp. Neurol. 153: 59-76.

Dado, R.J., J.T. Katter and J.R. Giesler (1994a) Spinothalamic and spinohypothalamic tract neurons in the cervical enlargements of rats I. Locations of antidromically identified axons in tha thalamus and hypothalamus. J. Neurophysiol. 71: 959-980.

Dado, R.J., J.T. Katter and J.R. Giesler (1994b) Spinothalamic and spinohypothalamic tract neurons in the cervical enlargements of rats II. Responses to innocuous and noxious mechanical and thermal stimuli. J. Neurophysiol. 71: 981-1002.

Dado, R.J., J.T. Katter and J.R. Giesler (1994c) Spinothalamic and spinohypothalamic tract neurons in the cervical enlargements of rats III. locations of antidromically identified axons in the cervical cord white matter. J. Neurophysiol. 71: 1003-1021.

Dailenbach, K.M. (1927) The temperature spots and endorgans. Amer. J. Psychol. 39: 402-427.

Deuchar, E.M. (1975). Xenopus: the South African Clawed frog. In D. Publications (Ed.), Nueva York:

Dodt, E., A.P. Skouby and Y. Zoterman (1953) The effects of cholinergic substances on the discharges from thermal receptors. Acta Physiol Scand. 28: 101-114.

Donaldson, H.H. (1885) On the temperature sense. Mind. 10: 399-416.

Duce, I.R. and P. Keen (1977) An ultrastructural Classification of the neuronal cell bodies of the rat dorsal root ganglion using zinc iodine-osmiun impregnation. Cell. Tiss Res. 185: 263-277.

Ebbesson, S.O.E. (1967) Ascending axon degeneration following hemisection of the spinal cord in the tegu lizard (*Tupinambis nigropunctatus*). Brain Res. 5: 178-206.

Ebbesson, S.O.E. (1969) Brainstem afferents from the spinal cord in a sample of reptilian and amphibian species. Ann. N.Y. Acad. Sci. 167: 80-101.

Ebbesson, S.O.E. (1976) Morphology of the spinal cord. In Springer-Verlag. (eds): Frog neurobiology. Berlin-Heidelberg-New York: pp. 679-706.

Ebbesson, S.O.E. (1978) Somatosensory pathways in lizards: the identification of the medial lemniscus and related structures. In: Greenberg and Maclean (eds): Behavior and Neurology of Lizards. U.S. Department of Health, Education and Welfare, Rockville 91-104.

Ebbesson, S.O.E. and K.C. Hodde (1981) Ascending spinal systems in the nurse shark Ginglymostoma cirratum. Cell Tissue Res. 216: 313-331.

Fernandez de Molina, A., J.L. Garcia-Sanchez and A. Ruiz Marcos (1966) Frog's dorsal column nuclei potentials generated by stimulation of the sciatic nerve. Trabajo del Instituto Cajal de investigaciones biologicas. 58: 171-184.

Ferrington, D.G., L.S. Sorkin and W.D. Willis (1987) Response of spinothalamic tract cell in the superficial dorsal horn of the primate lumbar spinal cord. J. Physiol. 388: 681-703.

Flelds, H.L., G.M. Wagner and S.D. Amerson (1975) Some properties of spinal neurons projecting to the medial brain-stem reticular formation. Exp. Neurol. 47: 118-134.

Fields, H.L., C.H. Clanton and S.D. Anderson (1977) Somatosensory properties of spinoreticular neurons in the cat. Brain Res. 120: 49-66.

Finger, T.E. (1978) Efferent neurons of the teleost cerebellum. Brain Res. 153: 608-614.

Finger, T.E. (1981) Fish that taste with their feet: spinal sensory pathways in the sea robin, *Prionotus carolinus*. Biol. Bull. 161: 343.

Foerster, O. and O. Gagel (1932) Die vorderseitenstragdurchneidung beim menscchen. Eine klinisch-pathophysiologisch-anatomische studie. Z. Ges. Neurol. Psychiat. 138: 1-92.

Forehand, C.J. and P.B. Farel (1982) Spinal cord development in anuran larvae. I. primary and secondary neurons. J. Comp. Neurol. 209: 386-394.

Frank, E. and M. Westerfield (1982) Synaptic organization of sensory and motor neurones innervating triceps brachii muscle in the bullfrog. J. Physiol 324: 479-494.

Fuller, P.M. and S.O.E. Ebbenson (1973) Central projections of the facial nerve in the bullfrog (*Rana catesbeiana*). J. Morphol. 152: 193-200.

Fuller, P.M. (1974) Projections of the vestibular nuclear complex in the bullfrog (*Rana catesbeiana*). Brain Behav. Evol. 10: 157-169.

Funke, K. and R. Necker (1986) Cells of origin of ascending pathways in the spinal cord of the pigeon. Neurosci. Let. 71: 25-30.

Funke, K. (1988) Spinal projections to the dorsal column nuclei in pigeons. Neurosci. Let. 295-300.

Funke, K. (1989) Somatosensory areas in the telencephalon of the pigeon II. Spinal pathways and different connections. Exp. Brain Res. 76: 620-638.

Fyffe, R.E.W. (1979) The morphology of group II afferents fibre collaterals. J. Physiol. 296: 39-40.

Fyffe, R.E.W. (1992) Laminar organization of prymary afferents terminations in the mammalian spinal cord. In S. A. Scott. (eds): Sensory neurons, diversity, development and plasticity. Oxford: Oxford University Press, pp. 131-142.

Gasser, H.S. (1950) Unmedullated fibers originating in dorsal root ganglia. J. Gen. Physiol. 3: 651-690.

Gasser, H.S. (1955) Propierties of dorsal root unmedulated fibers on the two sides of the ganglion. J. Gen. Physiol. 38: 709-728. Gaupp, E. (1899). A Ecker's und R. Wiedersheim's anatomie des frosches. In Lehre vom nerven- und Gefässsystem. (Vol. 2, Vieweg, Braunschweig.:

Georgopoulos, A.P. (1977) Stimulus-response relations in high-threshold mechanothermal fibers innervating primate glabrous skin. Brain Res. 128: 547-552.

Giesler, G.J. and K.D. Cliffer (1985) Postsynaptic dorsal column pathway of the rat. II. Evidence against an important role in nocicepcion. Brain Res. 326: 347-356.

Glees, P. and J. Soler (1951) Fibre content of the posterior column and synaptic connections of nucleus gracilis Z. Zellforsch. 36: 381-400.

Goldby, F. and L.R. Robinson (1962) The central connexions of dorsal spinal nerve roots and the ascending tracts in the spinal cord of *Lacerta viridis*. J. Anat. Lond. 96: 153-170.

Goldscheider, A. (1884) The specific energy of the sensory nerves of the skin. Vth world congress on pain. Hamburg. 47-67.

González, A., H.J. ten Donkelaar and R. de Boer-van Huizen (1984) Cerebellar connections in *Xenopus laevis*. An HRP study. Anat Embryol. 169: 167-176.

González, A., A. Muñoz and M. Muñoz (1993) Trigeminal primary afferent projections to the spinal cord of the frog, *Rana ridibunda*. J.Morphol. 217: 137-146.

Goowin, G.M., D.I. McCloskey and P.B. Matthews (1972) The persistence of appreciable kinesthesia after paralysing joint afferents but preserving muscle afferents. Brain Res. 37: 326-329.

Grant, G. (1962) Spinal course and somatotopically localized termination of the spinocerebellar tracts. An experimental study in the cat. Physiol Scand. 56 (Suppl.), 193: 1-45.

Grantyn, R., A.I. Shapovalov and B.I. Shiriaev (1982) Combined morphological and electrophysiological description of connections between single primary afferents fibres and individual motoneurons in the frog spinal cord. Exp Brain Res. 48: 459-462.

Grantyn, R., A.I. Shapovalov and B.I. Shiriaev (1984a) Tracing of frog sensory-motor synapses by intracellular of horseradish peroxidase. J. Physiol (Lond). 349: 441-458.

Grantyn, R., A.I. Shapovalov and B.I. Shiriaev (1984b) Relation between structural and release parameters at the frog sensory-motor synapse. J. Physiol (Lond). 349: 459-474.

Hamano, K., H. Mannen and N. Ishizuka (1978) Reconstruction of trajectory of primary afferent collaterals in the dorsal horn of the cat spinal cord, using Golgi-stained serial sections. J. Comp. Neurol. 181: 1-16.

Hardy, J.D., H. Goodell and H.G. Wolff (1951) The influence of skin temperature upon the pain threshold as evoked thermal radiation. Science 114: 149-150.

Harrington, T. and M.M. Merzenich (1970) Neural coding in the sense of touch: human sensations of skin indentation compared with the response of slowly adapting mechanoreceptive afferents innervating the hair skin of monkeys. Exp. Brain. Res. 10: 251-264.

Hayle, T.H. (1973a) A comparative study of spinal projections to the brain (except cerebellum) in three classes of poikilothermic vertebrates. J. Comp. Neurol 149: 463-476.

Hayle, T.H. (1973b) A comparative study of spinocerebellar systems in three classes of poikilothermic vertebrates. J. Comp. Neurol. 149: 477-496.

Head, H. (1920). Studies in neurology. In Oxford Univ. Press. (Vol. 1, pp. 55-65 y 225-329). Oxford:

Heier, P. (1948) Fundamental principles in the structure of the brain: A study of the brain of *Petromyzon fluviatilis*. Acta Anat. 5 (Suppl. 8): 1-213.

Hensel, H. (1950) Temperaturempfindung und intracutane waermebewegung. Pfluegers Arch. 252: 165-215. Hensel, H. and Y. Zotterman (1951) The effects of

Hensel, H. and Y. Zotterman (1951) The effects of menthanol on the thermoreceptors. Acta Physiol. Scand 24: 27-34.

Hensel, H. (1973) Cutaneous thermoreceptors. In springer (eds): Somatosensory system. Nueva york: pp. 79-110.

Hensel, H. (1974) Thermoreceptors. Ann Rev. Physiol. 36: 233-249.

Herrick, C.J. (1914) The medulla oblongata of larval Amblystoma. J. Comp. Neurol 24: 343-427.

Herrick, C.J. (1930) The medulla oblongata of Necturus. J. Comp. Neurol 50: 1-96.

Herrick, C.J. (1939) The cerebrum of Amblystoma tigrinum in midlarval stages. J. Comp. Neurol 70: 249-266.

Herrick, C.J. (1948) The brain of the tiger salamander, Ambystoma tigrinum. Chicago: University of Chicago Press

Herrick, C.J. and G.H. Bishop (1957) A comparative survey of the spinal lemniscus systems. In "Reticular Formation of the Brain" (eds. Jasper, H.H., Proctor, L.D., Knighton, R.S., Noshay, W.C., Costello, R.T.) 353-360.

Holder, N., J.D.W. Clarke, N. Stephens, S.W. Wilson, C. Orsi, T. Bloomer and D.A. Tonge (1991) Continuous growth of the motor system in axolotl. J. Comp. Neurol. 303:

Hongo, T., N. Ishizuka, H. Mannen, N. Ishizuka and D. Sasaki (1978) Axonal trajectory of single gruoup 1a and 1b fibers in the cat spinal cord. Neurosci. Lett. 8: 321-328.

Hongo, T., N. Kudo, S. Sasaki, M. Yamashita, K. Yoshida, N. Ishizuka and H. Mannen (1987) Trajectory of group Ia and Ib fibers from the hind-limb muscle at the L3 and L4 segments of the spinal cord of the cat. J. Comp. Neurol. 262: 159-194.

Hoogland, P.V. (1981) Spinothalamic projections in a lizard, Varanus exanthematicus: An HRP study. J. Comp. Neurol 198: 7-12.

Horch, K.W., F.J. Clack and P.R. Burgess (1975) Awareness of knee joint angle under static conditions. J. Neurophysiol 38: 1436-1447.

Hubbard, I.J. and O. Oscarsson (1962) Localization of the cell bodies of the ventral spinocerebellar tract in lumbar segments of the cat. J. Comp. Neurol. 118: 199-204.

Huber, G.C. and E.C. Grosby (1926) On thalamic and tectal nuclei and fibers paths in the brain of the american alligator. J. Comp. Neurol. 40: 97-227.

Hughes, A.F.W. (1957) The development of the primary sensory system in *Xenopus laevis*. J. Anat. 91: 323-338.

Hughes, A. and P.A. Tschumi (1958) The factors controlling the development of the dorsal root ganglia

and ventral horn in Xenopus laevis. J. Anat. 92: 498-527.

Hulshof, J.B.E., R. Boer-van Huizen and H.J. ten Donkelaar (1987) The distribution of motoneurons supplying hind limb muscles in the clawed toad, *Xenopus laevis*. Acta Morphol Neerl-Scand. 25: 1-16.

Hunt, S.P., P.W. Mantyh and J.V. Priestley (1992) The organization of biochemically characterized sensory. In S. A. Scott. (eds): Sensory neurons, diversity, development and plasticity. Oxford: Oxford University Press, pp. 60-76.

Ishizuka, N., H. Mannen, T. Honho and S. Sasaki (1979) Trayectory of group 1A afferent fibers stained with horseradish peroxidase in the lumbosacral spinal cord of the cat: three-dimensional reconstructions from spinal serial sections. J. Comp. Neurol. 186: 189-212.

Ito, M. (1984). The cerebellum and neural control. In Nueva York: Raven Press

Ito, H., T. Murakami, T. Fukuoka and R. Kishida (1986) Thalamic fiber connections in a teleost (*Sebasticus* marmoratus): Visual, Somatosensory, Octavolateral and Cerebellar Relay Region to the telencephalon. J. Comp. Neurol 250: 215-227.

Jacobs, V.L. (1968) an experimental study of the course and termination of the spino-cerebellar systems in a lizard (*Lacerta viridis*). Brain Res. 11: 154-176.

Jankowska, E., J. Rastad and P. Zarzecki (1979) Segmental and supraspinal imput to cell of origin of non-primay fibers in the feline dorsal columns. J. Physiol. 290: 185-200.

Jhavery, S. and E. Frank (1983) Central projections of the brachial nerve in bullfrogs: Muscle and cutaneus afferents project to different regions of the spinal cord. J.Comp.Neurol 221: 304-312.

Jones, M.W., C.J. Hodge, A.V. Apkarian and R.T. Stevens (1985) A dorsolateral spinothalamic pathway in cat. Brain Res. 335: 188-193.

Jones, M.W., C.J. Hodge, A.V. Apkarian and R.T. Stevens (1987) The spinothalamic tract: an examination of the cell of origin of the dorsolateral and spinothalamic pathways in cats. J. Comp. Neurol. 260: 349-361.

Joseph, B.S. and D.G. Whitlock (1968a) Central projection of slectted spinal dorsal roots in anuran amphibians. Anat Rec. 160: 279-288.

Joseph, B.S. and W.D. G. (1968b) Central projections of brachial and lumbar dorsal roots in reptiles. J. Comp. Neurol. 132: 469-484.

Ju, G. (1984) Direct connections between hypothalamus and lumbar spinal cord in rabbits. Sci. Sin. 27: 789-799.

Kamogawa, H. and G.J. Bennet (1986) Dorsal column postsynaptic neurons in the cat are excited by myelinated nociceptors. Brain Res. 364: 386-390.

Kanui, T.I. (1988) Receptive field organization and electrophysiological response of spinal cord thermoreactive neurones in the rat. Exp. Brain Res. 71: 508-514.

Kappers, C.U.A., G.C. Huber and E.C. Crosby (1936). The comparative anatomy of the nervous system of vertebrates including man. In Hafner (Ed.), Nueva York:

Karten, H.J. (1963) Ascending pathways from the spinal cord in the pigeon (Columba livia). Proc XVIth Int. Congr. Zool., 2: 23.

Karten, H.J. and A.M. Rezvin (1966) The afferent connections of the nucleus rotundus in the pigeon. Brain Res. 2: 368-377.

Karten, H.J. and W. Hodos (1967) A sterotaxic atlas of the brain of the pigeon (*Columba livia*). Johns Hopkins University Press

Katz, B. (1961) The terminations of the afferent nerve in the muscle spindle of the frog. Phil. Trans. 243: 221-240.

Kenshalo, D.R., R.B. Leonard, J.M. Chung and W.D. Willis (1979) Response of primate spinothalamic neurons to graded and to repeated noxious heat stimuli. J. Neurophysiol. 42: 1370-1389.

Kerr, F.W.L. (1975) Pain a central inhibitiry balance theory. Mayo Clinic Proc. 50: 685-690.

Kevetter, G.A. and W.D. Willis (1984) Collateralization of the spinothalamic tract: New methodology to support or deny phylogenetic theories. Brain Res. Rev. 7: 1-14. Knibestöl, M. and A.B. Vallbo (1970) Single unit

analysis of mecanoreceptor activity from the human glaborous skin. Acta Physiol Scand. 80: 178-195.

Knibestöl, M. (1973) Stimulus-response functions of rapidly adapting mechanoreceptors in the human glabrous skin area. J. Physiol. 232: 427-452.

Koerber, H.R. and P.B. Brown (1980) Projections of two hindlim nerves to cat dorsal horn. J. Neurophysiol 44: 259-269.

Koerber, H.R. and P.B. Brown (1982) Somatotopic organization of hindlimb cutaneous nerve projections to cat dorsal horn. J. Neurophysiol. 48: 481-489.

Koerber, H.R. and L.M. Mendel (1992) Fuctional heterogeneity of dorsal root ganglion cells. In S. A. Scott. (eds): Sensory neurons, diversity, development and plasticity. Oxford: Oxford University Press, pp. 77-96.

Krause, W. (1859) Ueber nervenendigungen. Z. Rat. Med. 5: 28-43.

Kreht, H. (1940) Die markhaltige Fasersysteme im gehirn der Anuren und Urodelen und ihre Mylogenie; zugleich ein kritischer Beitrag zu den Flechsigschen myelogenettischen Grundgesetzen. I. Einleitung und medulla oblongata. Z. Mikrosk-Anat Forsch 41: 417-432.

Kruger, L. and P. Witkovsky (1961) A functional analysis of neurons in the dorsal column nuclei and spinal nucleus of the trigeminal in the reptile (*Alligator mississippiensis*). J. Comp. Neurol. 117: 97-105.

Kumazawa, T. and E.R. Perl (1977) Primate cutaneous sensory units with unmyelinated (C) afferents fibers. J. Neurophysiol. 40: 1325-1338.

Kuo, D.C., G.C.H. Yang, D.S. Yamaskai and G.M. Kraithamer (1982) A wide field electron microscopic analisys of y the fiber constituents of the major splanchnic nerve in the cat. J. Comp. Neurol. 210: 49-58.

Kuru, M. (1956) The spino-bulbar tracts and the pelvic sensory vagus. Futher contributions to the teory of the sensory dual innervation of the viscera. J. Comp. Neurol. 104: 207-231.

Kusuma, A. (1979) The organization of the spinal cord in reptiles with different locomotor pathways. Doctoral Thesis, Univ. of Nijmegen (The Netherlands)

Künzle, H. and W. Wooson (1982) Mesodiencephalic and other target regions of the ascending spinal projections in the turtle: *Pseudemys scripta elegans*. J.Comp. Neurol 212: 349-364.

Künzle, H. (1983) Spinocerebellar projections in the turtle. Observations on their origin and terminal organization. Exp. Brain Res. 53: 129-141.

Künzle, M.S. (1994) Connections of the basal telencephalic areas c and d in the turtle brain. Anat Embriol 189: 339-359.

LaMotte, R.H. (1978) Comparison of response of warm and nociceptive C-fibers afferents in monkey with human judgments of thermal pain. J. Neurophysiol. 41: 509-528.

Landgren, S. and H. Silfvenius (1969) Projection to cerebral cortex of group I muscle afferents from the cat's hind limb. J. Physiol. 200: 353-372.

Landgren, S. and H. Silfvenius (1971) Nucleus Z, the medullary relay in the projection path to the cerebral cortex of group I muscle afferents from the cat's hind limb. J. Physiol. 218: 551-571.

Larsell, O. (1967) The comparative anatomy and histology of the cerebellum from Myxinoids through birds. Univ. Minnesota Press, Minneapolis 291.

Larson-Prior, L.J. and W.L.R. Cruce (1992) The red nucleus and mesencephalic tegmentum in a ranid amphibian: A cytoarchitectonic and HRP connectional study. Brain Behav Evol. 40: 273-286.

Lawson, S.N. (1979) The postnatal development of large light and small dark neurons in mouse dorsal root ganglion: a statiscal analysis of cell numbers and size. J. Neurocytol 8: 275-294.

Lawson, S.N. (1992) Morphological and biochemical cell types of sensory neurons. In S. A. Scott. (eds): Sensory neurons, diversity, development and plasticity. Oxford: Oxford University Press, pp. 27-59.

Leek, B.F. (1972) Abdominal visceral receptors. In E. Neil. (eds): Handbook of sensory physiology. Enteroceptors. Springer, Nueva York: pp. 113-160.

Leha, J., D. Menétrey and J. De Pommery (1988) Neuroptides in long ascending spinal tract cells in the rat: evidence for parallel processing of ascending information. Neuroscience 24: 195-207.

Lewis, T. and E.E. Pochin (1938) The double pain response of the human skin to a single stimulus. Clin. Sci. 3: 67-76.

Lewis, T. (1942). Pain. In New York: MacMillan

Lichtman, J.W. and E. Frank (1984) Physiological evidence for specifity of synaptic connections between individual sensory and motor neurons in the brachial spinal cord of the bullfrog. J. Neurosci. 4: 1745-1753.

Lieberman, A.R. (1976) Sensory ganglia. In Chapman and Hall (eds): The peripheral nerve. Londres: pp. 188-278.

Light, A.R. and E.R. Perl (1979) Spinal termination of functionally identified primary afferent neurons with slowly conducting myelinated fibres. J. Comp. Neurol. 186: 133-150.

Light, A.R. and R.G. Durkovic (1984) Features of laminar and somatotopic organization of lumbar spinal cord units receiving cutaneous inputs from bindlimb receptive fields. J. Neurophysiol 52: 449-458.

Lima, D. and A. Coimbra (1990) Structural types of marginal (lamina I) neurons projecting to the dorsal reticular nucleus of the medulla oblongata. Neuroscience 34: 591-607. Lima, D. (1990) A spinomedullary projection terminating in the dorsal reticular nucleus in the rat. Neuroscience 34: 577-591.

Lindblom, U. (1962) The relation between stimulus and discharge in a rapidly adapting touch receptor. Acta Physiol. Scand. 56: 349-361.

Lindblom, U. and L. Lund (1966) The discharge from vibration sensitive receptors in the monkey foot. Exp. Neurol. 15: 401-417.

Liu, C.N. and W.W. Chambers (1957) Experimental study of anatomical organization of frog's spinal cord. Anat. Rec. 127: 326.

Liu, C.N. (1969) Discussion. Brazier MAB. UCLA Forum in medical Sciences nº 11 193-204.

Liuzzi, F.J., M.S. Beattie and J.C. Bresnahan (1984) The relatonship of dorsal root afferents to motoneuron somata and dendrites in the adult bullfrog: a light and electron microscopic study using horseradish peroxidase. Neuroscience 11: 951-961.

Lloyd, D.P.C. and A.K. McIntyre (1950) Dorsal column conduction of group I muscle afferent impulse and their relay through Clarke's column. J. Neurophysiol. 13: 39-54.

Loewy, A.D. and R.M. Spyer (1990). Central regulation of autonomic functions. In Arthur D. Loewy and K. Michael Spyer (Ed.), Oxford: Oxford University Press

Low, J.S.T., J. Mantle-St and D.J. Tracey (1986) Nucleus Z in the rat: Spinal afferents from collaterals of dorsal spinocerebellar tact neurons. J. Comp. Neurol. 243: 510-526.

Lundberg, A. and F. Weight (1971) Functional organization of conexions to the ventral spinocerebellar tract. Exp. Brain Res. 12: 295-316.

Macefield, G., S.C. Gandevia and D. Burke (1990) Perceptual response to microstimulation of single afferents innervating joints, muscle, and skin of the human hand. J. Physiol. 429: 113-129.

Magendie, F. (1822) Experiences sur les fonctions des racines des nerfs rachidiens. J. Physiol. Exp. Pathol. 2: 276-279.

Manns, M.D. (1973) Clarke's Column and the dorsal spinocerenellar tract: A review. Brain Behav. Evol. 7: 34-83.

Marahashi, J., K. Mizugushi and I. Tasaki (1952) Action currents in single fibres elicited by stimulation of the skin of the toad and the cat. J. Physiol. 117: 129-151.

Maruhashi, J., K. Mizuguchi and I. Tasaki (1952) Action current in single afferent nerve fibres elicited by stimulation of the skin of the toad and the cat. J. Physiol 117: 129-151.

Marushashi, J., M. K. and I. Tasaki (1952) Action currents in sigle afferent nerve fibres elicited by stimulation of the skin of the toad and the cat. J. Physiol. (Lond) 117: 129-151.

Matesz, C. and G. Székely (1978) The motor column and sensory projections of the branchial cranial nerves in the frog. J. Comp. Neurol. 178: 157-176.

Matsushita, M. and N. Okado (1981) Spinocerebekkar projections to lobulles I and II of the anterior lobe in the cat, as studied by retrograd transport of hoseradish peroxidase. J. Comp. Neurol. 197: 411-424.

Matsushita, M. and T. Tanami (1983) Contralateral termination of prymary afferent axons in the sacral and caudal segments of the cat, as studied by anterograde transport of horseradish peroxidase. J. Comp. Neurol. 220: 206-218.

Maxwell, D.J. and H.R. Koerber (1986) Fine structure of collateral axons originating from feline spinocervical tract neurons. Brain Res. 363: 199-203.

McCloskey, D.I. (1973) Differences by tween the senses of movement and position shown by effects of loading and vibration of muscle in man. Brain Res. 63: 119-131.

McMahon, S.B. and P.D. Wall (1983) A system of rat spinal cord lamina I cells projecting-through the contralateral dorsolateral funiculus. J. Comp. Neurol 214: 217-223.

McMahon, S.B. and P.D. Wall (1985) Electrophysiological mapping of brainstein projections of spinal cord lamina I cell in the rat. Brain Res. 333: 19-26.

Meissner, G. (1859) Untersuchungen ueber den Tastsinn. Z. Rat. Med. 7: 92-118.

Menetrey, D., A. Chaouch and J.M. Besson (1980) Location and properties of dorsal horn neurons at origin of spinoreticular tract in lumbar enlargement of the rat. J. Neurophysiol 44: 862-877.

Menétrey, D. and A.I. Basbaum (1987) Spinal and trigeminal projections to the nucleus of the solitary tract: A possible substrate for somatovisceral and viscerovisceral reflex activation. J. Comp. Neurol. 255: 439-450.

Mense, S. and N.R. Prabhakar (1986) Spinal termination of nociceptive afferents fibres from deep tissues in the cat. Neurosci. Lett. 66: 169-174.

Menshah, P.L. (1974) The course and distribution of the descending fibers of the lateral funiculus of the amphibian spinal cord. Thesis. University of California Milne, R.J., R.D. Foreman and W.D. Willis (1982) Response of primate spinothalmaic neurons located in the sacral intermediomedial gray (Stilling's niucleus) to propioceptive input from the tail. Brain Res. 234: 227-236.

Minderhound, J.M. (1967) Observation on the supraoptic decussations in the albino rat. J. Comp. Neurol. 129: 297-312.

Morin, F. (1955) A new spinal pathway for cutaneus impulses. Amer. J. Physiol 183: 245-252.

Motorina, M.V., Z.A. Tamarova, A.I. Shapovalov and B.I. Shiriaev (1982) Investigation of synaptic connections between thew frog spinal cord primary afferents and motoneurons by means of HRP intracellular injection. Neurophysiology 14: 60-68.

Mountcastle, V.B. (1973) Central nervous mechanisms in mecanoreceptive sensibility. In (eds): Handbook of Physiology - the nervous system III. pp. 789-878.

Murakami, T. and H. Ito (1985) Long ascending projections of the spinal dorsal horn in a teleost, Sebasticus marmoratus. Brain Res. 346: 168-170.

Müller, J. (1940-42). Elements of physiology. In W. Baly (Ed.), (Vol. 2, Londres:

Neary, T.J. and W. Wilczynski (1977) Ascending thalamic projections from the obex region in ranid frogs. Brain Res. 138: 529-533. Neary, T.J. and W. Wilczynski (1986) Auditory

Neary, T.J. and W. Wilczynski (1986) Auditory pathways to the hypothalamus in ranid frogs. Neurosci. Let. 71: 142-146.

Neary, T.J. (1988) Forebrain auditory pathways in ranid frogs. In Fritzsch B, Ryan MJ, Wilczynski W,

Hetherington TE, Walkowiak W. The evolution of amphibian auditory system. New York, Willey, 233-252.

Necker, R. (1989) Cell of origin of spinothalamic, spinotectal, spinoreticular and spinocerebellar pathways in the pigeon as studied by the retrograde transport of horseradish peroxidase. J. himforsch 30: 33-43.

Necker, R. (1990) Cells of origin of ascending and descending as well as branching fibers in the cervical spinal cord of the pigeon. Neurosci. Let. 119: 1-4.

Neisser, U. (1959) Temperature thresholds for cutaneous pain. J. Appl. Physiol. 14: 368-372.

Nieuwenhuys, R. (1964) Comparative anatomy of the spinal cord. Prg. Brain Res. 11: 1-57.

Nieuwenhuys, R. and M. Cornelisz (1971) Ascending projections from the spinal cord in the axolot (Ambystoma mexicanum). Ana. Rec. 169: 388.

Nijensohn, D.E. and F.W.L. Kerr (1975) The ascending projections of the dorsolateral funiculus of the spinal cord in the primate. J. Comp. Neurol 161: 459-470.

Nikundiwe, A.M., R. de Boer van Huizen and H.J. ten Donkelaar (1982) Dorsal root projections in the clawed toad (*Xenopus laevis*) as demonstrated by anterograde labeling with horseradish peroxidase. Neuroscience 7: 2089-2103.

Noordenbos, W. and P.D. Wall (1976) Diverse sensory functions with an almost totally divided spinal cord. A case of spinal cord transection of part of one anterolateral quadrant. Pain 2: 185-195.

Nordin, M. (1990) Low-Threshold mechanoreceptive and nociceptive units with unmyelinated (C) fibers in the human supraorbital nerve. J. Physiol. 426: 229-240.

Northcutt, R.G. and M.B. Pritz (1978) A spinothalamic pathway to the dorsal ventricular ridge in the spectacled caiman *Caima crocodilus*. Anat. Rec. 190: 618-619.

Northcutt, R.G. and S.O.E. Ebbesson (1980) Ascending spinal pathways in the sea lamprey. Soc. for Neurosci. Abstr. 6: 628.

Nyberg, G. and A. Blomqvist (1984) The central projections on muscle afferents fibers to the lower medulla and upper spinal cord: an anatomical study in the cat with the transganglionic transport method. J. Comp. Neurol. 230: 99-109.

Oka, Y., M. Stou and K. Ueda (1986) Ascending pathways from the spinal cord in the Himé salmon (Landlocked red salmon, *Oncorhynchus nerka*). J. Comp. Neurol. 254: 104-112.

Opdam, P. and R. Nieuwenhuys (1976) Topological analysis of the brain stem of the axolotl Ambystoma mexicanum. J. Comp. Neurol. 165: 285-306.

Oscarsson, O. (1973) Fuctional organization of spinocerebellar paths. In A. Iggo. (eds): Handbook of sensory physiology,. New York: Springer-Verlag, pp. 339-380.

Ottoson, W.T. (1976) Cutaneous mechanoreceptors. In R. Llinás and W. Precht (eds): Frog Neurobiology. Berlin-Heidelberg New York: Springer- Verlag, pp. 629-642.

Pearson, A.A. (1936) The acustico-lateral centers and the cerebellum, with fiber connections of fishes. J. Comp Neurol 65: 201-294.

Pedersen, R. (1973) Ascending spinal connections in three species of sidenecked turtle" *Podocnemis unifilis*, *Pelusios subniger, Pelomedusa surufa*. Anat. Rec. 175: 409 (Abstr.). Perl, E.R., D.G. Whitlock and J.R. Gentry (1962) Cutaneous projection to second order neurons of the dorsal column system. J. Neurophysiol 25: 337-358.

Pert, E.R. (1992) Function of dorsal root ganglion neurons: an overview. In S. A. Scott. (eds): Sensory neurons, diversity, development and plasticity. Oxford: Oxford University Press, pp. 3-26.

Peschanski, M. and J.M. Besson (1984) A spinoreticulo-thalamic pathway in the rat: an anatomical study with reference to pain transmission. Neurosci. 12: 165-178.

Poggio, G.F. and V.B. Mountcastle (1960) A study of the fuctional contributions of the lemniscal and spinothalamic systems to somatic sensibility. Bull. Johns Hopkins Hosp. 106: 266-316.

Poggio, G.F. and V.B. Mountcastle (1963) The fuctional properties of ventrobasal thalamic neurons studied in unanesthetized monkeys. J. Neurophysiol 26: 775-806. Potter, H.D. (1965) Mesencephalic auditory region of the bullfrog. J. Neurophysiol 28: 1132-1154.

Price, D.D., J.W. Hu, R. Dubner and R. Gracely (1977) Peripheral suppression of first pain and central summation of secon pain evoked by noxius heat pulses. Pain 3: 57-68.

Price, D.D., R.L. Hayes, M. Ruda and R. Dubner (1978) Spatial and temporal transformations of imput to spinothalamic tract neurons and their relation to somatic sensations. J. Neurophysiol. 41: 933-947.

Pritz, M.B. (1980) Parallels in the organization of auditory and visual systems in crocodiles. In S.O.E. Ebbesson (eds): Comparative Neurology of the Telencephalon. Plenum Publishing Corporation., pp. 331-342.

Pritz, M.B. (1983) The dorsal column nucleus in a reptile. Caiman crocodilus. Neurosci. Lett. 39: 119-123. Pritz, M.B. and M.E. Stritzel (1986) Dorsal funicular projections to the dorsal column nucleus in a reptile, Caiman cocodrilus. J. Comp. Neurol 249: 1-12.

Pritz, M. and M.E. Stritzel (1989) Reptilian somatosensory misbrain: Identification based on input from the spinal cord and dorsal column nucleus. Brain Behav Evol 33: 1-14.

Pritz, M.B. and M.E. Stritzel (1994) Anatomical identification of a postsynaptic dorsal column system in a reptile, *Caiman crocodilus*. Brain Behav. Evol. 43: 233-243.

Rambourg, A., Y. Clermont and A. Beaudet (1983) Ultraetructural features of six types of neurons in rat dorsal root ganglia. J. Neurocytol. 12: 47-66.

Ramon y Cajal, S. (1909). Histoligie du Système Nerveux de l'homme et des vertebrates. In Maloine (Ed.), (Vol. 1, Paris:

Rastad, J., E. Jankowska and J. Westman (1977) Arborization of initial axon collaterals of spinocervical tract cells stained intracellulary with horseradish peroxidase. Brain Res. 135: 1-10.

Rexed, B. (1952) The cytoarchitectonic organization of the spinal cord in he rat. J. Comp. Neurol. 96: 415-466. Rexed, B. (1954) A cytoarchitectonic atlas of the spinal cord in the rat. J. Comp. Neurol. 100: 297-380.

Rexed, B. (1964) Some aspects of the cytoarchitectonics and synaptology of the spinal cord. Prog. Brain. Res. 11: 58-92. Riedel, W. (1976) Warm receptors in the dorsal abdominal wall of the rabbit. Pfluegers Arch. 361: 205-206.

Riss, W., R.A. Pedersen, J.S. Jakway and C.B. Ware (1972) Levels of function and their representation in the vertebrate thalamus. Brain Behav. and Evol. 6: 26-41.

Roberts, A. and J.D.W. Clarke (1983) The sensory systems of embryos of the newt: *Triturus vulgaris*. J. Comp. Physiol. 152: 529-534.

Roland, P.E. and H. Ladegaard-Pedersen (1977) A quantitative analisis of sensations of tendon and of kinaesthesia in man. Brain. 100: 671-692.

Ronan, M. and G. Northcutt (1990) Projections ascending from the spinal cord to the brain in Petromyzontid and Mysinoid Agnathans. J. Comp. Neurol 291: 491-508.

Rosenthal, B.M. and W.L.R. Cruce (1985) Distribution and ultrastructure of primary afferent axons in Lissauer's tract in the Northern Leopard Frog (*Rana pipiens*). Brain Behav. Evol. 27: 195-214.

Roth, G. and D.B. Wake (1985) The structure of the brainstem and cervical spinal cord in lungless salamanders (Family *Plethodontidae*) and its relation to feeding. J. Comp. Neurol. 241: 99-110.

Ruffini, A. (1894) Sur un nouvel organe nerveux terminal et sur la presence des corpuscles Golgi-Manzzoni dans le conjunctif sous-cutane de la pulpe des doigts de l'homme. Arch. Ital Biol. 21: 249-265.

Rushmer, D.S. (1970) Respons of Purkinje cells in the frog cerebellum to electrical and natural stimulation. Brain. Res. 33: 315-335.

Rushmer, D.S. and D.J. Woodward (1971) Responses of Purkinje cells in the frog cerebellum to electrical and natural stimulation. Brain Res. 33: 315-335.

Rustioni, A. and R.J. Weimberg (1989) The somatosensory system. In A. Björklund, T Hökfelt, and L.W. Swanson (eds): Handbook of chemical neuroanatomy. Vol. 7: Integrated systems of the CNS, Part II. Central Visual, Auditory, Somatosensory, Gustatory. Amsterdam: Elsevier. 219-321.

Schady, W.J.L., H.E. Torebjörk and J.L. Ochoa (1983) Peripheral projections of nerve fibres in the human median nerve. Brain Res. 277: 249-261.

Schneider, A. and R. Necker (1989) Spinothalamic projections in the pigeon. Brain Res. 484: 139-149.

Shapovalov, A.I. and B.I. Shiriaev (1984) Synaptic connexions between primary afferents and thoracic motoneurons in the frog. Exp. Brain. Res. 53: 467-472. Sherrington, C.S. (1898) Experiments in examination of the peripheral distribution of the fibers of the posterior roots of some spinal nerves. Parts II. Phill. Trans. 190: 45-186.

Shiriaev, B.I. and O.V. Shupliakov (1986) Synaptic organization of dorsal root projections to lumbar motoneurons in the clawed toad (*Xenopus laevis*). Exp. Brain Res. 63: 135-142.

Siemen, M. and H. Künzle (1994a) Afferent and efferent connections os the dorsal column nuclear complex and adjacent regions in the turtle. J. Brain Res. 35: 79-102.

Siemen, M. and H. Künzle (1994b) Connections of the basal telencephalic areas c and d in the turtle brain. Anatomy and Embryology 189: 339-359.

Silver, M.L. (1942) The motoneurons of the spinal cord of the frog. J. Comp. Neurol. 77: 1-40.

Silvey, G.E., R.L. Gulley and R.A. Davidoff (1974) The frog dorsal column nucleus. Brain. Res. 73: 421-437. Sinclair, D.C. (1955) Cutaneous sensation and the

doctrine of specific energy. Brain 78: 584-614.

Sinclair, D. and B.A.R. Stokes (1964) The production and characteristics of "second pain". Brain 87: 609-618. Sinclair, D. (1981) Mechanisms of cutaneous sensation. Oxford Univ. Press.

Sinn, R. (1913) Beitrag zur Kenntnis der medulla oblongata der Voegel. Monatschr. F. Psych. Neurol. 33: 1-39.

Smeets, W.J.A.J. (1982) The afferent connections of the tectum mesencephali in two chondrichthyans, the shark *Scyliorhinus canicula* and the ray *Raja clavata*. J. Comp. Neurol. 205: 139-152.

Smeets, W.J.A.J., R. Nieuwenhuys and B.L. Roberts (1983). The central nervous system of cartolaginous fishes. Structure fuctional correlations. In Springer-Verlag (Ed.), Berlin Heidelberg-New York:

Smith, C.F. and E. Frank (1988a) Specificity of sensory projections to the spinal cord during development in bullfrogs. J. Comp. Neurol. 269: 96-108.

Smith, C.F. and E. Frank (1988b) Peripheral spicification of sensory connections in the spinal cord. Brain Behav. Evol. 31: 227-242.

Snow, P.J., P.K. Rose and B. A.G (1976) Tracing axons and axon collaterals of spinal neurons using intracellular injection of horseradish peroxidase. Science N.Y. 191: 114-124.

Sotelo, C. and I. Grofová (1976) Ultrastrutural features of the spinal cord. In Springer (eds): Frog Neurobiology. Berlin-Heidelberg New York: pp. 707-727.

Spray, D.C. (1976) Pain and temperature receptors of anurans. In R. Llinás and W. Precht (eds): Frog Neurobiology. Berlin-Heidelberg New York: Springer-Verlag, pp. 607-628.

Studnica, F.K. (1895) Ein beitrag zur vergleichenden histologie und histogenese des rüskenmarkes. Ber Königl Böhm Gesells Wiss Mathemar-Naturwiss classe 51: 1-32.

Sugiura, Y. and C.L. Lee (1986) Central projection of identified, unmyelinated (C) afferent fibers innervating mammalian skin. Science 234: 358-361.

Sugiura, Y., N. Terui and Y. Hosoya (1989) Difference in distribution of central terminals between visceral and somatic unmyelinated (c) primary afferent fibers. J. Neurophysiol. 62: 834-840.

Surmeier, D.J., C.N. Honda and W.D. Willis (1988) Natural grouping of primate spinothalamic neurons based on cutaneous stimulation. Physiologycal and anatomical features. J. Neurophysiol. 59: 833-860.

Svensson, B.A., J. Westman and J. Rastad (1985) Light and electron microscopic study of neurones in the feline lateral cervical nucleus with a descending projection. Brain Res. 361: 114-124.

Swett, J.E., S.B. McMahon and P.D. Wall (1985) Long ascending projections to the midbrain from cell of lamina I and nucleus of the dorsolateral funiculus of the rat spinal cord. J. Comp. Neurol. 238: 401-416.

Székely, G. (1976) The morphology of motoneurons and dorsal root fibers in the frog spinal cord. Brain Res. 103: 275-290.

Székely, G., N. Antal and T. Gorcs (1980) Direct dorsal root projection onto the cerebellum in the frog.

Székely, G., K. Matesz, R.E. Baker and M. Antal (1982) The termination of cutaneous nerves in the dorsal horn of the spinal cord in normal and in skin-rotated frogs. Exp. Brain. Res. 45: 19-28.

Székely, G. and M. Antal (1984) Segregation of mucles and cuteneous afferent fibre terminals in the brachial spinal cord of the frog. J. Himforsch. 23: 671-675.

ten Donkelaar, H.J. and R. de Boer-van Huizen (1978) A possible pain control system in a non-mammalian vertebrate (a lizard *Gekko gecko*). Neurosci. Let. 83: 65-70.

ten Donkelaar, H.J. (1982) Organization of descending pathways to the spinal cord in clawed toad (Xenopus laevis). Prog. Brain Res. 57: 2297-2311.

ten Donkelaar, H.J. and R. de Boer-van Huizen (1991) Observations on the development of ascending spinal pathways in the clawed toad, *Xenopus laevis*. Anat. Embryol. 183: 589-603.

tenDonkelaar, H.J. (1992) Development of motor systems: A comparative approach. Eur. J. Morphol. 30: 9-22.

Thors, F. (1980) De ontwikkeling van het ruggemerg van de klauwpad, *Xenopus laevis*. Thesis. University of Nijmegen.

Tretjakoff, D. (1909) Das Nervensystem von Ammocoetes. I. Das ruckenmark. Arch. Mikrosk. Anat. 73: 607-680.

Urbán, L. and G. Szekely (1982) The dorsal column nuclei of the frog. 7: 1187-1196.

Vallbo, A.B. (1981) Sensations evoked from the glabrous skin of the human hand by electrical stimulation of unitary mechanosensitive afferents. Brain Res. 215: 359-363.

Vallbo, A.B., K.A. Olsson, K.G. Westberg and F.J. Clark (1984) Microstimulation of single tactile afferents from the human hand. Brain 107: 727-749.

van den Akker, L.M. (1970) An anatomical outline of the spinal cord of the pigeon. Thesis, University of Leiden (The Netherlands)

van Gehuchten, A. (1897) La moèlle èpinière des larves bactraciens (*Salamandra maculosa*). Arch. Biol. 15: 599-619.

Vesselkin, N.P., A. Agayan and L.M. Nomokonova (1971) A study of thalamo-telencephalic afferent systems in frogs. Brain, Behav. Evol. 4: 295-306.

Vesselkin, N.P. and N. Kovacevic (1973) Non-olfactory projections of frog telencephalon. Neurophysiology (Kiev) 5: 537-544.

Vielvoye, G.J. (1977) Spinocerebellar tracts in the lerhorn (*Callus domesticus*). Doctoral Thesis Univ. Leiden (The Netherlandds)

von Frey, M. (1896) Treatise on the sensory fuctions of the human skin. Vth congress on pain, Hamburg.

von Frey, M. (1906) The distribution of afferent nerves in the skin. J.A.M.A. 47: 645-648.

von Frey, M. (1910) Physiologie der Sinnesorgane der Menschlichen Haut. Ergebnisse Physiol. 9: 351-368.

Wahnschaffe, U., U. Bartsch and B. Fritzsch (1987) Metamorphic changes within the lateral-line system of Anura. Anat. Embryol. 175: 431-442.

Weddell, G. (1955) Somesthesis and the chemical senses. Ann. Rev. Psychol 6: 19-136.

White, J.C. and W.H. Sweet (1969). Pain and the Neurosurgeon. In Thomas (Ed.), Springfield:

Wicht, H. and W. Himstedt (1988) Topologic and connectional analysis of the dorsal thalamus of *Triturus alpestris* (Amphibia Urodela Salamandridae). J. Comp. Neurol. 267: 545-561.

Wilczynski, W. (1981) Afferents to the midbrain auditory center in the bullfrog *Rana catesbeiana*. J. Comp. Neurol. 198: 421-433.

Wilczynski, W. and T.J. Neary (1986) Terminal field organization in the bullfrog torus semicircularis. Soc. Neurosci. Abstr. 12: 105.

Wild, J.M. (1983) Connectional analysis of the somatosensory system of the pigeon. Neurosci. Abstr. 9: 244.

Wild, J.M. (1985) The avian somatosensory system. I. primary spinal afferent input to the spinal cord and brainstem in the pigeon (*Columba livia*). 240: 377-395. Wild, J.M. (1989) Avian somatosensory system II. Ascending projections of the dorsal column and external cunneate nuclei in the pigeon. J. Comp. neurol 287: 1-18.

Willer, J.C. and D. Albe-Fessard (1983) Futher studies on the role afferent input from relatively large diameter fibers in transmission of nociceptive messages in humans. Brain Res. 278: 318-321.

Willis, W.D., D.L. Trevino, J.D. Coulter and R.A. Maunz (1974) Responses of primate spinothalamic tract neurons to natural stimulation of hindlimb. J. Neurophysiol 37: 358-372.

Willis, W.D. and R.E. Coggeshall (1991) Sensory mechanisms of the spinal cord. Eds. Plenum press. New York and London

Wilson, P., D.E.R. Meyers and P.J. Snow (1986) The detailed somatotopic organization of the dorsal horn in the lumbosacral enlargement of the cat spinal cord. J. Neurophysiol 55: 604-617.

Winter, D.L. (1971) Receptor characteristics and conduction velocities in bladder afferents. J. Psychiat. Res. 8: 225-235.

Wolf, S. and J.D. Hardy (1941) Studies on pain. Observation on pain due to local cooling and on factors involved in the "cold pressor" effects. J. Clin. Invest. 20:

Woodburne, R.T. (1939) Certain phylogenetic anatomical relations of localizing significance for the mammalian central nervous system. J. Comp. Neurol. 71: 215-257.

Yamada, J. and K. Otani (1978) The spinoperiventricular fiber system in the rabbit, rat, and cat. Exp. Neurol. 61: 395-406.

Yamamori, T. (1970) A light and electron microscopic study in the postnatal development of spinal ganglia in rats. Acta Ant. Nippon. 45: 191-204.

Yezierski, R.P. (1988) Spinomesencephalic tract: projections from the lumbosacral spinal cord of the rat cat and monkey. J. Comp. Neurol 267: 131-146.

Yoss, R.E. (1952) Studies of the spinal cord, part I. Topographic localization within the dorsal spinocerebellar tract in *Macaca mulata*. J. Comp. Neurol. 97: 5-20.

Zeehandelaar, I. (1921) Ontogenese und phylogenese der hinterstrangkerne in verband der sensibilitaet. Fol. Neurobiol. 12: 1-133.

Zelman, F.P., C.M. Leonard, L.-M. Kow and D. Pfaff (1978) Ascending of the lateral columns of the rat spinal

cord: A study using the silver impregnation and horseradish peroxidase techniques. 62: 298-334.

# CAPÍTULO 2

Consideraciones técnicas

- **2.1.-** The use of <u>in vitro preparations</u> of the isolated amphibian central neuvous system in neuroanatomy and electrophysiology
- 2.2.- Comentarios

The use of <u>in vitro</u> preparations of the isolated amphibian central nervous system in neuroanatomy and electrophysiology

2.1

### Harald Luksch<sup>1</sup>, Wolfgang Walkowiak<sup>1</sup>, Alberto Muñoz<sup>2,3</sup>, Hans J. ten Donkelaar<sup>2</sup>

- 1 Zoologisches Institut der Universität zu Köln, Weyertal 119, D-50923 Köln
- 2 Katholieke Universiteit Nijmegen, Department of Anatomy and Embryology, Faculty of Medical Sciences, P.O. Box 9101, NL-6500 HB Nijmegen
- 3 Departamento de Biología Celular, Facultad de Ciencias Biológicas, Universidad Complutense de Madrid, E-28040 Madrid

## Based on: Journal of Neuroscience Methods (Submitted)

Keywords: Amphibians, isolated brain preparation, tract-tracing techniques, intracellular recording and staining, auditory pathways.

#### ABSTRACT

In the present study an isolated preparation of the complete anuran central nervous system (CNS) is described which can be kept alive for several days and allows tracing, immunohistochemical and electrophysiological studies. A simple perfusion chamber is being used in which the isolated CNS preparation is superfused with oxygenated Ringer. The use of an isolated CNS has many advantages including: 1) virtually all areas are easily accessible at the same time without having the problem of blood vessels that hinder access; 2) large lesions and massive tracer applications are possible without survival problems of the animal, and tracers will not be translocated by blood circulation; 3) since pulsations caused by the pressure changes of blood circulation do not occur, intracellular recordings are comparatively easy and stable, and 4) this approach offers the possibility to work on the same brain for

several days by storing the preparation in a refrigerator overnight at low temperatures, thus allowing extensive utilization of a single preparation and a reduction of experimental animals required. Some applications to the anuran auditory system illustrate that the isolated anuran CNS is well-suited for a variety of neuroanatomical and physiological techniques.

### INTRODUCTION

Many isolated preparations of the central nervous system (CNS) have been developed and their success often depends on special features of the preparation that enable it to survive outside the body or on elaborate support strategies that reduce the effects due to absence of a blood supply. A factor that enhances viability of isolated CNS preparations is a decreased need for oxygen and metabolic substrates and this can be achieved in some cases by cooling mammalian preparations but is often found naturally in non-mammalian species.

Among such species are the lamprey and freshwater turtles. The lamprey spinal cord is thin, has no intrinsic blood vessels, and is oxygenated directly from the cerebrospinal fluid (CSF). The brain stem has intrinsic blood vessels, but is probably also oxygenated to a large extent from the CSF (Brodin and Grillner, 1990). Rovainen (1967a,b) took advantage of these favourable conditions by developing an *in vitro* preparation of the lamprey nervous system. The isolated spinal cord can be maintained *in vitro* for periods of 2-3 days at a temperature around 7-10<sup>m</sup>C (Wallén et al., 1985). The brain stem is somewhat more sensitive but can often be maintained during a similar period of time (Brodin and Grillner, 1990). The turtle brain has an unusual resistance to anoxia (Lutz et al., 1985; Hounsgaard and Nicholson, 1990). *In vitro* preparations of isolated parts of the CNS, especially of the telencephalon (e.g., Connors and Kriegstein, 1986; Kriegstein and Connors, 1986; Larson-Prior et al., 1991) and of an isolated cerebellum-brain stem-spinal cord preparation (Keifer and Houk, 1989; Keifer et al., 1992; Sarrafizadeh and Houk, 1994), are used increasingly for combined tracing and electrophysiological studies as well as for pharmacological manipulations of the turtle brain.

For most mammalian tissues it is not possible to maintain adequate physiological integrity without perfusing the vascular system with some form of blood substitute, usually oxygenated artificial CSF. An example of such a preparation is the isolated guinea-pig brain (Llinás et al., 1981; Mühlethaler et al., 1993). Isolated neonatal CNS or brainstem-spinal cord preparations, however, can be kept alive without perfusion of the vascular system. An in vitro brainstem-spinal cord preparation is available for studies on the central control of respiration and locomotion (e.g., Smith and Feldman, 1987). An isolated CNS preparation of the newly born South American opossum, Monodelphis domestica, is a very attractive model for studies on the development and regeneration of synaptic interactions (e.g., Nicholls et al., 1990; Møllgård et al., 1994).

In vitro preparations of the amphibian CNS are increasingly used in physiological studies. Various approaches have been described including the use of a superfused preparation of *Xenopus laevis*  embryos (e.g., Kahn and Roberts, 1982; Roberts and Clarke, 1982; Roberts et al., 1986), brain slices (Holohean et al., 1990), brainstem preparations (Schmidt, 1976; Schaffer 1982; Cochran et al., 1987; Straka and Dieringer, 1993; Atzori and Nistri, 1994; Dicke and Roth, 1994; McLean et al., 1995), and combined spinal cord-muscle preparations (Sagawa et al., 1987; Wheatley and Stein, 1992). So far, the use of isolated CNS preparations for tracing experiments in anurans was restricted to a simple in vitro horseradish peroxidase (HRP) technique in which perfused and subsequently fixed brains are used (McCormick and Braford, 1984; González and Muñoz, 1987), and the cobalt-labeling procedure for fixed brains (e.g., Székely and Gallyas, 1975; Görcs et al., 1979). Straka and Dieringer (1991) used an isolated brainstem-spinal cord preparation for HRP tracing.

In the present study an isolated preparation of the complete anuran CNS will be described which can be kept alive for several days and allows tracing, immunohistochemical and electrophysiological studies. A simple perfusion chamber (modified after Schaffer, 1982) is being used in which the isolated CNS preparation is superfused with oxygenated Ringer. The use of an isolated CNS has many advantages. Some applications on the anuran auditory system will show that the isolated CNS is well-suited for a variety of neuroanatomical and physiological techniques.

#### MATERIAL AND METHODS

Various anuran species were tested for the applicability in vitro: Rana temporaria, R. perezi, Bombina orientalis, Discoglossus pictus, and Xenopus laevis. Animals were deeply anesthetized with tricaine methanesulphonate (MS 222) and cooled to a body temperature of 5<sup>µ</sup>C. The heart was exposed by rapid thoracotomy in order to perfuse the animal transcardially with approximately 40 ml iced Ringer solution (75 mM NaCl, 25 mM NaHCO<sub>3</sub>, 2 mM CaCl<sub>2</sub>, 2 mM KCl, 0.5 mM MgCl<sub>2</sub>, 11 mM glucose) that had been oxygenized with carbogen (95%  $O_2$ , 5%  $CO_2$ ) to a pH of 7.3 (Straka and Dieringer, 1993). Head and vertebral column were then severed and the skin removed to avoid the spread of cutaneous toxins. The following preparation was carried out under a dissecting microscope in a dish the bottom of which was coated with Sylgard (Dow Corning) and covered with ice-cold Ringer solution. The skull was fixed with stainless steel needles sticked into the Sylgard layer. The complete brain and spinal cord were isolated by a dorsal or ventral approach by removing the overlaying bony tissue of the skull and the vertebrae. The nerves were cut with microscissors, special care was taken with the cranial nerves and especially the branchlets of the auditory nerve. After isolation which took approx. 10-15 min, the CNS was transferred into another dish with fresh iced Ringer. Subsequently, the pituitary gland, the dura mater, and the choroid plexus were removed to facilitate oxygen diffusion into the tissue. For ethical reasons, the telencephalon was removed (Mühlethaler et al., 1993). The application of tracer substances usually followed immediately. In case of intracellular studies, the CNS was isolated one day prior to electrophysiological recordings and was stored overnight at 6°C in oxygenated Ringer solution.

#### Tracing techniques

Two different approaches, crystalline application and the injection of aqueous solutions, were used for the application of tracers. These were horseradish peroxidase (HRP, Sigma), biocytin (Ne-Biotinyl-L-lysine, Sigma), biotinylated dextran amines with molecular weights of 3,000 (3 kD BDA, D-7135) or 10,000 (10 kD BDA, D-1956), and various 10 kD or 3 kD lysine-fixable dextran amines conjugated to different fluorochromes, either to tetramethylrhodamine (RDA, D-1817 and D-3308, respectively) or fluorescein (FDA, D-1820 and D-3306, respectively). All dextran amines were purchased from Molecular Probes (Eugene, OR). For massive application of tracers, e.g., after hemisections or in case of superficial application sites, the substances were applied as small crystals dried onto the tips of glass microelectrodes or sharp tungsten needles. Since all substances tested are hydrophilic, the application area had to be rather dry to avoid undesirable spread of the dissolving tracer. A reduction of spread was achieved by taking the brain out of the Ringer solution for a short moment, and rapidly positioning the crystal with the aid of a micromanipulator. Immediately after the application, the brains were submerged again into the Ringer solution that was subsequently changed several times.

For the application of the dextran aminecoupled fluorescent dyes into deeper brain areas, we tested the pressure injection of aqueous solutions (approx. 5-10% substance in distilled water) via glass microelectrodes that were broken down to tip diameters of 10-30  $\mu$ m and attached to a Hamilton syringe. The electrodes were then positioned with a micromanipulator and 1-10  $\mu$ l of the solution were injected. With this approach, the injection speed turned out to be a crucial parameter since rapid injection (more than 1  $\mu$ l per min) resulted in leakage of the substance along the penetration track.

### Intracellular recordings of electrically evoked activity in the auditory pathway

During recording, the CNS was transferred into a perfusion chamber (modified after Schaffer, 1982) and fixed with stainless steel pins to the Sylgard covered floor of the chamber (Fig. 1). The chamber was mounted on an X-Y-table which allowed to move the CNS horizontally in relation to the recording electrode. The preparation was permanently superfused with freshly oxygenated Ringer solution. The flow rate was adjusted to 3-6 ml/min by a peristaltic pump (Masterflex Pump Controller). The solution was guided into the chamber by silicone tubing that ran through an ice container to maintain a superfusate temperature of 16<sup>rd</sup> (Straka and Dieringer, 1993). Temperature and pH or oxygenation, respectively, were monitored continuously.

Single branchlets of the auditory nerve were stimulated electrically with suction electrodes. Stimuli consisted of monophasic square pulses (200  $\mu$ s duration, 1-50  $\mu$ A, 0.5 Hz repetition rate) and were delivered via an isolation unit (WPI, stimulus isolator A 360) which in turn was controlled by an interval generator (WPI, digipulser series 1800). Recordings were done with glass microelectrodes with an impedance of 80-120 M $\Omega$  when filled with 1-2 M potassium-acetate. After removal of the pia mater, electrodes were positioned stereotactically and lowered into the brain with a piezo-driven system (Märzhäuser; PM 10-1). Recorded potentials were amplified (WPI, Cyto 720), digitized (Instrutech, VR100 A) and displayed on a personal computer screen for on-line control and stored on hard disk for off-line analysis.

For intracellular staining, electrodes were filled with a 3.5-4% solution of biocytin in 0.1-0.3 M potassium-acetate. Due to the low molarity of the solution, the impedance of the electrodes was much higher (up to 500 M $\Omega$ ), but this did not influence the recordings significantly. Staining of a neuron was achieved by applying a constant current (1-3 nA) with the iontophoresis unit of the intracellular amplifier for 1-10 min. The membrane potential was controlled every minute to ensure the intracellular position of the electrode. After iontophoresis, the electrode was immediately retracted out of the brain.

#### (Immuno)histochemical procedures

After extracellular application of the tracers or finishing the electrophysiological recordings the CNS was stored in Ringer at room temperature, the solution was changed several times and the pH was monitored to ensure sufficient oxygenation. Over night, brains were put into freshly oxygenized Ringer (pH 7.3; approx. 300 ml) and the container sealed airtight and kept at  $6^{\mu}C$  to slow down metabolism and oxygen consumption of the tissue. After 12 hrs at this temperature, the pH of the Ringer solution typically increased to 7.6-7.8. With this day/night protocol, transport times were usually 16-18 hrs for the 3 kD BDA or fluorescent dextran amines, 20 hrs for biocytin, the 10 kD BDA and the fluorescent dextran amines, and 45 hrs for HRP.

The subsequent steps were dependent on the tracer applied. In the case of HRP and biocytin, brains were fixed with 4% paraformaldehyde and 1.25% glutaraldehyde in phosphate buffer (pH 7.4), or 4% paraformaldehyde in phosphate buffer (pH 7.4) for BDA. Brains were then embedded in gelatine, polyacrylamide or embedding medium (Reichert-Jung) for sectioning on a freezing microtome or in agar or polyacrylamide for sectioning on a vibratome. Section thickness was 20 µm if the sections were directly mounted onto slides after sectioning, and 40-50 m if the following steps were carried out with freefloating sections. For the localization of biocytin and BDA, endogenous peroxidases were blocked by incubation in 0.5%  $H_2O_2$  in phosphate buffer for 15 min. Sections were then rinsed several times, washed with 0.5% Triton-X 100 for 10 min and subsequently incubated with streptavidin-coupled HRP (Amersham; dilution 1:100 in phosphate buffer) for 2 hrs. Like the tracer HRP, the streptavidin (biocytin)- or avidinbiotin (BDA)-coupled HRP was then visualized with the chromogen DAB following a modified protocol of Adams (1981); the peroxide was produced by a glucose-oxidase-reaction (Shu et al., 1988). After the DAB procedure, sections were lightly counterstained with neutral red, dehydrated and coverslipped.

In the case of the fluorescent dextran amines, brains were fixed with 4% paraformaldehyde and sectioned on a freezing microtome (40  $\mu$ m) or on the vibratome (100  $\mu$ m). If immunohistochemical localization of neurotransmitters was desired additionally, standard procedures with fluorescent secondary antibodies were applied. Sections were then dried quickly to avoid fluorescence fading and coverslipped with anti-fading medium (Serva, Fluoromount) or a glycerin-gelatin medium.

#### RESULTS

#### Tracing techniques

The application of HRP showed the wellknown characteristics of this tracer: a dense spot of non-incorporated enzyme at the application site and intense retrograde as well as anterograde labeling of neuronal structures (Fig. 2A). The enzyme was taken up by the soma, by terminal structures, and by damaged fibers of passage; transport speed was approximately 0.5 mm/h. In contrast, the application of biocytin yielded intensely stained neurons at the application site without a diffuse background, indicating that most of the tracer had been taken up by the cells (Fig. 2B). Transport speed was high (2 mm/h) and comparable to the situation in vivo (Luksch and Walkowiak, in preparation). The uptake of biocytin was not restricted to the soma, and resulted in anterograde as well as retrograde labeling. Anterograde labeling (Fig. 2C) was stronger compared to HRP, whereas retrograde labeling seemed to be less intense. However, even the retrograde label achieved with HRP did not stain entire neurons but comprised only primary and secondary dendrites. 10 kD BDA labeling was comparable to that described for biocytin. A much faster axonal transport was observed, however, for 3 kD BDA. With this fast dextran amine Golgi-like labeling of the entire neurons including secondary and tertiary dendrites was achieved (Fig. 2D).

Figure 3 shows an experiment in *Xenopus laevis* in which ascending projections to the torus semicircularis are demonstrated with 3 kD BDA. Labeled cells are found in acoustic and vestibular cell areas, in lateral line related structures, in the dorsal column nucleus, in a lateral cervical nucleus and in the spinal cord. Comparable data were obtained from the other species studied.

The use of the dextran amine-coupled fluorescent dyes (FDA, RDA) yielded best results when applied in crystalline form, both for 3 kD and 10 kD dextran amines, probably since the concentrations in the tissue achieved with the injection of aqueous solutions were too low to result in intense labeling of cells. The transport speed of the 10 kD fluorescent dextran amines was slightly faster for the rhodamine-coupled derivative (3 mm/h) than for the fluorescein-coupled derivative (2 mm/h). Both tracers were transported retrogradely as well as anterogradely, and led to an intense staining of somata and terminal structures up to a distance of 20 mm (Fig. 4A). Like for 3 kD BDA, a faster axonal transport was observed when using 3 kD fluorescent dextran amines.

In the experiments in which fluorescent dextran amine-tracers and immunohistochemical techniques for neuromodulator localization were combined, the protocols did not interfere with each other significantly. Figure 4B shows an example of retrogradely labeled neurons of the nucleus laminaris (torus semicircularis) traced with rhodamine-coupled dextran amin (red) and terminal structures containing the neuromodulator leucine-enkephalin detected with a secondary antibody coupled to FITC (green).

#### Intracellular recordings

The electrical stimulation at the auditory branchlets of the statoacoustic nerve led to neuronal responses in several nuclei of the auditory brainstem. Neurons in other areas of the brain, e.g., in the tectum mesencephali, did not show responses, indicating that the electrical stimulation selectively excited the auditory pathway. The quality of the nerve preparation had a strong influence on the stimulation current that was necessary to excite auditory nuclei; stimulation current was usually 2-10 µA in good preparations but had to be increased up to 50  $\mu$ A if the nerve had been bruised or pulled. Intracellular recordings were possible for up to four days after CNS isolation without a noticeable loss of activity, a decrease of membrane potential in penetrated neurons, or any sign of tissue degeneration in stained structures compared to structures stained in vivo. However, the stimulation current had to be increased probably due to the squeezing of the branchlets with the suction electrode. The longest recording time for one individual neuron was 4.5 h which exceeded that in comparable in vivo recordings by far.

Intracellular application of biocytin yielded intensely stained neurons (Fig. 5). The tracer was distributed homogeneously in soma, dendrites and axonal structures; no gradient was observable so that the recording site in the neuron could not be detected. Transport of the tracer within the cell was fast (approx. 2 mm/h) and comparable to the transport speed found after extracellular biocytin application. The quality of cell staining depended on the duration of the iontophoreses and the current applied but also on electrode characteristics; in some cases, neurons seemed to be completely stained after application of 1 nA for only 1 min. Occasionally, we observed ensembles of stained neurons (2-3) lying closely together. Such simultaneously stained neuronal ensembles were likely no artifacts due to accidental extracellular application of biocytin, since they were also observed when the resting membrane potential of the injected neuron was still high after the iontophoreses, indicating an intracellular position of the electrode. These simultaneously stained neurons usually showed similar dendritic and axonal patterns.

#### Immunohistochemical procedures

The survival time of the preparation, which was defined as the time between the dissection of the animal and the fixation of the isolated CNS, ranged between 16 hrs for the 3 kD dextran amine tracing and 4 days for intracellular biocytin studies. Even after the longest survival times, no sign of tissue degeneration was observed. Labeled neuronal structures had an inconspicuous appearence and could be traced over long distances (several centimeters) without any sign of inhomogeneous tracer distribution or membrane disruption indicating that the fixation by immersion was sufficiently fast to avoid tissue damage. In general, every result obtained in the in vitro preparation was comparable to results collected with an in vivo approach. In those experiments where immunohistochemistry for neuromodulator localization followed the transport phase of the tracer, the distribution of the immunolabeled terminals (e.g., leucine-enkephalin) was indistinguishable from the

pattern found in *in vivo* brains that had been fixed by perfusion of an anesthetized animal.

Each of the various procedures tested for the localization of HRP, biocytin or BDA gave comparable staining of labeled structures. However, since the diffusion of the streptavidin- or avidin-biotin-coupled HRP complexes is limited, the thickness of the sections should not exceed 70  $\mu$ m when processed free-floating or 20  $\mu$ m when processed already mounted onto slides. Staining of erythrocytes that sometimes remained after the perfusion disappeared completely after the blocking of endogeneous peroxidases with H<sub>2</sub>O<sub>2</sub>; this step was only possible for biocytin and BDA.

#### DISCUSSION

#### Methodological considerations

Various physiological studies using in vitro brain approaches in amphibians have been published during the last decades, comprising superfused CNS preparations of Xenopus laevis embryos (e.g., Kahn and Roberts, 1982; Roberts and Clarke, 1982; Roberts et al., 1986) brain slices (Holohean et al., 1990), brainstem preparations (Schmidt, 1976; Schaffer, 1982; Cochran et al., 1987; Straka and Dieringer, 1993; Atzori and Nistri, 1994; Dicke and Roth 1994; McLean et al., 1995) and combined spinal cord-musculature preparations (Sagawa et al., 1987; Wheatley and Stein, 1992). To our knowledge, the use of the complete isolated CNS for tracing experiments has only been reported for HRP (McCormick and Braford, 1984; González and Muñoz, 1987; Straka and Dieringer, 1991).

A crucial question for the evaluation of data collected in such a preparation is whether they are comparable to findings in vivo. We think that the transferability is supported by several arguments. First, no cellular degeneration of the brains was observed even after several days in the Ringer solution; intracellularly stained neurons as well as anterogradely and retrogradely labeled structures showed no differences to neuronal structures stained in vivo (Straka and Dieringer, 1991; Walkowiak and Luksch, 1994; Muñoz et al., 1995). The pattern of labeling in experiments such as the one shown in Fig. 3 is comparable to that obtained in in vivo experiments (e.g., Wilczynski, 1981; Will et al., 1985; Feng and Lin, 1991). Second, intracellularly recorded neurons had resting potentials of up to -90 mV even after three days in the Ringer solution, indicating a good physiological state of the brain. Third, our intracellular recordings showed that the electrical stimulation of the auditory nerve elicited reactions in all nuclei of the auditory pathway including the dorsal medullary nucleus, the superior olive, various structures in the tegmentum mesencephali, and the torus semicircularis, but not in other brain areas. Since afferents to the midbrain include at least two synapses (up to four), the essential neuronal circuits seem to be intact. Additionally, we did not find general differences when recording in vitro and in vivo (Luksch and Walkowiak, 1993). Fourth. the immunohistochemical data on neuromodulator localization showed no differences when compared with results yielded in another study (Luksch and Walkowiak, 1992) where the brains were fixated by perfusion of the anesthetized animal. This finding indicates that even after several days in vitro cellular synthesis and transport systems are functioning properly. Taken together, we think that in amphibians, the *in vitro* approach maintains a physiological status of the brain, allowing anatomical as well as physiological studies.

Of the tracing substances applied extracellularly, HRP has been used for several decades and is well characterized (Mesulam, 1982). In combination with a heavy metal-intensification (Adams, 1981) and the tissue preservating glucoseoxidase-modification (Shu et al., 1988), anterogradely and retrogradely labeled structures are intensely stained. However, even if the retrograde transport exceeds the anterograde one, retrograde labeling of neurons comprises only the soma and the main dendrites. Biocytin has been introduced as an extracellular neuronal tracer only recently (King et al., 1989). The uptake of biocytin seems to rely on a specific, sodium- and ATP-dependent mechanism at the soma (King et al., 1989), and biocytin was therefore characterized as an anterograde tracer. The amount of retrograde transport is described contradictory in the literature (King et al., 1989; Diamond et al., 1991; Izzo, 1991; Kenan-Vaknin et al., 1992) and might depend on the density of terminal structures at the application site (Lapper and Bolam, 1991). In our experiments, retrogradely labeled structures were only weakly stained but comprised all known afferents. The main advantages of biocytin are the fast and strong anterograde transport, a comparatively weak background at the application site which allows the precise identification of the neurons labeled, the possibility to block endogenous peroxidases and the sensitivity and variability of the detection system which allows DAB-precipitation as well as fluorescent label. Biotinylated dextran amines have similar advantages. These tracers are transported retrogradely as well as anterogradely. The retrogradely labeled neurons have an excellent dendritic filling, whereas the tracer can be identified at long distances from the injection site. Particularly the fast 3 kD BDA is extremely useful in the *in vitro* preparation.

Fluorescent dyes coupled to dextran amines have been introduced as neuronal tracers several years ago (Glover et al., 1986; Fritzsch and Wilm, 1990; Nance and Burns, 1990). Small dextran amines with a molecular weight of 3,000 diffuse faster than the larger 10 kD ones (Popov and Poo, 1992; Tao and Nicholson, 1992; Fritzsch, 1993). In the in vitro approach, the substances have characteristics comparable to the in vivo situation, i.e. they are transported anterogradely and retrogradely with 2-3 mm/h and are taken up by intact as well as damaged cells (Glover et al., 1986). Similar findings have been reported in a mammalian brain slice preparation (Boulton et al., 1992). The main advantages of the fluorescent dextran amines are their fast and bidirectional transport, the uptake by all cellular compartments, the possibility to apply two or three differently coupled substances for easy double and triple labeling and the ease of localization. Moreover, double-labeled structures can be analyzed with confocal laser scanning microscopy in great detail. One disadvantage is the instability of some fluorescent dyes (e.g., FITC) to ethanol treatment, thus allowing only very rapid dehydration or coverslipping with water-soluble media. Another problem arises if the combination of fluorescent tracers with immunohistochemistry is desired since

glutaraldehyde, which may be necessary for the binding of the antibody to its epitope, can not be used because it causes autofluorescence.

The use of biocytin for the intracellular staining of neurons has several advantages: Very short application time is required for complete staining of a neuron, the substance is transported rapidly and distributes homogeneously in the cell and the axon, the tip diameter of the electrodes can be small since the molecule is small and does not tend to clog, and the detection system is very sensitive. The finding of simultaneously stained neuronal ensembles is probably due to coupling of these cells via gap junctions (Simpson et al., 1977) rather than unspecific application of biocytin. Similar effects have been reported by other groups working with biocytin (Kawaguchi et al., 1989; Kita and Armstrong, 1991; Wiggers and Roth, 1994; Schulte-Mattler and Luhmann, 1995). This hypothesis is supported by the fact that simultaneously stained neurons usually had comparable dendritic and axonal organization. However, since a comparable percentage of neuronal ensembles is found in vivo, this finding can not be attributed to the isolated brain preparation.

#### General considerations

In our experiments with the isolated brains of amphibians, we did not find any technical limitations concerning the application of tracing substances. We have not tested the application of tracers via iontophoretic injections into deep tissue areas. However, since neuronal circuits remain intact and can be activated across several synapses, the exact localization of nuclear boundaries by multi-unit recordings and the subsequent stereotactic tracer injection (e.g., HRP and biocytin) as described elsewhere (Luksch and Walkowiak, in preparation) should be possible. Moreover, we have not tested the combination of biocytin-application and immunohistochemistry with different DAB-protocols as described by Norgren and Lehman (1990) or Veenman et al. (1992) to yield different light-stable reaction products, which should be possible as well. In general, we think that every tracing technique developed *in vivo* may be applied in an isolated brain preparation as well.

The analysis of physiological parameters in an isolated brain has some obvious limitations, e.g., a stimulation of sensory systems with physiological stimuli is excluded. On the other hand, the stimulation of sensory systems with electrical stimuli leads to an excitation of complete sensory pathways and complex reaction patterns in individual neurons comparable to the findings in intact animals. We therefore think that the basic physiology of neurons can be investigated well in vitro. In some cases, the complete deafferentiation of the brain even may be an advantage of this preparation. In vivo, many physiological parameters of the animal and the environment can not be controlled entirely, e.g., variations due to experimental conditions such as immobilization stress, changing oxygen supply, or the influence of other sensory modalities. The 'constancy' of these parameters is guaranteed in the isolated brain, offering the possibility to analyze the response of single neurons and networks to a reliably constant stimulation and to manipulate the network via the application of neuropharmacological agents or via the stimulation of different afferents.

Besides the limitations for physiological work mentioned above, the use of an isolated CNS has many advantages. First, virtually all areas are easily accessible at the same time without having the problem of blood vessels that hinder access. Second, large lesions and massive tracer applications are possible without survival problems of the animal, and tracers will not be translocated by blood circulation. Third, as pulsations caused by the pressure changes of blood circulation do not occur. intracellular recordings are comparatively easy and stable. Additionally, since the meninges can be completely removed, the penetration of the electrode is facilitated and no limitations to the electrode shape exist. Fourth, this approach offers the possibility to work on the same brain for several days by storing the tissue in a refrigerator overnight at low temperatures, thus allowing extensive utilization of a single preparation and a reduction of experimental animals required. Taken together, we think that the isolated frog CNS is well suited for a variety of neuroanatomical standard procedures and, furthermore, may bridge the gap between isolated cellular physiology and the analysis of complex brain functions.

The present study shows that an isolated anuran CNS preparation has many advantages. Adequate physiological integrity can be maintained without perfusing the vascular system as is necessary in isolated mammalian brains (e.g., Llinás et al., 1981; Mühlethaler et al., 1993). Therefore, no elaborate support strategy is necessary to reduce the effects due to absence of a blood supply. Isolated CNS preparations of anuran brains can be used with equal success and ease as the lamprey (e.g., Brodin and Grillner, 1990) and turtle (e.g., Hounsgaard and Nicholson, 1990; Keifer et al., 1992) preparations.

#### ACKNOWLEDGEMENTS

We are especially grateful to Drs. Norbert Dieringer, Hans Straka and Andreas Kunkel for introducing us into the preparation of the isolated brain. Thanks are due to Drs. Wolfgang Wiggers and Joachim Strake for additional technical details, and to Mrs. Marlies Grosmann and Mrs. Roelie de Boer-van Huizen for technical assistance. This work was supported by a DFG grant (Wa 446/3-2), a grant of the Spanish Government (DGICYT PB 93-0083) and a NATO Collaborative Research Grant (930542) to H.J. ten Donkelaar.



Figure 1: A, Recording chamber with Sylgard-covered bottom; B, Recording situation for electrophysiology: an isolated brain preparation of *Discoglossus pictus* is stimulated at a single branchlet of the auditory nerve via a suction electrode (1), the recording electrode (2) is lowered into the midbrain.



Figure 2: Examples of HRP, biocytin and BDA labeling in *in vitro* preparations. A, Retrogradely labeled neurons in the contralateral caudal and dorsal medullary nucleus as well as anterogradely labeled structures after application of HRP into the dorsal medullary nucleus of *Discoglossus pictus*, x175; B, Retrogradely labeled neurons in the contralateral dorsal medullary and vestibular nucleus after application of biocytin into the dorsal medullary and vestibular nucleus after application of biocytin into the dorsal medullary area of *D. pictus*, x175; C, Anterogradely labeled fibers and terminal structures in the contralateral principal nucleus of the torus semicircularis after application of biocytin into the dorsal medullary area of *D. pictus*, x175; D, Retrogradely labeled neuron in the contralateral reticular formation of *Xenopus laevis* at the obex level after application of 3 kD BDA to the torus semicircularis, x190.



Figure 3: A, A representative experiment showing the application of 3 kD BDA to the torus semicircularis of *Xenopus laevis*; B, Example of labeling in the dorsal medullary nucleus, x170.



Figure 4: Double-labeled fluorescence preparations of *Discoglossus pictus*. A, Double labeling in the lateral part of the midbrain tegmentum. Red fluorescence: neurons of the lateral lemniscal nucleus retrogradely labeled with RDA from the contralateral torus; green fluorescence: fibers of the lateral lemniscus labeled with FDA from the ipsilateral torus, x250. B, Double labeling in the laminar nucleus of the torus semicircularis. Red fluorescence: neurons retrogradely labeled with RDA from the superior olive; green fluorescence: leucine-enkephalin-like immunoreactivity in the laminar nucleus, x250.



Figure 5: Intracellularly labeled neuron in the magnocellular nucleus of the torus semicircularis of *Discoglossus pictus*. Section thickness 20  $\mu$ m, x225.

#### REFERENCES

Adams, J.C. (1981) Heavy metal intensification of DABbased HRP reaction product, J. Histochem. Cytochem., 29: 775.

Atzori, M. and Nistri, A. (1994) Non-monotonic decay of excitatory synaptic transmission in the frog optic tectum following repetitive stimulation of the optic nerve in vitro, Exp. Brain Res., 102: 287-296.

Boulton, C.L., von Haebler, D. and Heinemann, U. (1992) Tracing of axonal connections by rhodaminedextran-amine in the rat hippocampal-entorhinal cortex slice preparation, Hippocampus, 2: 99-106.

Brodin, L. and Grillner, S. (1990) The lamprey CNS in vitro, an experimentally amenable model for synaptic transmission and integrative functions. In: H. Jahnsen (Ed.) Preparations of Vertebrate Central Nervous System *In Vitro*, Wiley, Chichester, pp. 103-153.

Cochran, S.L., Kasik, P. and Precht, W. (1987) Pharmacological aspects of excitatory synaptic transmission to second-order vestibular neurons in the frog, Synapse, 1: 102-123. Connors, B.W. and Kriegstein, A.R. (1986) Cellular

Connors, B.W. and Kriegstein, A.R. (1986) Cellular physiology of the turtle visual cortex: Distinctive properties of pyramidal and stellate neurons, J. Neurosci., 6: 164-177.

Diamond, J.T., Conley, M., Fitzpatrick, D. and Raczkowski, D. (1991) Evidence for seperate pathways within the tecto-geniculate projection in the tree shrew, Proc. Natl. Acad. Sci., 88: 1315-1319.

Dicke, U. and Roth, G. (1994) Tectal activation of premotor and motor networks during feeding in salamanders, Eur. J. Morphol., 32: 106-116.

Feng, A.S. and Lin, W. (1991) Differential innervation patterns of three divisions of frog auditory midbrain (torus semicircularis), J. Comp. Neurol., 306: 613-630.

Fritzsch, B. (1993) Fast axonal diffusion of 3000 molecular weight dextran amines, J. Neurosci. Meth., 50: 95-103.

Fritzsch, B. and Wilm, C. (1990) Dextran amines in neuronal tracing, Trends Neurosci., 13: 14.

Glover, J.C., Petursdottir, G. and Jansen, J.K.S. (1986) Fluorescent dextran amines used as axonal tracers in the nervous system of the chick embryo, J. Neurosci. Meth., 18: 243-254.

González, A. and Muñoz, M. (1987) Some connections of the area octavolateralis of *Pleurodeles waltlii*. A study with horseradish peroxidase under in vitro conditions, Brain Res., 423: 338-342.

Görcs, T., Antal, M., Olah, E. and Székely, G. (1979) An improved cobalt labelling technique with complex compounds, Acta Biol. Acad. Sci. Hung., 30: 79-86.

Holohean, A.M., Hackman, J.C. and Davidoff, R.A. (1990) An *in vitro* study of the effects of serotonin on frog primary afferent terminals, Neurosci. Lett., 113: 175-180.

Hounsgaard J. and Nicholson, C. (1990) The isolated turtle brain and the physiology of neuronal circuits. In: H. Jahnsen (Ed.) Preparations of Vertebrate Central Nervous System *In Vitro*, Wiley, Chichester, pp. 279-294.

Izzo, P.N. (1991) A note on the use of biocytin in anterograde tracing studies in the central nervous system - application at both light and electron microscopic level, J. Neurosci. Meth., 36: 155-166.

Kahn, J.A. and Roberts, A. (1982) Experiments on the central pattern generator for swimming in amphibian embryos, Phil. Trans. Roy. Soc. (Lond.) B, 296: 229-243.

Kawaguchi, Y., Wilson, C.J. and Emson, P.C. (1989) Intracellular recording of identified neostriatal patch and matrix spiny cells in a slice preparation preserving cortical inputs, J. Neurophysiol., 62: 1052-1068.

Keifer, J. and Houk, J.C. (1989) An *in vitro* preparation for studying motor pattern in the cerebellorubrospinal circuit of the turtle, Neurosci. Lett., 97: 123-128.

Keifer, J., Vyas, D. and Houk, J.C. (1992) Sulforhodamine labeling of neural circuits engaged in motor pattern generation in the *in vitro* turtle brainstemcerebellum, J. Neurosci., 12: 3187-3199.

Kenan-Vaknin, G., Katz, H. and Malach. R. (1992) Golgi-like staining of visual cortex cells obtained by extracellular biocytin application *in vitro*, Brain Res., 571: 338-341.

King, M.A., Louis, P.M., Hunter, B.E. and Walker, D.W. (1989) Biocytin: A versatile anterograde neuroanatomical tract-tracing alternative, Brain Res., 497: 361-367.

Kita, H. and Armstrong, W. (1991) A biotin-containing compound N-(2-aminoethyl)biotinamide for intracellular labeling and neuronal tracing studies - comparison with biocytin, J. Neurosci. Meth., 37: 141-150.

Kriegstein, A.R. and Connors, B.W. (1986) Cellular physiology of the turtle visual cortex: Synaptic properties and intrinsic circuitry, J. Neurosci., 7: 2488-2492.

Lapper, S.R. and Bolam, J.P. (1991) The anterograde and retrograde transport of neurobiotin in the central nervous system of the rat - comparison with biocytin, J. Neurosci. Meth., 392: 163-174.

Larson-Prior, L.J., Ulinski, P.S. and Slater, N.T. (1991) Excitatory amino acid receptor-mediated transmission in geniculocortical and intracortical pathways within visual cortex, J. Neurophysiol., 66: 293-306.

Llinás, R., Yarom, Y. and Sugimori, M. (1981) Isolated mammalian brain in vitro: New technique for analysis of electrical activity of neuronal circuit function, Fed. Proc., 40: 2240-2245.

Luksch, H. and Walkowiak, W. (1992) Distribution of neurotransmitter-like immunoreactivities in the torus semicircularis of three anuran species. In: N. Elsner and D.W. Richter (Eds.), Proc. 20th Göttingen Neurobiol. Conf., Thieme, Stuttgart-New York, p. 201.

Luksch, H. and Walkowiak, W. (1993) Comparison of intracellular *in vivo* and *in vitro* recordings in the auditory midbrain of the frog, Discoglossus pictus. In: N. Elsner and M. Heisenberg (Eds.), Proc. 21th Göttingen Neurobiol. Conf., Thieme, Stuttgart-New York, p. 233.

Lutz, P.L., Rosenthal, M. and Sick, T.J. (1985) Living without oxygen: Turtle brain as a model of anaerobic metabolism, Mol. Physiol., 8: 411-525.

McCormick, C.A. and Braford, M.R. (1984) A horseradish peroxidase procedure for tracing the central projections of peripheral nerves in exsanguinated anamniotes, Neurosci. Lett., 51: 207-212.

McLean, H.A., Kimura, N., Kogo, N., Perry, S.F. and Remmers, J.E. (1995) Fictive respiratory rhythm in the isolated brainstem of frogs, J. Comp. Physiol. A, 176: 703-713.

Mesulam, M.-M. (1982) Tracing Neuron Connections with Horseradish Peroxidase. Wiley, Chichester.

Møllgård, K., Balslev, Y., Stagaard-Janas, M., Treherne, J.M., Saunders, N.R. and Nicholls, J.G. (1994) Development of spinal cord in the isolated CNS of a neonatal mammal (the opossum *Monodelphis domestica*) maintained in longterm culture. J. Neurocytol., 23: 151-165.

Mühlethaler, M., de Curtis, M., Walton, K. and Llinás, R. (1993) The isolated and perfused brain of the guineapig *in vitro*, Eur. J. Neurosci., 5: 915-926.

Muñoz, A., Muñoz, M., González, A and ten Donkelaar, J.J. (1995) The anuran dorsal column nucleus: Organization, immunohistochemical chracterization and fiber connections in *Rana perezi* and *Xenopus laevis*, J. Comp. Neurol., in press.

Nance, D.M. and Burns, J (1990) Fluorescent dextrans as sensitive anterograde neuroanatomical tracers: Applications and pitfalls, Brain Res. Bull., 25: 139-145.

Nicholls, J.G., Stewart, R.R., Erulkar, S.D. and Saunders, N.R. (1990) Reflexes, fictive respiration and cell division in the brain and spinal cord of the newborn opossum, *Monodelphis domestica*, J. Exp. Biol., 152: 1-15.

Norgren, R.B. and Lehmann, M.N. (1990) A new chromogen for use in HRP-tract tracing and double-label immunocytochemistry, Brain Res. Bull., 25: 393-396.

Popov, S. and Poo, M.-M. (1992) Diffusional transport of macromolecules in developing nerve processes, J. Neurosci., 12: 77-85.

Roberts, A. and Clarke, J.D.W. (1982) The neuroanatomy of an amphibian embryo spinal cord, Phil. Trans. Roy. Soc. (Lond.) B, 296: 195-212.

Roberts, A., Soffe, S.R. and Dale, N. (1986) Spinal interneurons and swimming in frog embryos. In: S. Grillner, R. Herman, P.S.G. Stein and D. Stuart (Eds), Neurobiology of Vertebrate Locomotion, MacMillan, London, pp. 279-306

Rovainen, C.M. (1967a) Physiological and anatomical studies on large neurons of central nervous system of the sea lamprey (*Petromyzon marinus*). I. Müller and Mauthner cells, J. Neurophysiol., 30: 1000-1023.

Rovainen, C.M. (1967b) Physiological and anatomical studies on large neurons of central nervous system of the sea lamprey (*Petromyzon marinus*). II. Dorsal cells and giant interneurons, J. Neurophysiol., 30: 1024-1042.

Sagawa, S., Oka, J.I. and Fukuda, H. (1987) Descending and segmental spinal pathways differently regulate contractions of antagonistic muscles of frogs *in vitro*, Comp. Biochem. Physiol., 2: 437-441.

Sarrafizadeh, R. and Houk, J.C. (1994) Anatomical organization of the limb premotor network in the turtle (*Chrysemys picta*) revealed by in vitro transport of biocytin and neurobiotin, J. Comp. Neurol., 344: 137-159.

Schaffer, D.D. (1982) Comparative and descriptive aspects of neural correlates of mating calling in the isolated brainstem of Rana p. pipiens, Comp. Biochem. Physiol., 72: 371-375.

Schmidt, R.S. (1976) Neural correlates of frog calling. Isolated brain stem, J. Comp. Physiol., 108: 99-113. Schulte-Mattler, M. and Luhmann, H.J. (1995) Dyecoupling reveals intrinsic microcircuitry in rat neocortex. In: N. Elsner and R. Menzel (Eds.), Proc., 23rd Göttingen Neurobiol. Conf., Thieme, Stuttgart-New York, p. 559.

Shu, S., Ju, G. and Fan, L. (1988) The glucose oxidase-DAB-nickel method in peroxidase histochemistry of the nervous system, Neurosci. Lett., 85: 169-171.

Simpson, I.B., Rose, R.W. and Loewenstein, W.R. (1977) Size limits of molecules permeating the junctional channel, Science 195: 294-296.

Smith, J.C. and Feldman, J.L. (1987) In vitro brainstemspinal cord preparations for study of motor systems for mammalian respiration and locomotion, J. Neurosci, Meth., 21: 321-333.

Straka, H. and Dieringer, N. (1991) Internuclear neurons in the ocular motor system of frogs, J. Comp. Neurol., 312: 537-548.

Straka, H. and Dieringer, N. (1993) Electrophysiological and pharmacological characterization of vestibular inputs to identified frog abducens motoneurons and internuclear neurons *in vitro*, Eur. J. Neurosci., 5: 251-260.

Székely, G. and Gallyas, F. (1975) Intensification of cobaltous sulphide precipitate in frog nervous tissue. Acta Biol. Acad. Sci. Hung., 26: 175-188.

Tao, L. and Nicholson, C. (1992) Quantitative optical imaging of the diffusion of dextrans of different molecular weights in rat cerebral cortex, Soc. Neurosci. Abstr., 18: 967.

Veenman, C.L., Reiner, A. and Honig, M.G. (1992): Biotinylated dextran amine as an anterograde tracer for single-labeling and double-labeling studies, J. Neurosci. Meth., 41: 239-254.

Walkowiak, W. and Luksch, H. (1994) Sensory motor interfacing in acoustic behavior of anurans, Amer. Zool., 34: 153-173.

Wallén, P., Grillner, S., Feldman, J. and Bergelt, S. (1985) Dorsal and ventral myotome motoneurons and their input during fictive locomotion in lamprey, J. Neurosci., 5: 654-661.

Wheatley, M. and Stein, R.B. (1992) An *in vitro* preparation of the mudpuppy for simultaneous intracellular and electromyographic recording during locomotion, J. Neurosci. Meth., 42: 129-138.

Wiggers, W. and Roth, G. (1994) Depth perception in salamanders: The wiring of visual maps, Eur. J. Morphol., 32: 311-314.

Wilczynsky, W. (1981) Afferents to the midbrain auditory center in the bullfrog, Rana catesbeiana, J. Comp. Neurol., 198: 421-433.

Will, U., Lühede, G. and Görner, P. (1985) The area octavo-lateralis in *Xenopus laevis*. II. Second order projections and cytoarchitecture, Cell Tiss. Res., 239: 147-161.

The use of <u>in vitro</u> preparations of the isolated amphibian central nervous system in neuroanatomy and electrophysiology

**COMENTARIOS** 

2.2

En el presente capítulo se describe, en diversas especies de anfibios, un protocolo para la realización de una preparación subperfundida con solución de Ringer oxigenada, en la que el sistema nervioso central (SNC) completo, y aislado del cuerpo del animal, puede mantenerse vivo durante varios días, permitiendo así la realización de estudios de trazado neuronal, con aplicaciones de trazadores tanto extracelular como intracelularmente; inmunohistoquímicos y electrofisiológicos.

La viabilidad de las preparaciones in vitro del SNC depende de aspectos como las estrategias de disminución de las necesidades metabólicas, así como el abastecimiento de oxígeno y sustratos metabólicos, que reducen los efectos causados por la ausencia del riego sanguíneo. En la mayoría de los tejidos de mamíferos no es posible mantener una integridad fisiológica adecuada, sin perfundir el sistema vascular con algún tipo de sustituto de la sangre. Un ejemplo de esto es la preparación del cerebro aislado de cobaya (Llinás y cols., 1981; Mühlethaler y cols., 1993). Sin embargo, las preparaciones aisladas del SNC neonatal de la zarigüella (Nicholls y cols., el 1990; Møllgård y cols., 1994), o preparaciones de tronco cerebral-médula espinal (Smith y Feldman, 1987), pueden mantenerse vivas sin perfusión del sistema vascular.

En vertebrados no mamíferos se han obtenido preparaciones *in vitro* viables para el estudio de partes del SNC, debido a características particulares. Así, en el caso del cerebro de la tortuga, al presentar una resistencia inusual a la anoxia (Lutz y cols., 1985; Hounsgaard y Nicholson, 1990), se han podido conseguir fácilmente preparaciones *in vitro*  (Connors y Kriegstein, 1986; Kriegstein y Connors, 1986; Keifer y Houk, 1989; Larson-Prior y cols., 1991 Keifer y cols., 1992; Sarrafizadeh y Houk, 1994). En la lamprea la médula es delgada, carece de vasos sanguíneos intrínsecos, y es oxigenada directamente desde el fluído cerebroespinal; otras regiones, como el tronco cerebral, presentan vasos sanguíneos intrínsecos, pero probablemente son también oxigenadas en gran medida desde el fluído cerebroespinal (Brodin y Grillner, 1990). Todo ello favorece la realización de preparaciones *in vitro* viables (Rovainen, 1967a,b; Wallén y cols., 1985; Brodin y Grillner, 1990).

En anfibios se han descrito diversos protocolos para el estudio in vitro del SCN. incluyendo la utilización de una preparación subperfundida de embriones de Xenopus laevis (Kahn y Roberts, 1982; Roberts y Clarke, 1982; Roberts y cols., 1986), de secciones de cerebros (Holohean y cols., 1990), y de tronco cerebral (Schmidt, 1976; Schaffer 1982; Cochran y cols., 1987; Straka y Dieringer, 1993; Atzori y Nistri, 1994; Dicke y Roth, 1994; McLean y cols., 1995), así como preparaciones combinadas de médula espinal y musculatura (Sagawa y cols., 1987; Wheatley y Stein, 1992). Hasta el momento, el uso de preparaciones del SNC completo y aislado, para experimentos de trazado neuronal en anuros, se ha limitado a la utilización de la técnica de la peroxidasa de rábano (HRP) in vitro, en la que se emplean cerebros perfundidos y consecutivamente fijados (McCormick y Braford, 1984; González y Muñoz, 1987), y del método de marcaje con cobalto en cerebros fijados (Székely y Gallyas, 1975; Görcs y cols., 1979). Straka y Dieringer (1991) utilizaron una preparación de tronco cerebral y médula espinal aislados para trazado neuronal con HRP.

En el presente trabajo se han empleado técnicas de trazado neuronal, mediante la aplicación extracelular de distintos trazadores en forma de cristal, y de inyecciones de soluciones acuosas. En general, las características observadas in vitro en cuanto a la captación del trazador, y a la direccionalidad y velocidad de su transporte son, en su mayor parte, comparables a las descritas mediante la aplicación in vivo de HRP (Adams, 1981; Mesulam, 1982), biocitina (King y cols., 1989; Diamond y cols., 1991; Izzo, 1991; Lapper y Bolam, 1991; Kenan-Vaknin y cols., 1992) y dextrano aminas con pesos moleculares de 3 kD o 10 kD, biotiniladas o conjugadas a diferentes fluorocromos, como tetrametilrodamina o fluoresceina (Glover y cols., 1986; Fritzsch y Wilm, 1990; Nance y Burns, 1990). Debido a que las dextrano aminas, de bajo peso molecular (3 kD), combinadas con biotina o con sustancias fluorescentes, difunden más rápidamente que las de alto peso molecular (10 kD) (Popov y Poo, 1992; Tao y Nicholson, 1992; Fritzsch, 1993), su utilización resulta sumamente útil en la preparación in vitro. Así, con la dextrano amina de 3kD combinada con biotina, se logró un rápido marcaje neuronal completo, incluyendo dendritas secundarias y terciarias.

Mediante la aplicación intracelular de trazadores *in vitro* se consiguen los mismos resultados que con las aplicaciones *in vivo*. Así con biocitina, ocasionalmente, observamos conjuntos de neuronas, simultáneamente marcadas, con patrones similares de morfologías dendríticas y axonales, probablemente debido al acoplamiento de dichas células por medio de uniones de tipo "gap" (Simpson y cols., 1977) a través de las cuales difunde la biocitina, más que a una aplicación inespecífica de ésta. Estos resultados, sin embargo, no pueden atribuirse a la preparación de cerebro aislado, debido a que se han descrito datos similares en estudios realizados *in vivo* (Kawaguchi y cols., 1989; Kita y Armstrong, 1991; Wiggers y Roth, 1994; Schulte-Mattler y Luhmann, 1995).

En los experimentos neuroanatómicos realizados en el presente trabajo, no hemos encontrado ninguna limitación técnica, en cuanto a la aplicación de los trazadores *in vitro*, ni signos de degeneración celular, incluso después de varios días en la solución de Ringer. Igualmente, no hemos observado diferencias morfológicas con respecto a los estudios realizados *in vivo*, en cuanto a las neuronas marcadas intracelularmente, las estructuras marcadas anterógrada y retrógradamente (Straka y Dieringer, 1991; Walkowiak y Luksch, 1994; Muñoz y cols., 1995), así como a su patrón de conectividad (Wilczynski, 1981; y cols., 1985; Feng y Lin, 1991).

Los resultados obtenidos en los experimentos inmunohistoquímicos, para la detección de Leu-Encefalina, y su combinación con técnicas de trazado con dextrano aminas *in vitro*, no mostraron diferencias, con los obtenidos en un estudio previo (Luksch y Walkowiak, 1992), en el que los cerebros fueron fijados, mediante la perfusión de los animales anestesiados. Esto indica que, incluso después de varios días en condiciones *in vitro*, los sistemas celulares de síntesis y transporte funcionan adecuadamente.

La estimulación eléctrica in vitro de las ramas auditivas del nervio estatoacústico condujo a la producción de respuestas selectivas en varios núcleos auditivos del tronco cerebral, incluyendo el núcleo rombencefálico dorsal, la oliva superior, diversas estructuras del tegmento mesencefálico, y el torus semicircularis; hasta cuatro días después del aislamiento del SNC, sin observarse una pérdida apreciable de actividad o una disminución del potencial de membrana de las neuronas penetradas, que mantuvieron potenciales de reposo de hasta -90 mV. El tiempo de registro más largo para una neurona individual fue de 4.5h, el cual excede considerablemente al conseguido en un registro comparable in vivo. Debido a que las aferencias al mesencéfalo incluyen por lo menos dos sinapsis (hasta cuatro), los circuitos neuronales esenciales parecen conservarse en esta preparación, la cual mantiene un buen estado fisiológico del cerebro. Además, en general no hemos observado diferencias entre los registros in vitro e in vivo (Luksch y Walkowiak, 1993).

El análisis de parámetros fisiológicos, en un cerebro aislado, tiene algunas limitaciones obvias, como la imposibilidad de excitar sistemas sensoriales mediante estímulos fisiológicos. Por otra parte, su excitación con estímulos eléctricos conduce a la estimulación de las vías sensitivas completas, y a patrones de reacciones complejas en neuronas individuales, comparables a los hallazgos en animales intactos. En algunos casos, la desaferentación completa del cerebro puede suponer una ventaja de esta preparación. *In vivo*, muchos parámetros fisiológicos del animal y el ambiente no pueden ser enteramente controlados, como por ejemplo, las variaciones debidas a las condiciones experimentales tales como el estrés causado por la inmovilización, el cambio en el abastecimiento del oxígeno, o la influencia de otras modalidades sensoriales. La preparación del cerebro aislado asegura la estabilidad de estos parámetros, ofreciendo la posibilidad de analizar la respuesta de neuronas únicas y de redes neuronales, ante una estimulación fehacientemente constante, y de manipulaciones por medio de la aplicación de agentes neurofarmacológicos o por la estimulación de distintas aferencias.

La utilización del SNC aislado ofrece numerosas ventajas para la realización de experimentos tanto anatómicos como electrofisiológicos. Todas las áreas cerebrales son fácilmente accesibles sin el inconveniente de la presencia de vasos sanguíneos que dificultan o impiden el acceso. En los experimentos anatómicos es posible realizar grandes lesiones y aplicaciones masivas de trazador, sin los problemas consecuentes de supervivencia del animal; además, los trazadores no son transportados por la circulación sanguínea. Los registros electrofisiológicos son comparativamente más fáciles de realizar y más estables que mediante la aproximación in vivo, al no producirse las pulsaciones ocasionadas por los cambios de presión de la circulación sanguínea. Igualmente, al ser posible extraer completamente las meninges, la penetración de los electrodos se facilita, y por tanto no existen limitaciones en cuanto a su forma. Esta aproximación experimental ofrece la posibilidad de trabajar sobre el mismo cerebro durante varios días, almacenando el tejido en un refrigerador durante la noche a temperaturas bajas, y permitiendo así su utilización

continuada, con la consiguiente reducción del número de animales de experimentación.

# CAPÍTULO 3

Proyecciones espinales ascendentes

- **3.1.-** Spinothalamic projections in amphibians as revealed with anterograde tracing techniques
- **3.2.-** Spinal ascending pathways in amphibians: cells of origin and main targets
- 3.3.- Comentarios

Spinothalamic projections in amphibians as revealed with anterograde tracing techniques

#### A. Muñoz, M. Muñoz, A. González and H.J. ABSTRACT ten Donkelaar\*

Department of Cell Biology, Faculty of Biology, Complutense University of Madrid, 28040 Madrid, Spain

\*Department of Anatomy and Embryology, Faculty of Medical Sciences, University of Nijmegen, Nijmegen, The Netherlands

Based on: Neuroscience Letters 171 (1994) 81-84

Key words: Spinal cord, Thalamus, Spinothalamic projections, Somatosensory systems, Amphibians, Anurans, Urodeles, Tract-tracing.

. Direct spinothalamic pathways were demonstrated in anurans (Rana ridibunda, Xenopus laevis) and in the ribbed newt, Pleurodeles waltl. With the powerful anterograde tracers Phaseolus vulgaris leucoagglutinin and biotinylated dextran amine rather extensive spinothalamic projections were found, including the ventromedial thalamic nucleus, the dorsal thalamus and several posterior diencephalic nuclei (anurans), and the neuropil lateral to the pars ventralis thalami as well as to the anteroventral and posterodorsal zones (P. waltl), respectively.

72

3.1
The presence of spinothalamic pathways appears to be a shared character in the brain of amniotes. Thus, thalamic projections from second order neurons in the spinal cord have been extensively shown for mammals<sup>25</sup>, birds<sup>18</sup>, and reptiles<sup>2,3,8</sup>. Among anamniotes, a distinct spinothalamic projection has been demonstrated only in an advanced galeomorph shark, the nurse shark, *Ginglymostoma cirratum*<sup>4</sup>, possibly a case of a non-homologous, independently evolved character<sup>17</sup>.

In amphibians so far tracing studies failed to spinothalamic projections, both in show anurans3,5,13 and in urodeles14,23. In the present study the presence (or lack) of spinothalamic projections in amphibians was studied with modern, more powerful, anterogradely transported tracers. Therefore, in the anurans Rana ridibunda and Xenopus laevis, and in the ribbed newt, Pleurodeles waltl, the tracers Phaseolus vulgaris leucoagglutinin (PHA-L) or biotinylated dextran amine (BDA) were selectively applied into the spinal cord, and the anterograde labeled fibers and terminals in the thalamus were examined. In addition, several experiments with spinal injections of horseradish peroxidase (HRP) were available for each species. Part of the results have been published in abstract form<sup>10</sup>.

The data presented are based on a total of 18 adult specimens of *Rana ridibunda*, six of *Xenopus laevis* and 10 of *Pleurodeles waltl*. In addition several BDA experiments were done in late tadpole stages of *X. laevis*. The animals were commercially obtained (*R. ridibunda* and *X. laevis*) or captured in the wild (with permission from the Spanish Government) in the surroundings of Madrid (*P. waltl*). All experiments were carried out under surgical anaesthesia with tricaine methanesulphonate (MS 222, Sandoz). In three series of experiments, the tracers HRP (6 cases), PHA-L (11 cases), and BDA (12 cases) were injected unilaterally into the cervical, thoracic (three cases) or lumbar (three tadpoles) spinal cord. In addition, in five cases, BDA was applied to the spinal cord as crystals (recrystallized from a saturated solution of the tracer in distilled water).

All injections were made iontophoretically by applying 5-8  $\mu$ A positive pulsed current (7s on/7s off) to the tracer solution (15% HRP, 2% PHA-L or 10% BDA) in a glass micropipette (outer tip diameter 20-30  $\mu$ m) for a period of 15-30 min. The animals were allowed to survive for 6-10 days in the experiments with PHA-L or BDA, and for 15-20 days in the HRP experiments. They were then reanesthetized with an overdose of MS 222 and perfused transcardially with 0.1 M phosphate buffer (pH 7.4), followed by a fixative containing 1% paraformaldehyde and 2.5% glutaraldehyde (for PHA-L and BDA experiments) or 1% paraformaldehyde and 1.25% glutaraldehyde (for HRP experiments) in 0.1 phosphate buffer. The brain and spinal cord were removed and further fixed for one to four hours in the perfusion mixture. They were then immersed in a mixture of phosphate buffer and 30% sucrose solution for three to five hours at 4<sup>a</sup>C, subsequently embedded in a solution of 15% gelatin with 30% sucrose added, and stored overnight in a 4% formaldehyde solution at room temperature. Frozen sections were cut at 40 µm thickness in the frontal plane on a freezing microtome. Histochemistry for HRP followed a heavy metal intensification of the diaminobenzidine (DAB)based HRP reaction product. For the immunostaining

for PHA-L the indirect peroxidase- antiperoxidase (PAP) technique was used, for visualizing BDA, a Vectastain ABC Elite Kit (Vector Laboratories).

The data obtained in Rana ridibunda and Xenopus laevis were largely comparable and a common pattern was found for the spinothalamic projection from the cervical spinal cord, as depicted in Fig. 1 for Rana ridibunda. The nomenclature for the different thalamic neuronal groups follows that of Neary and Northcutt<sup>12</sup> for the bullfrog, Ranacatesbeiana. Application of each of the tracers into the spinal cord at cervical levels always resulted in the labeling of ascending fiber systems, among which a small contingent of fibers is seen to arise in the dorsal gray of the spinal cord, passing ventrally to cross the midline ventral to the central canal at the same spinal levels or immediately rostral to their cells of origin. In the contralateral spinal cord the fibers turn rostrally and ascend as the spinal lemniscus in the ventrolateral funiculus. A few uncrossed fibers course rostrally in the ventral funiculus. Throughout the brainstem, most of the fibers of the spinal lemniscus terminate in rhombencephalic and mesencephalic centers. However, a small component of fibers proceeds rostrally in the ventrolateral aspect of the caudal diencephalon. Here, most fibers bend dorsomedially towards diverse targets. At caudal diencephalic levels the dorsomedial part of the nucleus of the posterior tubercle (TP) is heavily innervated by labeled fibers, some of these terminate here whereas some others continue toward the dorsal thalamus (Figs. 1D, E; 3A, B). In the ventral thalamus, the ventromedial nucleus receives the heaviest spinal projection as thin varicose fibers that cross almost all of its cell layers (Figs. 1A, B; 3C). Fibers that reach this nucleus pass through the

dorsal and ventral parts of the ventrolateral thalamic nucleus where also varicose structures were found. A small contingent of fibers arborizes within the limits of the posterior entopeduncular nucleus. In all the nuclei, their caudal aspect is always the most densely innervated. Fewer and scattered labeled fibers reach the dorsal thalamus and innervate the central, posterior and, to a lesser extent, anterior nuclei. In addition, the posterodorsal and posteroventral lateral nuclei are also scarcely innervated although numerous passing fibers cross towards the medially located dorsal thalamic nuclei. In the experiments where the application of the tracer was located in the thoracic spinal cord, similar, although less intensive, labeling was found in the thalamus, where the ventromedial nucleus receives the heaviest innervation. In three Xenopus laevis tadpole stages (stages 54, 57) a few ascending fibers from the lumbar spinal cord were found to innervate the ventral thalamus.

In the ribbed newt, Pleurodeles waltl, HRP applications to the spinal cord failed to label ascending projections as far as the diencephalon<sup>10</sup>. However, after PHA-L injections and BDA applications into the cervical spinal cord, a consistent and conspicuous labeling in the thalamus was found (Fig. 2). The different thalamic areas are termed after Wicht and Himstedt<sup>23</sup> for *Triturus alpestris*. As for anurans, ascending spinal fibers, predominantly contralateral to the injection side, ascend in the ventrolateral funiculus, pass through the brainstem, and enter the mesodiencephalic transition area. Here, the fibers form a neuropil just lateral to the cells in the pars ventralis thalami (Figs. 2B-E; 3D) that can be considered as a rostral continuation of a profuse terminal zone in the area equivalent to the torus semicircularis in the dorsal mesencephalic tegmentum. Rostrally, labeled fibers ascend up to the level of the habenular commissure (Fig. 2A). Varicose fibers distribute primarily to the internal zone of the white matter, avoiding the periventricular grey where almost all of the cells are located. Within the dorsal thalamus, a few scattered fibers terminate in the posterodorsal zone (Figs. 2D; 3E) and also in the anteroventral zone (Figs. 2C; 3F).

The present study demonstrates the presence of rather distinct, direct spinothalamic pathways in amphibians. The use of new and more powerful anterograde tracers has made it possible to identify even fine terminal fields and thin scattered fibers in the thalamus of the three species studied. Although in anterograde degeneration studies in amphibians, distinct dorsal and ventral ascending bundles were found in the ventrolateral spinal funiculus, the most rostrally located targets of ascending spinal pathways were found in the mesencephalic tectum in the  $axolotl^{14}$  and in the midbrain tegmentum of anurans $^{3,5}$ , respectively. The lack of degenerating fibers or terminals in thalamic areas only confirmed other studies 5,17 in various anamniotes in which spinothalamic projections could not be observed. With the sole exception of an advanced galeomorph shark (the nurse shark, Ginglymostoma cirratum) in which spinal projections were found to reach the thalamus<sup>4</sup>. it seemed likely that spinothalamic pathways evolved in amniotes only.

In anurans, however, electrophysiological studies<sup>20</sup> suggested a bilateral processing of somatic information all along the rostrocaudal extent of the dorsal thalamus. In the telencephalon several areas respond to somatic stimuli including the dorsal

pallium, the medial pallium, the septum and the striatum<sup>15,20</sup>. All these telencephalic centers have been identified to receive thalamic information from the various nuclei that respond to somatic signals. Thus, the medial pallium, and the dorsal pallium are innervated by the anterior thalamic nucleus<sup>7,11,16,21</sup>. The striatum receives thalamic information from the central dorsal thalamus<sup>22,24</sup>, and this thalamic region can be viewed as an intermediate in the transition of somatic information to the striatum. The present study shows that, apart from somatosensory information relayed to the thalamus via the dorsal column nucleus<sup>9,13</sup>, a direct channel from the spinal cord is also available in anurans.

In urodeles, relatively few experimental data are available on ascending somatosensory pathways. In a pioneer study in *Ambystoma tigrinum*, Herrick<sup>6</sup> suggested a spinal innervation of the medial zone of the sensory dorsal thalamus. Anterograde degeneration<sup>14</sup> or HRP<sup>23</sup> studies could not confirm Herrick's ideas. In the present study in *Pleurodeles waltl* a small projection to the posterodorsal and anteroventral zones of the dorsal thalamus was shown. Both zones have been demonstrated to project to the striatum and the medial pallium, respectively<sup>1,19,23</sup>. Therefore, in urodeles as in anurans, various dorsal thalamic structures might serve as relay stations for somatosensory information from the spinal cord to the telencephalon.

The present study was supported by a Grant from the Spanish Government (DGICYT PB 90-0628) to A. González and a NATO Collaborative Research Grant (930542) to H.J. ten Donkelaar. The authors would like to thank Mrs. Ine Bergervoet-Vernooy for technical assistance, and Mrs. Marion van de Coevering for help in preparing the electronic manuscript.



Figure 1: Camera lucida drawings of transverse sections through the diencephalon from rostral (A) to caudal (E) for a representative experiment with BDA application to the cervical spinal cord (shaded areas in F) of *Rana ridibunda*. Abbreviations: A, C, anterior and central thalamic nuclei; cho, chiasma opticum; La, Lpd, Lpv, lateral thalamic nucleus, anterior, posterodorsal and posteroventral divisions; Mg, magnocellular preoptic nucleus; NPv, nucleus of the periventricular organ; P, posterior thalamic nucleus; SC, suprachiasmatic nucleus; tm, tectum mesencephali; TP, nucleus of the posterior tuberculum; VH, ventral hypothalamic nucleus; Vld, Vlv, ventrolateral thalamic nucleus, dorsal and ventral parts; VM, ventromedial and thalamic nucleus.



a -

Figure 2: Diagrams of transverse sections through the diencephalon from rostral (A) to caudal (E) for a representative experiment with a PHA-L injection into the spinal cord of (shaded area in F). Abbreviations: cho, chiasma opticum; hb, habenula; pop, nucleus preopticus, pars posterior; pvt, pars ventralis thalami; zav, zona anteroventralis; zpd, zona posterodorsalis.



Figure 3: Photomicrographs of anterogradely BDA or PHA-L labeled fibers in *Rana ridibunda* (A-C) and *Pleurodeles waltl* (D-F). A: a bundle of fibers ascending toward the posterior tubercle; B, C, varicose fibers progressing dorsally to innervate the central (B) and ventromedial (C) thalamic nuclei; D: a dense arborization in the pars ventralis thalami contralateral side; E, F, a few, scattered fibers that reach the zona posterodorsalis or the zona anteroventralis in both the contralateral (E) and ipsilateral (F) side. Scale bars = 140  $\mu$ m (A-C), 50  $\mu$ m (D), 200  $\mu$ m (E), and 100  $\mu$ m (F). For abbreviations see Figs. 1 and 2.

#### REFERENCES

- 1 Dubé, L., Clairambault, P. and Malacarne, G., Striatal afferents in the newt *Triturus cristatus*, Brain Behav. Evol., 35 (1990) 212-226.
- 2 Ebbesson, S.O.E., Ascending axon degeneration following hemisection of the spinal cord in the tegu lizard (*Tupinambis nigropunctatus*), Brain Res., 5 (1967) 178-206.
- 3 Ebbesson, S.O.E., Brain stem afferents from the spinal cord in a sample of reptilian and amphibian species, Ann. N.Y. Acad. Sci., 167 (1969) 80-101.
- 4 Ebbesson, S.O.E. and Hodde, K.C., Ascending spinal systems in the nurse shark *Ginglymostoma* cirratum, Cell Tissue Res., 216 (1981) 313-331.
- 5 Hayle, T.H., A comparative study of spinal projections to the brain (except cerebellum) in three classes of poikilothermic vertebrates, J. Comp. Neurol., 149 (1973) 463-476.
- 6 Herrick, C.J., The cerebrum of Amblystoma tigrinum in midlarval stages, J. Comp. Neurol., 70 (1939) 249-266.
- 7 Kicliter, E., Some telencephalic connections in the frog, Rana pipiens, J. Comp. Neurol., 185 (1979) 75-86.
- 8 Künzle, H. and Woodson, W., Mesodiencephalic and other target regions of the ascending spinal projections in the turtle: *Pseudemys scripta elegans*, J. Comp. Neurol., 212 (1982) 349-364.
- 9 Muñoz, A., de Boer-van Huizen, R., Bergervoet-Vernooy, I. and ten Donkelaar, H.J., Early development of dorsal column-medial lemniscal projections in the clawed toad, *Xenopus laevis*, *Dev. Brain Res.*, 74 (1993) 291-294.
- 10 Muñoz, M., Muñoz, A., González, A., Machín, C. and Rúa, C., Ascending projections from the spinal cord in the anuran amphibian *Rana ridibunda*, *Eur.* J. Neurosci. Suppl., 4 (1991) 24.
- Neary, T.J., Anterior thalamic nucleus projections to the dorsal pallium in ranid frogs, *Neurosci. Lett.*, 51 (1984) 213-218.
- 12 Neary, T.J. and Northcutt, R.G., Nuclear organization of the bullfrog diencephalon, J. Comp. Neurol., 213 (1983) 262-278.
- 13 Neary, T.J. and Wilczynski, W., Ascending thalamic projections from the obex region in ranid frogs, *Brain Res.*, 138 (1977) 529-533.
- 14 Nieuwenhuys, R. and Cornelisz, M., Ascending projections from the spinal cord in the axolotl (Ambystoma mexicanum), Anat. Rec., 169 (1971) 388.
- 15 Northcutt, R.G., Pallial projection of sciatic, ulnar and mandibular branch of the trigeminal afferents in the frog (*Rana catesbeiana*), Anat. Rec., 166 (1970) 356.
- 16 Northcutt, R.G. and Ronan, M., Afferent and efferent connections of the bullfrog medial pallium, *Brain Behav. Evol.*, 40 (1992) 1-16.
- 17 Ronan, M.C. and Northcutt, R.G., Projections ascending from the spinal cord to the brain in petromyzontid and myxinoid Agnathans, J. Comp. Neurol., 291 (1990) 491-508.
- 18 Schneider, A. and Necker, R., Spinothalamic projection in the pigeon, *Brain Res.*, 484 (1989) 139-149.

- 19 Sassoè Pognetto, M., Pairault, C., Clairambault P. and Fasolo, A., The connections of the anterior pallium in *Pleurodeles waltl* and *Triturus carnifex*: an HRP study, J. Hirnforsch., 32 (1991) 397-407.
- 20 Vesselkin, N.P., Agayan, A.L. and Nomokonova, L.M., A study of thalamo-telencephalic afferent systems in frogs, *Brain Behav. Evol.*, 4 (1971) 295-306.
- 21 Vesselkin, N.P. and Ermakova, T.V., Studies on the thalamic connections in the frog *Rana temporaria* by the peroxidase technique, *J. Evol. Biochem. Physiol.*, 14 (1978) 117-122.
- 22 Vesselkin, N.P., Ermakova, T.V., Kenigfest, N.B. and Goikovic, G., The striatal connections in frog *Rana temporaria*: an HRP study. J. Hirnforsch., 21 (1980) 381-392.
- 23 Wicht, H. and Himstedt, W., Topologic and connectional analysis of the dorsal thalamus of *Triturus alpestris*, J. Comp. Neurol., 267 (1988) 545-561.
- 24 Wilczynski, W. and Northcutt, R.G., Connections of the bullfrog striatum: afferent organization, J. Comp. Neurol., 214 (1983) 321-332.
- 25 Willis, W.D. and Coggeshall, R.E., Sensory Mechanisms of the Spinal Cord, 2nd edn., Plenum, New York, 1991.

Spinal ascending pathways in amphibians: cells of origin and main targets

A. Muñoz<sup>1,2</sup>, M. Muñoz<sup>1</sup>, A. González<sup>1</sup> and H.J. ten Donkelaar<sup>2</sup>

<sup>1</sup>Department of Cell Biology, Universidad Complutense de Madrid (Spain).

<sup>2</sup>Department of Anatomy and Embryology, University of Nijmegen, Nijmegen (The Netherlands)

Based on: J. Comp. Neurol. (Submitted)

Key words: evolution, somatosensory systems, ascending sensory pathways, thalamus, torus semicircularis

### ABSTRACT

As part of a research programme on the evolution of somatosensory systems in vertebrates, the various components of ascending spinal projections were studied with *in vivo* and *in vitro* tract-tracing techniques in representative species of amphibians (the large green frog, *Rana perezi*, the clawed toad, *Xenopus laevis*, and the ribbed newt *Pleurodeles waltl*). Three main ascending sensory channels were demonstrated:

1) Ascending projections via the *dorsal* funiculus include primary and non-primary projections that ascend to terminate mainly in the dorsal column nucleus at obex levels. A small component ascends farther rostralwards to terminate in the reticular formation, the octavolateral area, the trigeminal nuclear complex and in the granular layer of the cerebellum.

2) Projections ascending via the *dorsolateral* funiculus reach other spinal and supraspinal targets

3.2

than the dorsal funicular fibers, mainly ipsilaterally. At upper cervical cord and obex levels, many fibers innervate a region considered the amphibian homologue of the lateral cervical nucleus of mammals. In the medulla, these fibers ascend ventral to the descending trigeminal tract to terminate in the dorsal column and the solitary tract nuclei, and more rostrally, in the reticular formation, the descending trigeminal nucleus, and the medial aspect of the ventral octaval nucleus. Major projections reach the area between the facial motor nucleus and the ventral octaval nucleus, and a mediolateral subcerebellar band. These projections arise in neurons located mainly in the ipsilateral deep dorsal and lateral fields throughout the spinal cord.

3) Ascending spinal projections via the ventral quadrant of the spinal cord (the ventral and ventrolateral funiculi) ascend throughout the brainstem up to the diencephalon. Along its course this component innervates various parts of the reticular formation, the octavolateral area, the granular layer of the cerebellum, the region ventromedial and ventrolateral to the isthmic nucleus and the subcerebellar region. In the mesencephalon, the torus semicircularis, the midbrain tegmentum and, sparsely, the tectum mesencephali are innervated. Beyond the midbrain various dorsal and particularly ventral thalamic nuclei and the posterior tubercle are innervated by this ascending sensory channel. The cells of origin of some of these projections were observed in the dorsal, and to a lesser extent, in the lateral and ventral spinal fields of the spinal cord.

Evidence for the presence of these three main ascending sensory channels throughout vertebrates will be discussed. The presence of such channels appears to be a shared character in the brain of both amniotes and anamniotes.

#### INTRODUCTION

In terrestrial vertebrates, two basic systems of ascending spinal projections are found (see Willis and Coggeshall, 1991): 1) a primary afferent ascending spinal projection via the dorsal funiculus to the dorsal column nucleus, and 2) a secondary afferent projection via the lateral funiculus to the reticular formation, mesencephalon and thalamus. Recent studies in amphibians (A.Muñoz et al., 1994b, 1995 a,b) show that the dorsal column nucleus also receives nonprimary spinal afferents, and that the dorsolateral funiculus innervates an anuran homologue of the mammalian lateral cervical nucleus. Both the dorsal column nucleus and the lateral cervical nucleus innervate the contralateral thalamus via the medial lemniscus. Moreover, distinct spinothalamic projections are present in amphibians (A. Muñoz et al., 1994a). These studies suggest that the classical subdivision of ascending spinal projections into two systems is too simple.

The present study will show that *three* main ascending sensory channels are present in vertebrates: an ipsilateral projection via the dorsal funiculus to the dorsal column nucleus, a mainly ipsilateral projection via the dorsolateral funiculus to a lateral cervical nucleus and various rhombencephalic centers, and mainly contralateral projections via the ventral and ventrolateral funiculi (the ventral quadrant of the spinal cord) to the brainstem and the thalamus. The ascending projections that course in the various funiculi of the spinal cord were studied with modern tract-tracing techniques in three representative species of amphibians: two anuran species, the Spanish large green frog, Rana perezi (formely R. ridibunda), and the South African clawed toad, Xenopus laevis, and one urodele species, the ribbed newt, Pleurodeles waltl. The targets of ascending spinal fibers were studied with anterograde tracers (Phaseolus vulgarisleucoagglutinin, horseradish peroxidase and biotinylated dextran amine). The cells of origin of some of these ascending sensory pathways were analyzed by retrograde labeling with HRP and BDA. The latter part of this study was done on an in vitro preparation of the anuran central nervous system, i.e. an isolated brain preparation (Luksch et al., 1996). It will be shown that the presence of three ascending sensory channels is common to vertebrates, and is a shared character in the brain of both amniotes and anamniotes.

#### MATERIALS AND TECHNIQUES

The present study is based on data obtained in the anuran species *Rana perezi* and *Xenopus laevis* and in the urodele *Pleurodeles waltl*. A total number of 20 adult *R. perezi*, 9 adult and 10 young adult *X. laevis* and 12 adult *P. waltl* were used. The animals were obtained from laboratory stock of the Department of Cell Biology, University Complutense of Madrid (*R. perezi* and *P. waltl*), and the Department of Animal Physiology, University of Nijmegen (*X. laevis*). In vivo and in vitro approaches were used.

In vivo tract-tracing experiments. All experiments were carried out under surgical anesthesia with MS 222 (Sigma). The anterograde tracer *Phaseolus vulgaris*-leucoagglutinin (PHA-L, Sigma) as well as the bidirectionally transported tracers horseradish peroxidase (HRP, Boehringer), and biotinylated dextran amine (BDA 10kD, Molecular Probes) were applied to the dorsal horn of various spinal segments. Tracer solutions (a 10% HRP; a 2% PHA-L or a 10% BDA solution in 0.1M phosphate buffer-PB-pH 7.4) were iontophoretically injected during 5-10 min using a 5-10 µA positive pulled current (7 s on/7 s off) at cervical (brachial), thoracic or lumbar spinal cord levels in R. perezi and X. laevis. Cervical injections were made in *P. waltl.* In another set of experiments, BDA was recrystallized from distilled water onto fine sharp tungsten needles or glass micropipettes and applied dorsally at different spinal levels as well as in the torus semicircularis and the ventral thalamus of Rana perezi and P. waltl. Survival times varied from 5 to 10 days. The animals were then reanesthetized and perfused transcardially with isotonic saline followed by a fixative containing 4% paraformaldehyde for the PHA-L and BDA experiments, 1.5% paraformaldehyde and 2% glutaraldehyde for the HRP cases, in PB. The brain and spinal cord were removed, postfixed for four hours, cryoprotected in a 30% sucrose solution in PB, and embedded in gelatin or polyacrylamide (see ten Donkelaar and de Boer van Huizen, 1991). The brains were cut transversally at 40 µm on a freezing microtome. Histochemistry for HRP followed the heavy metal intensification of the diaminobenzidine (DAB)-based HRP reaction product according to Adams (1981). For visualizing BDA, an avidine biotin complex (Vectastain ABC Elite Kit, Vector Laboratories) was used. PHA-L was visualized with an indirect peroxidase anti-peroxidase (PAP) technique (Sternberg, 1979) using as antibodies: 1) goat anti-PHA-L (Vector) (1:2000) overnight at 4°C, 2) donkey anti-goat (Nordic) (1:50) for 1h at room temperature, and 3) PAP goat (Sigma) (1:800) for 1h at room temperature. The antibodies were diluted in 0.1% Triton X-100 in a 0.05M Tris/saline pH 7.6 buffer. In some cases the sections were subsequently osmificated for 30-45 min in 0.1% osmium tetroxide in PB. Selected sections were counterstained with 0.1% cresyl violet solution. The sections were mounted on gelatin coated glasses and coverslipped with Entellan (gelatin embedded sections) or glyceringelatin (polyacrylamide embedded sections).

In vitro tract-tracing experiments. In 10 young adult Xenopus laevis, an in vitro approach was used according to Luksch et al. (1996) based on Cochran et al. (1987). The animals were deeply anesthetized with a 0.2% solution of MS222 and perfused transcardially with iced Ringer solution (75 mM NaCl, 25 mM NaHCO<sub>3</sub>, 2 mM CaCl<sub>2</sub>, 2 mM KCl, 0.5 mM MgCl<sub>2</sub>, 11 mM glucose; pH 7.4). The brains were removed, submerged in the same iced Ringer solution, and cut at middiencephalic or midmesencephalic levels. Applications of 3kD BDA (Molecular Probes, D-7135), recrystallized at the tip of sharp tungsten needles or glass micropipettes, were made with the help of a micromanipulator at the ventral thalamus, the torus semicircularis or at the dorsal horn of the spinal cord. The brains were kept for 18 hours at 15°C in continuously oxygenated Ringer solution (pH 7.4) with carbogen, and subsequently processed as described for the in vivo BDA experiments.

The nomenclature used in this study is largely based on studies by Ebbesson (1976) on the spinal cord, by Opdam and co-workers (Opdam and Nieuwenhuys, 1976; Opdam et al., 1976) and Nikundiwe and Nieuwenhuys (1983) on the brain stem, by Potter (1965) on the midbrain, by Neary and Northcutt (1983) and Wicht and Himstedt (1988) on the diencephalon, and by Northcutt and Kicliter (1980) on the telencephalon.

#### RESULTS

In the present study essentially two types of experiments were carried out. The supraspinal targets of ascending pathways from the spinal cord were studied with anterograde tracers. The cells of origin of spinoreticular, spinotoral and spinothalamic projections were studied with retrograde tracers. Some general remarks are appropriate here. Following iontophoretic or dry application of tracers, rather extensive ascending projections were found. The spinal cord gives rise to distinct, mainly ipsilateral, ascending projections via the dorsal funiculus and the dorsolateral funiculus and predominantly contralateral projections via the ventral funiculus and ventrolateral funiculus (the ventral quadrant of the spinal cord). Due to the proximity of the dorsal horn to the dorsal funiculus and the dorsolateral funiculus, tracer application to the dorsal horn often led to the involvement of the dorsal funiculus and the dorsolateral funiculus resulting in the uptake of tracers by its fibers. In line with previous studies (Antal et al., 1980; Nikundiwe et al., 1982; Jhaveri and Frank, 1983; ten Donkelaar and de Boer-van Huizen, 1991; A.Muñoz et al., 1995a) in such cases spinal dorsal root primary afferents were labeled as well as fibers of the postsynaptic dorsal column system. Moreover, in experiments with tracer applications at cervical and upper thoracic levels, cell groups such as the nucleus of the descending trigeminal tract or their descending fiber projections may have incorporated the tracer from the injection sites. In such cases, trigeminal primary afferents were labeled between the dorsal funiculus and the dorsolateral funiculus in line with previous data

(González and Muñoz, 1987; González et al., 1993). For the sake of clarity, anterograde tracing data will be presented for the dorsal and dorsolateral funiculi, and for the ventral quadrant, separately. In figures 1, 2 and 8 labeled fibers are indicated *only* for the dorsal funicular and the dorsolateral funicular components, whereas in figures 3 and 9 only the labeled fibers passing via the ventral and ventrolateral funiculi are shown. First, the anterograde tracing experiments in the anuran species studied will be discussed, followed by retrograde tracing data on the cells of origin of ascending spinal pathways, and finally the data obtained in *Pleurodeles waltl*.

#### Anterograde tracing experiments in anurans

In a first set of experiments, unilateral applications of the tracers PHA-L, HRP or BDA were made into the dorsal horn of the cervical spinal cord of *Rana perezi* and *Xenopus laevis*. The injections affected the dorsal and lateral spinal fields and, occasionally, the dorsal funiculus and the dorsolateral funiculus. More ventral injections affected the ventromedial, ventrolateral and lateral motor spinal fields, and the ventral and ventrolateral funiculi.

## Ascending spinal projections passing via the dorsal and dorsolateral funiculi

In those experiments that affected the dorsal and lateral grey spinal fields, rostral to the injection site anterogradely labeled fibers were observed in the dorsal funiculus and the dorsolateral funiculus that innervate different supraspinal targets. Two experiments will be described. Following a lumbar BDA application (Fig. 1) labeled fibers could be traced via the ipsilateral dorsal and dorsolateral funiculi.

Most of the labeled fibers ascending in the dorsolateral funiculus turn dorsomedially at upper cervical segments and at the level of the obex, and profusely innervate the neurons in the dorsolateral grey. At these levels, the labeled fibers that course in the dorsal funiculus massively innervate the medial portion of the dorsal column nucleus (DCN) and the caudal aspect of the nucleus of the solitary tract (Fig. 1 F-H). Only a few fibers terminate in the contralateral DCN. Just caudal to the obex a band-shaped area in the grey was found where terminal fibers originating in the dorsal funiculus and dorsolateral funiculus overlap (Fig. 1 F-H). This band borders an unlabeled zone in the dorsolateral margin of the obex region. The latter zone is known to be occupied by descending fibers of the trigeminal tract and the cells related to them (González et al., 1993). The labeled fibers in both funiculi could be traced into the rhombencephalon where they shift to more ventrolateral positions. The fibers from the dorsolateral funiculus terminate diffusely in the lateral reticular zone dorsal to the IXth and Xth motor nuclei (Fig. 1 B-D). Some fibers continue rostrally and innervate the subcerebellar region (Fig. 1A). Labeled fibers ascending via the dorsal funiculus could be traced to the dorsolateral aspect of the rhombencephalon where they innervate the lateral cells of the reticular formation, the area of the nucleus of the descending tract of the trigeminal nerve and the ventral region of the octavolateral area. Only a few fibers reach the level of the trigeminal nerve root and no dorsal funicular fibers were labeled in the cerebellum and the subcerebellar region.

Following tracer applications to the thoracic cord the pattern of labeling in the brain stem is essentially the same Fig. 4A). Two distinct sites of termination of ascending spinal DLF fibers should be emphasized: a zone in the lateral reticular formation between the IXth and VIIth motor nuclei and more, rostrally, the subcerebellar region with a few fibers entering the caudal aspect of the granular cell layer of the cerebellar plate. A small fiber bundle continues rostrally to terminate in the posterodorsal tegmental nucleus of the mesencephalon. The innervation of the mesencephalic tegmentum is more abundant in Rana perezi than in Xenopus laevis. The fibers ascending from the thoracic cord via the dorsal funiculus innervate the nucleus of the solitary tract, the nucleus of the descending trigeminal tract and the octavolateral rhombencephalic area. In X. laevis, these fibers reach the zone of the lateral line nuclei and tracts while in R. perezi, in which no lateral line system is present in the adult, the innervation is restricted to the ventral nucleus of the VIIIth nerve. The rostralmost fibers in both species studied reach the subcerebellar region with some fibers entering the cerebellar granule cell layer.

After a cervical BDA application (Fig. 2) the innervation pattern is similar to that in lumbar and thoracic cases, although the amount of labeling in the lateral reticular formation between the IXth and the Xth motor nuclei (Figs. 2 F-H; 4 C,D) in the subcerebellar region and in the cerebellum (Figs. 2 A, B; 4 B) is much higher. The rostralmost fibers from the dorsolateral funiculus abundantly innervate the caudal aspect of the mesencephalic posterodorsal tegmental nucleus. Some fibers even decussate in the anterior medullary velum to its contralateral part.

Finally, it should be emphasized that, following unilateral application of tracers to lumbar, thoracic or cervical spinal cord, a small contralateral component of ascending fibers in the dorsolateral funiculus was always labeled. This may be due to spread of the tracer to the contralateral side or its uptake by dendrites of contralateral spinal neurons extending into the site of tracer application.

### Ascending spinal projections passing via the ventral quadrant of the spinal cord

In those experiments that affected the dorsal, lateral or ventral spinal fields, a distinct, bilaterally ascending system from the spinal cord was labeled in the ventral quadrant of the white matter. One of these experiments is shown in figure 3, a cervical BDA application. It should be noted that in experiments with tracer applications restricted to the dorsal horn (dorsal and lateral spinal fields) more labeled fibers were found contralaterally than ipsilaterally in the ventral and ventrolateral funiculi. In cases in which tracer applications affected the ventral horn often the ventral and ventrolateral funiculi were damaged resulting in a higher amount of ipsilaterally than contralaterally labeled fibers. This is presumably due to labeling of damaged, crossed fibers. The axons of contralaterally projecting spinal cells could be traced from the injection site ventrally and medially, decussating to the contralateral side beneath the central canal, then turning rostralwards in the ventral and ventrolateral funiculi (Fig. 3V). Axons of ipsilaterally projecting cells were often seen to join the ipsilateral ventral funiculus. As the ventral quadrant component ascends in the rhombencephalon, it smoothly swings to more lateral and dorsolateral positions. Throughout the medulla it gives off thin varicose fibers to different targets (Fig. 3L-S). Most of the labeled fibers innervate structures in the caudal part of the brain stem. A progressive decrease in the amount of labeled fibers was observed as the ventral quadrant component

ascends to more rostral levels. Some of the thin terminal fibers that reach distinct medullary cell masses are collateral branches of thicker fibers located at the margin of the medulla that presumably ascend to more rostral levels. The inferior, middle and superior reticular nuclei receive an extensive innervation, from the ventral quadrant of the spinal cord, while the lateral reticular zone receives only a sparse innervation from this component. Additionally, the IX-Xth motor nuclei, raphe nuclei and the descending trigeminal tract are innervated. More rostrally, labeled fibers were observed in the area between the VIIth and Vth motor nuclei and more sparsely in the central grey at midrhombencephalic levels. A few, smooth fibers course dorsally into the octavolateral area to innervate the dorsal and ventral octaval nuclei (Fig. 3N, O). Some fibers enter the granular layer of the cerebellum where some of them cross the midline in the cerebellar commissure (Fig. 3J, K). A few fibers were observed in subcerebellar areas, just caudal to the isthmic nucleus. More rostrally, fine varicose fibers were observed ventromedial and ventrolateral to the conspicuous isthmic nucleus (Fig. 3 I) where the locus coeruleus and the nucleus of the lateral lemniscus are found. A sparse spinal innervation is also present in the band-shaped area located between the isthmic nucleus and the mesencephalic ventricle. The posterodorsal and posteroventral mesencephalic tegmental nuclei are sparsely innervated (Fig. 3G, H). At caudal mesencephalic levels ventral quadrant fibers turn dorsally along the lateral aspect of the midbrain and bend medially to terminate abundantly in the torus semicircularis (Figs. 3F-H; 4F). The principal, magnocellular and laminar toral nuclei receive spinal projections. A few fibers reach the midline where the commissural nucleus of the torus is located. Occasionally, some labeled fibers and terminals were

observed at the lateral aspect of the mesencephalic tectum (Fig. 4E). In both species, at more rostral mesencephalic levels, the anterodorsal and anteroventral tegmental nuclei, the red nucleus and the interstitial nucleus of the fasciculus longitudinalis medialis are slightly innervated by spinal ventral quadrant fibers (Fig. 3E). In addition, scattered fibers distribute to the pretoral and pretectal grey and some fibers cross in the posterior commissure. Beyond the midbrain, both dorsal and ventral thalamic areas are innervated by ventral quadrant fibers (Fig. 3A-D). A few thin, varicose fibers innervate the posterior and central dorsal thalamic nuclei whereas the anterior nucleus receives only a sparse spinal innervation. In addition, the posterodorsal and posteroventral lateral nuclei are also sparsely innervated. The ventromedial thalamic nucleus and the dorsal aspect of the posterior tubercle are far more densely innervated. The fibers reaching all cell layers of the ventromedial nucleus pass through the dorsal part and, especially, the ventral part of the ventrolateral thalamic nucleus where varicosities were also found among its cells. No labeling was found more rostrally in the diencephalon or in the telencephalon in any of the cases studied.

After thoracic or lumbar HRP or BDA applications, a largely similar pattern of anterograde labeling, although less conspicuous, was observed. In all cases innervation of the aforementioned rhombencephalic and midbrain tegmental areas was observed. A low density of labeled fibers was present in the torus semicircularis and in the ventral thalamus. The latter structure was only very sparsely innervated in experiments with lumbar spinal injections.

Retrograde tracer experiments in anurans

BDA was used to trace the cells of origin of components of ascending spinal pathways. The tracer was applied to the ventral part of the thalamus, to the torus semicircularis and to the reticular formation. In a first set of experiments in Rana perezi, BDA was applied to the ventral part of the thalamus or to the torus semicircularis, two main targets of ascending sensory pathways from the spinal cord. After BDA application to the ventral thalamus, retrogradely labeled cells were observed, predominantly contralaterally, in the sensory trigeminal nuclei, in the dorsal column nucleus, in the lateral cervical nucleus and in the cervical, and to a lesser extent, thoracic spinal cord. Round, triangular and irregularly- shaped neurons were observed mainly in the dorsal spinal field and a few cells were present in the lateral field. Some piramidal and bipolar cells with dendrites extending predominantly horizontally were observed in the ventral fields of cervical segments. Higher numbers of retrogradely labeled cells were seen in the spinal cord in experiments with BDA applications to the torus semicircularis.

In a second set of experiments in Xenopus laevis, an in vitro approach was used. In an isolated brain preparation of young adult X. laevis, 3kD BDA was applied to the ventral part of the thalamus, to the torus semicircularis and to the reticular formation (Fig. 5). In such isolated brain preparations a more extensive pattern of labeling was observed. In all experiments labeled cells were observed more contralaterally than ipsilaterally. In experiments with ventral thalamic BDA applications (Fig. 5A) numerous labeled cells were present in the cervical spinal cord (Fig. 6A), whereas only a few cells were found at thoracic levels and no lumbar neurons were labeled at all. Most of the cells (80%) were found contralateral to the application site, and about 20% ipsilaterally projecting spinothalamic cells were observed. Cells with round, bipolar and irregular morphology were found.

After BDA applications to the torus semicircularis more retrogradely labeled cells were seen in the spinal cord (Fig. 5B). Moreover, labeled cells were observed throughout the spinal cord. Again, more contralaterally (65%) than ipsilaterally (35%) labeled neurons were found. This proportion remained constant throughout the rostrocaudal extent of the spinal cord. Most spinotoral cells (about 80%) were found in the cervical cord (Fig. 6C, D, F, G), about 10% at the thoracic level and about 10% in the lumbar cord. At cervical levels, labeled neurons were found predominantly in the ventral part of the dorsal horn. Most neurons have round or bipolar cell bodies with dendritic trees extending laterally towards the dorsolateral funiculus and within the dorsal field. Additionally, triangular and irregularly-shaped large cells with dendrites extending dorsally, ventrally and medially were present. The axons of these cells course ventromedially, cross the midline ventral to the central canal and join the contralateral ventral funiculus where they turn rostrally. Triangular or bipolar, horizontally oriented, ipsilaterally projecting cells send their axons to the ipsilateral ventral funiculus. Some neurons were also present bilaterally in the lateral field. Round and triangular as well as larger multipolar neurons were labeled bilaterally in the ventral spinal grey within the ventromedial, ventrolateral and lateral motor fields. At thoracic levels, neurons were also found particularly in the ventral part of the dorsal field. Small, round and large irregular cells were observed. The dendrites of the large, irregular neurons are directed dorsally and medially within the dorsal field and laterally towards

the dorsolateral funiculus. A few labeled neurons were present in the ventromedial and ventrolateral spinal fields. In some cases, the axons of ipsilaterally labeled cells follow an unusual course and cross the midline twice at the spinal level where the cell body is located, first dorsal and then ventral to the central canal to ascend in the ipsilateral ventral funiculus. In the lumbar cord, the dorsally located cell bodies, most of them contralateral to the application site, are bipolar and oriented horizontally, with medially and laterally directed dendritic processes.

Experiments with BDA applications to the rhombencephalic reticular formation were analyzed only in those cases in which the spread of the tracer into the adjacent white matter was minimal. After BDA applications to the reticular formation at the level of the inferior reticular nucleus (Fig. 5C), retrogradely labeled neurons were observed thoughout the spinal cord up to upper lumbar levels. Most cells were present in the cervical cord (Fig. 6B, E). In striking contrast to the thalamic and toral tracer application experiments, a more equal percentage of ipsilaterally (45%) and contralaterally (55%) projecting neurons was found. Their axons ascend at the ventral aspect of the lateral funiculus dorsal to those corresponding to spinotoral and spinothalamic projecting neurons. Medium-sized and large, fusiform and irregularly shaped neurons were found bilaterally in the deep dorsal field and, to a lesser extent in more superficial parts of the dorsal field. Their dendrites extend thoughout the dorsal horn and their axons are directed ventromedially or ventrolaterally to join the contralateral or ipsilateral ventrolateral funiculus, respectively. Additionally, a higher amount of ipsilateral than contralateral, round, irregular and fusiform medium-sized neurons were found in the lateral spinal field, with axonal processes directed to the ventrolateral funiculus. A few small, mainly round neurons were found, bilaterally, in the ventral horn. They were located in the ventromedial and ventrolateral fields.

## Anterograde tracing experiments in an urodele

In the ribbed newt, *Pleurodeles waltl*, comparable experiments were carried out. Since hardly any data are available on dorsal root projections for urodeles, one experiment is included.

## Ascending spinal projections passing via the dorsal and dorsolateral funiculi

In figure 7 an experiment is shown in wich a main brachial dorsal root was cut and subsequently labeled with BDA. In such experiments two fiber tracts were labeled in the spinal cord: a medial bundle in the dorsal funiculus consisting of thick fibers, and a lateral group of thin fibers within the dorsal portion of the dorsolateral funiculus, i.e. Lissauer's tract. Both fiber systems project to widespread spinal and supraspinal regions. Here only the distribution of dorsal root afferents to supraspinal targets will be discussed. Within the dorsal funiculus, spinal primary afferents ascend, somatotopically arranged, to the obex region. Fibers originating at cervical segments are present in the lateral part of the dorsal funiculus. In experiments in which a lumbar dorsal root was labeled, fibers were found in a position medial to those of cervical origin. These ascending spinal projections via the dorsal funiculus outline the DCN at the obex level (Fig. 7M, O). The terminal fields in this area largely resemble the organization of the fibers in the

dorsal funiculus. Thus, with a certain degree of overlap in the projection, medially situated axons from lumbar dorsal root ganglion cells terminate on medial cells in the DCN, whereas laterally located fibers arising from cervical dorsal root ganglion cells end on more lateral cells. Most of the primary afferents terminate dorsal to the cells of the DCN. However, some fibers reach more ventrolaterally located positions. Rostral to the obex level, tightly packed dorsal funicular fibers gradually turn ventrolaterally and ascend throughout the medulla dorsal to the descending trigeminal tract (Fig. 7J-H). Lumbar primary afferents could not be traced far beyond the rostral limit of the DCN. However, brachial primary afferents extend as far rostrally as the cerebellum where they arborize profusely within the granular layer (Fig. 7A, P). Throughout the rhombencephalon, varicose fibers leave the tract and arborize within the white matter where dendrites of the adjacent periventricular cells of the reticular formation, the nucleus of the solitary tract, the nucleus of the descending trigeminal tract and of the octavolateral area can be contacted. More rostrally, at the level of the VIIth and VIIIth nerve roots, some fibers enter the lateral reticular zone and the ventral aspect of the octavolateralis area.

After BDA applications to the cervical spinal cord in *Pleurodeles*, labeled fibers could be traced to supraspinal targets via the dorsal, dorsolateral, ventral and ventrolateral funiculi organized in a way similar to that in anurans, although some differences were eminent. One experiment is shown in figure 8. BDA was applied to the midcervical segment of the spinal cord (Fig. 8P). The application site included the dorsal and dorsolateral funiculi. In the dorsal funiculus, two different, adjacent components of labeled fibers could be distinguished (Fig. 8M-O): a medial component composed of thick, tightly packed fibers, and a second more dorsolaterally located component that is flanked laterally by the descending trigeminal tract. At spinal levels both components are very close to each other. At the obex level the medial component occupies a dorsomedial position and its thick fibers do not give off varicose terminal fibers. In turn the more sparse lateral fibers give off abundant thin, varicose fibers in this region that enter the dorsalmost aspect of the grey. Slightly rostral to the obex level, a small wedged-shaped, non-labeled area separates both components of labeled fibers (Fig. 8L, M). This is more evident rostrally in the rhombencephalon as the dorsal funicular system, the descending trigeminal tract and the dorsolateral funicular system swing ventrolaterally (Fig. 9A, B). A progressive decrease in the number of fibers of the medial component of the dorsal funicular fibers was observed up to the level of the facial motor nucleus where they fade and terminate as thick, varicose fibers in the lobe of the lateral line. However, the lateral component of the dorsal funicular fibers gives off thin terminal branches that massively reach the dorsal grey at caudal medullary levels. More rostrally, many fibers arborize within the white matter adjacent to the tract but some fibers reach the adjacent reticular, octavolateral and trigeminal areas as they ascend in the medulla up to the cerebellum. Here they distribute profusely to the ipsilateral superficial half of the granular cell layer (Fig. 8A).

Anterogradely labeled thin fibers in the dorsolateral funiculus ascend throughout the medulla just ventral to the descending trigeminal tract (Figs. 8, 9A, B). Along their course varicose terminal fibers were observed within the tract but they rarely enter the adjacent grey in the lateral reticular formaion, and the descending trigeminal nucleus. Numerous varicose fibers and terminals were observed in the ventrolateral alar grey at the obex level and between the rostralmost root of the IXth-Xth complex and the root of the trigeminal nerve. Here some fibers turn toward the ventral aspect of the octavolateral area. More rostrally, at the level of the cerebellum, the dorsolateral funicular fibers shift dorsally and medially and profusely innervate the subcerebellar region. A few fibers enter the cerebellum. The rostralmost fibers sparsely innervate the caudal aspect of the mesencephalic tegmentum.

## Ascending spinal projections passing via the ventral quadrant of the spinal cord

In experiments with BDA applications to the dorsal grey at cervical spinal segments a predominantly contralateral bundle of labeled fibers could be traced via the ventral and ventrolateral funiculi to the brain stem and diencephalon (Fig. 9). Most fibers cross the midline below the central canal and enter the contralateral ventral funiculus. Here, the fibers bend rostrally and ascend to various rhombencephalic, mesencephalic and diencephalic areas. Like in anurans, ascending spinal fibers passing via the ventral quadrant swing to a more lateral position in the rhombencephalon (Fig. 9F-L). In the rhombencephalon, projections were found throughout the rostrocaudal extent of the reticular formation (Figs. 9G-L; 10C), to the area octavolateralis and to the cerebellum (Fig. 9F). At caudal mesencephalic levels the ascending fibers bend dorsolaterally and innervate the outer layers of the tectum mesencephali (Figs. 9D, E; 10D), the torus semicircularis (Figs. 9D; 10E), and the midbrain tegmentum. Within these areas, the labeled fibers are mainly distributed in the inner region of the external white fiber layer, and hardly reach the periventricular cell layer. In the diencephalon, a main projection field is present in the ventral thalamus, and forms a rostral continuation of that observed in the torus semicircularis and dorsal midbrain tegmentum. Generally, in the thalamus, the fibers distribute in the outer fiber layer but do not penetrate the periventricular cell layer. A few fibers pass dorsal to the intermediate sulcus and innervate both the posterodorsal and anteroventral dorsal thalamic areas.

#### Retrograde tracer experiments

In experiments in which BDA crystals were applied to the ventral thalamic region or to the torus semicircularis of *Pleurodeles waltl*, retrogradely labeled neurons were found in the spinal cord. The majority of spinothalamic or spinotoral cells was observed in the dorsal grey of the contralateral cervical spinal cord (e.g., Fig. 10F). However, a small number of labeled cells was also present in the lateral and ventral fields of the grey. A small component of ipsilateral cells was also observed. In this set of experiments, no labeled cells were found more caudally in the spinal cord.

#### DISCUSSION

In the present study the mainly anterograde tracer PHA-L and the bidirectionally transported tracers HRP and 10kD BDA were used in an *in vivo* approach in adult *Rana perezi*, *Xenopus laevis* and *Pleurodeles waltl*. Additionally, in an *in vitro* approach 3kD BDA was used in isolated brain preparations of young adult *Xenopus laevis*. The results with both the *in vivo* and *in vitro* tracing techniques were largely comparable. However, the 3kD BDA *in vitro* approach presents several advantages such as fast transport, the good neuronal labeling including secondary and tertiary dendrites, good labeling at long distances, the easy accessibility of all brain areas without the surgical and survival problems of the animals in cases with large lesion or massive tracer application and the precise tracer application under visual control (Luksch et al., 1996).

With these tracer techniques three main components of ascending spinal projections were demonstrated, i.e. a dorsal funicular component, a dorsolateral funicular component and a ventral quadrant component. In previous studies (A. Muñoz et al., 1994b, 1995a, b, 1996), spinal projections to the dorsal column and lateral cervical nuclei, respectively, were demonstrated. Additionally, the diencephalic targets of the ventral quadrant of the spinal cord were demonstrated (A. Muñoz et al., 1994a). In the present study the organization, the main targets and the cells of origin of some of the ascending spinal projections were investigated.

In previous studies, based on silver-stained material or making use of anterograde degeneration techniques, ascending spinal projections were demonstrated in anurans (Ebbesson, 1969; 1976; Hayle, 1973a,b) and in urodeles (Herrick, 1914, 1930; Herrick and Bishop, 1958; Nieuwenhuys and Cornelisz, 1971). In *Necturus*, Herrick (1930) considered the ascending spinal projections through the dorsolateral funiculus as a distinct ascending spinal system, independent of the spinal lemniscus, and named it "the spinobulbar tract". At rhombencephalic levels it is located just ventral to the descending trigeminal tract and dorsal to the reticular neuropil. Additionally, Herrick (1930) observed a conspicuous terminal field of this spinobulbar tract just caudal to the obex and, rostrally, a close relationship of this ascending system with the reticular formation up to the level of the trigeminal nerve root. However, in later studies the ascending spinal projections via the dorsolateral funiculus were considered part of the spinal lemniscus (Herrick, 1948; Herrick and Bishop, 1958; Hayle, 1973a). In the axolotl, Nieuwenhuys and Cornelisz (1971) reported a distinct ascending spinal system in the dorsal part of the lateral funiculus that innervates the medulla and the mesencephalic tectum. It was interpreted part of Herrick's spinal lemniscus.

Additionally, Herrick (1930) described the spinal lemniscus as a system that ascends through the ventral quadrant of the spinal cord, separate from the spinobulbar tract at rhombencephalic levels by the reticular neuropil area. Subsequently, Ebbesson (1976) subdivided the ascending spinal projections in Rana catesbeiana that pass via the "anterolateral funiculus" into medial and lateral components. His lateral component is comparable to the dorsolateral funicular component, whereas his medial component forms part of our ventral quadrant component. However, Ebbesson's medial component predominantly innervates the inferior and middle reticular fields and fades at rostral rhombencephalic and caudal mesencephalic levels and his lateral component apperently innervates more rostral mesencephalic levels and even the thalamus, at least from the upper cervical cord. In the present study, based on more sensitive tract-tracing techniques, the dorsolateral funicular component was found to fade at cerebellar and caudal mesencephalic levels, whereas the fibers of the ventral quadrant component, located more ventrally throughout the rhombencephalon, turn dorsally at the isthmic level and reach mesencephalic and diencephalic targets (A.Muñoz et al., 1994a).

The spinal ascending pathways demonstrated in the present study will now be discussed in an evolutionary perspective.

#### Dorsal funicular pathways in amphibians

Tracer applications to the dorsal spinal cord at cervical, thoracic and lumbar levels showed that the fibers ascending via the dorsal funiculus and their pattern of termination in the DCN region are organized somatotopically, as previously noted in anurans (Antal et al., 1980; Nikundiwe et al., 1982; M.Muñoz et al., 1991; A.Muñoz et al., 1995b). The dorsal funicular fibers are ascending collaterals of primary afferents from spinal dorsal roots and most likely include second order projections towards the DCN, i.e. the postsynaptic dorsal column system (ten Donkelaar and de Boer van Huizen, 1991; A.Muñoz et al., 1995b). Part of the dorsal funicular component innervates rhombencephalic structures rostral to the DCN, in line with data obtained after labeling of spinal primary afferents (Joseph and Whitlock, 1968a; Antal et al., 1980; Nikundiwe et al., 1982; M.Muñoz et al., 1991; A.Muñoz et al., 1995b). The postsynaptic dorsal column system may contribute to this innervation. The most rostral site of termination of dorsal funicular fibers is the cerebellum. No dorsal funicular fibers from lumbar segments reach the cerebellum in line with previous data (Nikundiwe et al., 1982).

In urodeles, the glossopharyngeal and vagal nerves are known to send descending branches into the dorsal funiculus (Roth and Wake, 1985). Our BDA experiments on the ascending pathways of the cervical spinal cord of *Pleurodeles* (Fig. 8) showed two different components of labeled fibers in the dorsal funiculus. The lateral component is comparable to the dorsal funiculus of anurans and includes primary and, most likely, also non-primary ascending fibers. This component terminates mainly in the dorsal column nucleus at the obex level, but could be traced through the medulla up to the cerebellum, in line with our data on spinal primary afferents from the second spinal dorsal root (Fig. 7). Roth and Wake (1985) did not mention second dorsal root primary afferents reaching the cerebellum in lungless salamanders. The medial dorsal funicular component ascends, tightly packed up, to the area of the lateral line, where it fades, and corresponds to afferent fibers of the second root of the glossopharyngeal nerve as described by Roth and Wake (1985). Descending branches of this cranial nerve later mingle with spinal dorsal root projections in the dorsal funiculus up to the level of the fourth spinal nerve. Between the two dorsal funicular components an unlabeled region was found in *Pleurodeles* that may correspond to the area in which Fritzsch (1988) observed descending branches of inner ear afferents in various urodele species.

The dorsal funiculus is mainly ained at the dorsal column nucleus at the spinomedullary border. This nucleus gives rise to a contralaterally ascending projection, i.e. the medial lemniscus, to the midbrain and thalamus (A. Muñoz et al., 1994b, 1995b).

#### Dorsal funicular pathways in vertebrates

Throughout vertebrates the dorsal funiculus contains ascending collaterals of primary spinal afferents reaching obex levels and continuing into the ipsilateral alar medulla (Table 1). In lampreys, dorsal funicular fibers innervate neurons at the obex level and in the octavolateral area (Northcutt and Ebbesson, 1980; Ronan and Northcutt, 1990). Dorsal column

fibers innervate the cerebellum as well (Ronan and Northcutt, 1990; Dubuc et al., 1993). In the Pacific hagfish, Eptatretus stouti, it is unclear whether dorsal column fibers reach the, little evident, cerebellum (Ronan and Northcutt, 1990). Although in the spotted dogfish, Scyliorhinus canicula, Hayle (1973a) was unable to trace degenerating fibers in the dorsal funiculus more than several segments rostral to spinal lesions, Ebbesson and Hodde (1981) demonstrated dorsal column fibers to a dorsal column nucleus at the obex level, and to the vestibular nuclear complex and cerebellar granule layer, in the nurse shark, Ginglymostoma cirratum. Experimental evidence for dorsal column pathways in bony fishes is limited. In the himé salmon, Oncorhynchus nerka, Oka et al. (1986) applied cobaltic lysine to the cut end of the spinal cord at the 10th to 15th spinal segment, and traced labeled axons via the dorsal funiculus to a site of termination at the obex level but hardly beyond the caudal medulla. Ronan and Northcutt (1990) reported the presence of dorsal funicular fibers to the obex and octavolateral area in African lungfishes (Protopterus species), bichirs (Polypterus species), and in the gar, Lepisosteus osseus. In bichirs and in gars, no dorsal funicular fibers were found to the cerebellum, but in lungfishes dorsal column fibers terminate in the granule layer of the cerebellum.

In amphibians, dorsal column fibers project to the ill-defined dorsal column nucleus, to the octavolateral area and to the cerebellum (Joseph and Whitlock, 1968a; Ebbesson, 1969, 1976; Hayle, 1973a.b; Antal et al., 1980; Székely et al., 1980; Nikundiwe et al., 1982; Urban and Székely, 1982; A. Muñoz et al., 1994b, 1995b, the present study; see Table 1). The dorsal column system has been extensively studied in reptiles. In the tegu lizard, Tupinambis teguixin (Ebbesson, 1967), a lacertid, Lacerta viridis (Jacobs, 1968), a boid snake, Constrictor constrictor (Ebbesson, 1969), a crocodilian, Caiman crocodilus (Ebbesson and Goodman, 1981), and in various turtle species, especially Pseudemys scripta elegans (Ebbesson, 1969; Pedersen, 1973; Künzle 1982; Künzle and Woodson, 1983) the dorsal funiculus innervates the dorsal column nucleus, vestibular nuclei, and the cerebellum. In Thamnophis sirtalis, Jacobs and Sis (1980) found a dorsal column projection to the DCN and to the descending vestibular nuclei. Comparable data were obtained in Iguana iguana (Joseph and Whitlock, 1968b) and in Python reticulatus (Kusuma and ten Donkelaar, 1980). In birds, Karten (1963) and van den Akker (1970) noted a spinal projection to the dorsal column nuclei of pigeons, Columba livia. In an HRP tracing study in pigeons, Wild (1985) found extensive dorsal root projections to the dorsal column nuclei, but also to a wide dorsolateral region of the rhombencephalon which includes the external cuncate nucleus, the lateral part of the nucleus of the descending trigeminal tract, and the principal trigeminal sensory nucleus. A discrete projection was also found to the nucleus of the solitary tract.

Ascending spinal fibers of both primary and non-primary origin are found in the dorsal funiculus of mammals (see Willis and Coggeshall, 1991 for review). The presence of non-primary spinal afferents to the dorsal column nucleus, i.e. the postsynaptic dorsal column system, has now been demonstrated throughout terrestrial vertebrates (e.g., Rustioni, 1973; Angaut-Petit, 1975a,b; Rustioni and Kaufman, 1977; Bennett et al., 1984; Giesler et al., 1984; Funke, 1988; ten Donkelaar and de Boer-van Huizen, 1991; Pritz and Stritzel, 1994; A. Muñoz et al., 1995b).

## Dorsolateral funicular pathways in amphibians

In the present study a well-developed system of ascending fibers in the amphibian dorsolateral funiculus was demonstrated. The intraspinal targets of these fibers include the dorsal and lateral spinal field and, especially, the lateral cervical nucleus (A. Muñoz et al., 1995a, 1996). In the rhombencephalon, several structures are innervated by dorsolateral funicular fibers including the nucleus of the solitary tract, a lateral reticular zone and a subcerebellar "parabrachial" region.

In a previous in vitro tract-tracing study in Xenopus laevis (A. Muñoz et al., 1996), ascending spinal fibers from all levels of the spinal cord, passing via the dorsolateral finuculus, were found to terminate in a cell area ventrolateral to the dorsal column nucleus. This cell area can be considered a possible homologue of the mammalian lateral cervical nucleus. Moreover, this cell area was found to project contralaterally to the torus semicircularis and to the ventral thalamus, both targets for somatosensory projections. Similar observations were made in Rana perezi (A. Muñoz et al., 1995a). Data in R. catesbeiana tadpoles (Forehand and Farel, 1982) also indicate the presence of a lateral cervical nucleus in this ranid frog. The amphibian lateral cervical nucleus is innervated by the spinocervical tract which arises, mainly ipsilaterally, in the ventral part of the dorsal horn throughout the spinal cord (A. Muñoz et al., 1995a, 1996). In the present study three main other, rhombencephalic targets were found. In Rana perezi as well as in X. laevis, dorsolateral funicular fibers

innervate the area of the nucleus of the solitary tract. Moreover, dendrites of the tyrosine hydroxylasepositive neurons, present in this nucleus, extend laterally and reach the dorsolateral funiculus (González and Smeets, 1994) where they may receive input from Lissauer's tract and from the dorsolateral funiculus. Spinal projections to the nucleus of the solitary tract were also observed in an anterograde degeneration study in R. catesbeiana. Hayle (1973a), however, did not report a spinosolitary projection in R. temporaria. In Pleurodeles waltl, the very poor segregation of neurons in the caudal part of the alar plate makes it difficult to distinguish which component may receive dorsolateral funicular afferents. In early studies in Ambystoma tigrinum, Herrick (1930) suggested that a spinobulbar tract, passing via the dorsolateral funiculus, contributes to a "general coordination field" at the obex level together with descending bulbospinal systems. The lateral reticular zone in the rhombencephalon is also innervated by dorsolateral funicular fibers. Especially, a lateral reticular area located dorsal to the VIIth-IX motor nuclei and ventral to the ventral nucleus of the VIIIth nerve, was labeled. In this area, retrogradely labeled cells were found after thoracic and cervical HRP injections (ten Donkelaar et al., 1981).

A distinct ipsilateral projection from the dorsolateral funiculus was found to a subcerebellar region. This projection reaches the caudal part of the secondary visceral nucleus as distinguished by Larsell (1923), Barnard (1936) and Opdam et al. (1976). Both Larsell (1923) and Barnard (1936) suggested a tractus visceralis secundarius connecting the nucleus of the solitary tract with the nucleus visceralis secundarius. This distinct nucleus lies ventrolateral to the nucleus cerebelli (Opdam et al., 1976). HRP injections including the subcerebellar region resulted in labeling of cells in the nucleus of the solitary tract (González et al., 1984; A. Muñoz et al., 95b; in preparation). Tract-tracing studies in ranid frogs (Neary and Wilczynski, 1986; Neary, 1988, 1995) and in the green treefrog, *Hyla arborea* (Allison and Wilczynski, 1991), showed a distinct projection of a secondary isthmal nucleus in a position comparable to that of the nucleus visceralis secundarius to the ventral hypothalamus and to the preoptic area. Spinal projections were also found to arise in this "parabrachial" area (ten Donkelaar et al., 1981). In line with Neary's (1995) study, this subcerebellar region will be described as the *parabrachial nucleus*.

# Dorsolateral funicular pathways in vertebrates

A spinocervical system, passing via the dorsolateral funiculus and terminating in a lateral cervical nucleus, appears much more common than previously thought (A. Muñoz et al., 1996). In anamniotes other than amphibians, evidence for a spinocervical tract projecting to a lateral cervical nucleus is at least suggestive. In petromyzontid and myxinoid agnathans (Northcutt and Ebbesson, 1980; Ronan and Northcutt, 1990) as well as in cartilaginous (Hayle, 1973a,b; Ebbesson and Hodde, 1981) and bony fishes (Hayle, 1973a,b; Finger, 1981; Ito et al., 1986) ascending spinal projections via the dorsal part of the lateral funiculus were demonstrated (Table 2). No separate site of termination, reminiscent of a lateral cervical nucleus, was noted. In reptiles, experimental evidence for the presence of a spinocervical tract comes from an anterograde degeneration study in the tegu lizard, Tupinambis teguixin. Ebbesson (1967) noted that at caudal brainstem levels some collateral fibers

leave the dorsolateral funiculus and innervate an area located dorsal to the hypoglossal nucleus and ventral to the DCN. A tracing study in Pseudemys scripta elegans (Künzle and Woodson, 1982) also suggests the presence of a spinocervical tract. In birds, van den Akker (1970) showed a "dorsolateral ascending bundle" in the dorsolateral funiculus that arises in neurons found in the deep part of the dorsal horn. At cervical levels, this bundle innervates the deep dorsal and central spinal grey. More recent tract-tracing studies showed various ascending non-primary spinal projections in the dorsolateral funiculus of the pigeon (Funke and Necker, 1986; Funke, 1988; Necker, 1991), most likely including the spinocervical tract. The spinocervical tract in mammals has been studied extensively (see Willis and Coggeshall, 1991). It forms the first part of a bisynaptic spinocervicothalamic pathway (Morin, 1955; Nijensohn and Kerr, 1975; Boivie, 1983). Tracttracing data in Xenopus laevis and in Rana perezi (A. Muñoz et al., 1995a, 1996) suggest the presence of an anuran homologue of the mammalian spinocervicothalamic system. Such a system is much more common in vertebrates than previously thought.

Spinosolitary projections passing via the dorsolateral funiculus were observed in reptiles (Ebbesson, 1967, 1969; Pedersen, 1973), in birds (Karten, 1963; Funke and Necker, 1986; Funke, 1988), and in mammals (e.g., Kuru, 1956; Hazlett et al., 1972; McMahon and Wall, 1983; Apkarian et al., 1985; Menétrey and Basbaum, 1987). Although in mammals the bulk of spinal projections to the reticular formation ascends via the anterolateral system (see Mehler, 1969; Willis and Coggeshall, 1991), ascending projections to different parts of the lateral reticular zone via the dorsolateral funiculus were demonstrated (e.g., Zemlan et al., 1978; Blomqvist and Berkley, 1992). In cats, Blomqvist and Berkley (1992) showed spinoreticular fibers to the area surrounding the facial motor nucleus. They suggested that this termination field includes the feline adrenergic  $C_1$  cell group in the rostral ventrolateral medulla caudal to the VIIth motor nucleus, and the catecholaminergic A5 group rostral to this motor nucleus. Neurons in the rostral ventrolateral medulla including the C1 cell group project to the spinal cord via the dorsolateral funiculus (e.g., Ross et al., 1984; Dampney et al., 1987; Tucker et al., 1987). Although in the amphibian rhombencephalon the presence of adrenergic and noradrenergic cell structures in the rhombencephalon is not entirely clear (González and Smeets, 1994), comparison with the data available for mammals suggest than in amphibians the lateral reticular zone at the level of the VIIth motor nucleus is involved in viscerosensory functions.

The main rostral site of termination of ascending dorsolateral funicular fibers is the amphibian homologue of the mammalian parabrachial nucleus. In mammals, the parabrachial complex includes various subnuclei and the nucleus of Kölliker-Fuse (Saper, 1995). The parabrachial nucleus receives the majority of the ascending projections from the nucleus of the solitary tract (e.g., Herbert et al., 1990) as well as extensive projections from the spinal cord and different trigeminal nuclei (Bernard et al., 1995; Feil and Herbert, 1995). Several studies showed ascending spinal projections to the parabrachial nucleus via the dorsolateral funiculus (Nijensohn and Kerr, 1975; Björkeland and Boivie, 1984; Hylden et al., 1986; Kitamura et al., 1993; Slugg and Light, 1994; Bernard et al., 1995; Feil and Herbert, 1995). This bilateral projection arises mainly in laminae I-II of the dorsal horn (see Bernard et al., 1995; Feil and Herbert, 1995). This projection most likely plays a crucial role in a variety of important physiological systems, ranging from pain and automatic control through arousal and ingestive behaviour (Saper, 1995).

Some dorsolateral funicular fibers in amphibians continue rostrally and reach the caudal aspect of the posterodorsal nucleus of the mesencephalic tegmentum. This sparse projection may be reminisent of the spinomesencephalic projection via the dorsolateral funiculus in mammals to the periaqueductal grey and the nucleus cuneiformis (e.g., Nijensohn and Kerr, 1975; Zemlan et al., 1984; Björkeland and Boivie, 1984; Hylden et al., 1986; Yezierski, 1988; Bernard et al., 1995).

#### Ventral quadrant pathways in amphibians

In the anurans studied, in line with previous studies (Ebbesson, 1969, 1976; Hayle, 1973a), the bulk of fibers ascending via the ventral quadrant of the spinal cord fades at rhombencephalic levels. Ventral quadrant pathways terminate mainly in the reticular formation, particularly in its caudal, inferior part. Additionally, in the present study ventral quadrant projections were found to be lateral reticular zone, the raphe nucleus, the nucleus of the descending tract of the trigeminal nerve, the octavolateral area and the inferior olive. More rostrally, the parabrachial nucleus receives as sparse ventral quadrant projection, in addition to the massive innervation by the dorsolateral funiculus.

In the rostral part of the rhombencephalon, some fibers turn dorsally and enter the cerebellar granule layer. These fibers presumably arise from neurons located mainly in the ventral horn of the spinal cord (González et al., 1984; Grover and Grüsser-Cornehls, 1984). This ventral spinocerebellar pathway forms one of the three spinocerebellar projections found in amphibians (Joseph and Whitlock, 1968a; Nieuwenhuys and Cornelisz, 1971; Hayle, 1973b; Ebbesson, 1976; Antal et al., 1980; González et al., 1984; Grover and Grüsser-Cornehls, 1984; the present study). The other two spinocerebellar projections in amphibians are the ipsilateral, primary spinocerebellar projection via the dorsal funiculus, and a non-primary dorsal spinocerebellar pathway via the dorsolateral funiculus that ascends through the medulla just ventral to the descending tract of the trigeminal nerve.

The ventral quadrant projection to the isthmic region found in the present study is in line with earlier anterograde degeneration studies (Ebbesson, 1969, 1976; Hayle, 1973a). In the mesencephalon, the torus semicircularis is the main target of ventral quadrant fibers. The torus semicircularis is a major integrating center for a number of sensory and nonsensory afferents in addition to auditory inputs, and may serve a role similar to the one the tectum mesencephali serves for the visual system (Wilczynski and Capranica, 1984). A distinct, mainly contralateral spinotoral projection arises from all levels of the anuran spinal cord. In Rana catesbeiana, Ebbesson (1976) noted a sparse spinomesencephalic projection to the magnocellular and laminar nuclei of the torus semicircularis. In the present study, a more extensive spinotoral innervation was found: spinal fibers mainly innervate the magnocellular and laminar nuclei but, although more sparsely, also the principal and commissural nuclei. More rostrally, labeled fibers were also found in the pretoral grey. Apart from this direct spinotoral input, the torus semicircularis

receives somatosensory input from the nucleus of the descending tract of the trigeminal nerve (M. Muñoz et al., 1994), from the dorsal column nucleus (Wilczynski, 1981; Wilczynski and Neary, 1986; A. Muñoz et al., 1994b, 1995b) and from the lateral cervical nucleus (A. Muñoz et al., 1995, 1996). In *R. pipiens*, Comer and Grobstein (1981a,b) demonstrated the involvement of the torus in tactually elicited prey acquisition behavior. A crude somatotopic map of the contralateral body surface was found for the torus (Comer and Grobstein, 1981c). A sparse spinal innervation of the tectum mesencephalic was observed.

In the present study a direct, rather extensive, mainly contralateral spinothalamic projection was demonstrated. This projection arises mainly from cervical, but also from thoracic and, more sparsely, from lumbar spinal segments. Most cells of origin were observed in the contralateral dorsal horn. Spinothalamic fibers innervate the posterior tubercle, the ventromedial and the ventrolateral thalamic nuclei. Only a few spinal fibers reach the posterior, central and anterior thalamic nuclei.

In Pleurodeles waltl, the ventral quadrant pathways are comparable to those in anurans. A notable exception in the well-developed spinotectal component. The pattern of termination of spinoreticular projection is largely comparable to that found with the classical silver impregnation techniques (Herrick, 1948; Herrick and Bishop, 1958) and with anterograde degeneration techniques (Nieuwenhuys and Cornelisz, 1971; Ebbesson et al., 1972). The indistinct torus semicircularis, characterized by its octavolateral input (González and Muñoz, 1987), receives spinal input via the ventral quadrant pathways. In Ambystoma tigrinum, Herrick (1914,

1948) suggested a direct spinotectal projection. In the axolotl, A. mexicanum, Nieuwenhuys and Cornelisz (1971) experimentally demonstrated this projection. Our BDA tracing experiments in P. waltl show a mainly contralateral spinotectal projection. Some contradictory data were obtained with regard to the origin of this projection. After unilateral HRP injections in Salamandra salamandra, Finkenstädt et al. (1983) found spinal projections from the dorsal grey matter of the ipsilateral spinal cord. Rettig (1984, 1988) in S. salamandra and Bolitoglossa subpalmata as well as Gruberg and Harris (1981) in A. tigrinum and A. mexicanum, however, found bilateral spinotectal projections from neurons in the ventral horn. With electrophysiological techniques, Gruberg and Harris (1981) observed a rostrocaudal and lateromedial somatosensory body reopresentation in layer 3 of the tectum, below the visual input.

In the present study a sparse spinal innervation of the anteroventral and posterodorsal thalamic zones was observed. A much denser spinothalamic projection was found to the ventral thalamus. In *Ambystoma tigrinum*, Gruberg and Solish (1978) reported the presence of degenerating fibers in the ventral thalamus after spinal hemisections. The ventral thalamus also receives retinal, tectal and tegmental input (Wicht and Himstedt, 1988).

#### Ventral quadrant pathways in vertebrates

Ventral quadrant pathways comparable to those demonstrated in amphibians are found in other jawed vertebrates. This system of ascending fibers, sometimes called the spinal lemniscus after an ascending system described by Herrick (1948) in Ambystoma tigrium, arises in spinal neurons and courses rostrally through the ventralateral spinal cord and brain stem. All vertebrates have in common a distinct spinoreticular pathway passing via the ventral quadrant (Table 3). A ventral spinocerebellar pathways seems to be restricted to gnathostomes. A spinotectal component is variable. Ventral quadrant fibers extend as far as the diencephalon in terrestrial vertebrates and in certain sharks. The ventral quadrant pathways therefore include spinoreticular, spinocerebellar, spinotectal, and spinothalamic tracts.

In agnathans, ventral quadrant pathways extend rostrally, along the ventrolateral border of the spinal and medullary central grey and consists of spinoreticular and possibly spinovestibular projections (Ronan and Northcutt, 1990). In the Pacific hagfish, Eptatretus stouti, spinal lemniscal fibers heavily terminate in the mesencephalic tectum, but no spinal projection to the thalamus was found. The spinal lemniscus of lampreys ascends to the isthmus level and may extend into the mesencephalic tegmentum. Retrograde tracer data (Ronan and Northcutt, 1990) indicate that a very small population of cells in the most rostral part of the lamprey spinal cord may project to the tectum and diencephalon. Unlike the spinocerebellar projections of jawed vertebrates which terminate in the cerebellar granule layer, spinal lemniscal fibers in lampreys do not enter the periventricular grey of the cerebellar region (Ronan and Northcutt, 1990).

Ventral quadrant fibers project to the reticular formation, particularly its caudal part, the cerebellar granule layer, and the tectum mesencephalic in the spotted dogfish, *Scyliorhinus canicula* (Hayle, 1973a,b) and the nurse shark, *Ginglymostoma*  cirratum (Ebbesson and Hodde, 1981). In the latter, advanced species, spinal fibers also reach a subtectal, intercollicular zone and the central nucleus of the thalamus. Spinoreticular fibers passing via the ventral quadrant of the spinal cord were demonstrated in representatives of three grades of ray-finned bony fishes: bichirs, Polypterus palmas (unpublished observations quoted by Ronan and Northcutt, 1990), gars, Lepisosteus osseus (unpublished observations quoted by Ronan and Northcutt, 1990), rudds, Scardinius erythrophtalmus (Hayle, 1973a,b), and one genus of lobe-finned bony fish, the African lungfish, *Protopterus* spp. (unpublished observations quoted by Ronan and Northcutt, 1990). The spinoreticular projections diminish rostrally and likely terminate most heavily in the caudal reticular formation. In the ray-finned fishes and probably in African lungfishes, spinal fibers project to the cerebellar granule layer, whereas a projection to the tectum was only observed in lungfishes (unpublished observations quoted by Ronan and Northcutt, 1990). In Sebastiscus marmoratus, Murakami and Ito (1985) showed ascending projection of the spinal dorsal horn, via the ventral quadrant of the cord, to the reticular formation, the vagal lobe, octaval nuclei, the cerebellum and the nucleus ventromedialis thalami. Ito et al. (1986) suggested that the spinothalamic projection may arise in the nucleus of the lateral funiculus and, therefore, form a cervicothalamic rather than a proper spinothalamic pathway. In the himé salmon, Oncorhynchus nerka, Oka et al. (1986) found, in addition to spinoreticular and spinocerebellar components of the spinal lemniscus, a projection to the lateral part of the torus semicircularis.

Ascending ventral quadrant pathways in reptiles were studied in a number of species - a boid

snake, Constrictor constrictor (Ebbesson, 1969); turtles, Pelusios subniger, Pelomedusa subrufa, Podocnemis unifilis (Pedersen, 1973), and, especially, Pseudemys scripta elegans (Ebbesson, 1969; Künzle and Woodson, 1982); lizards, a lacertid, Lacerta viridis (Goldby and Robinson, 1962), the tegu lizard, Tupinambis teguixin (Ebbesson, 1967), and the savannah monitor lizard, Varanus exanthematicus (Hoogland, 1981; ten Donkelaar et al., 1987), and caimans, Caiman crocodilus (Ebbesson and Goodman, 1981; Pritz and Stritzel, 1989). The bulk of the ventral quadrant fibers innervate the rhombencephalic reticular formation, particularly its caudal part. Other rhombencephalic centers such as the nucleus of the solitary tract, the vestibular nuclear complex, the perihypoglossal nuclear complex, the inferior olive, and the area in and around the facial motor nucleus are innervated by spinal fibers. The cerebellum receives an extensive spinal innervation (for retrograde tracer data see: Bangma and ten Donkelaar, 1982; Künzle, 1983). In the mesencephalon, the nucleus intercollicularis, a diffuse cell mass in the dorsolateral part of the caudal mesencephalon, receives an extensive spinal innervation (Ebbesson, 1967, 1969; ten Donkelaar et al., 1987). Evidence for spinal projections to the intercollicular nucleus was found in various vertebrates (RoBards et al., 1976) and is included in Table 3. In mammals, e.g., the North American opossum (Hazlett et al., 1972; RoBards et al., 1976), the intercollicular terminal zone receives afferents from the spinal cord, the dorsal column nuclei, and the somatosensory cortex. In lizards, so far only spinal afferents and a projection from the dorsal column nucleus (Ebbesson, 1978) were demonstrated. Another main mesencephalic target of spinal afferents in reptiles is the laminar nucleus of the torus semicircularis. This cell group

may be comparable to the mammalian periaqueductal grey (ten Donkelaar et al., 1987; Puelles et al., 1994).

Anterograde degeneration studies in lizards (Ebbesson, 1967; Kusuma, 1979; ten Donkelaar et al., 1987) showed the presence of a direct spinothalamic projection, terminating in a poorly differentiated region labeled dorsal intermediate nucleus (Ebbesson, 1967) of the dorsal thalamus. This nucleus may be comparable to the intralaminar nuclei of mammals (Ebbesson, 1967), which receive the so-called paleospinothalamic pathway (Mehler, 1957; Herrick and Bishop, 1958). In Caiman crocodilus, Ebbesson and Goodman (1981) demonstrated in addition to the dorsal spinothalamic tract (terminating in the nucleus medialis posterior), a ventral extension of the spinal lemniscus. This ventral pathway terminates in the ventrolateral thalamic nucleus. This connection is reminiscent of the mammalian spinothalamic projection (neospinothalamic tract) to the ventrobasal complex (e.g., Mehler, 1957, 1969). In a retrograde tracer study in Varanus, exanthematicus, Hoogland (1981) showed that the spinal cord projects to three thalamic areas: 1) an area that includes both the nucleus dorsolateralis and the nucleus intermedius dorsalis; b) the nucleus ventrolateralis; and c) the nucleus dorsomedialis. Moreover, it was demonstrated that these spinothalamic projections arise from different populations of neurons in the spinal cord. Comparable data were obtained in Pseudemys scripta elegans (Künzle and Woodson, 1982; Künzle and Schnyder, 1983).

In pigeons (*Columba livia*), ascending spinal projections were studied with anterograde degeneration (Karten, 1963; Karten and Rezvin, 1966;van den Akker, 1970), tract-tracing (Wild, 1983; Funke and

Necker, 1986; Necker, 1989, 1990; Schneider and Necker, 1989) and electrophysiological (Delius and Bennetto, 1972; Necker, 1989; Funke, 1989a,b) techniques. Funke and Necker (1986) and Necker (1989, 1990) found that ventral quadrant fibers mainly arise in neurons in the laminae V-VIII of the spinal gray, bilaterally at cervical levels, and contralaterally at lumbar levels. Ascending spinal projections to the raphe and medial reticular targets arise from neurons located in all spinal laminae, except II and IV (Necker, 1989; Schneider and Necker, 1989). In experiments with large HRP applications to the lateral aspect of the reticular formation labeled neurons were found bilaterally, in lamina I and, to a lesser extent, in laminae IV and IX (Necker, 1989). Spinocerebellar fibers form a distinct component of the ventral quadrant fibers (van den Akker, 1970; Vielvoye, 1977; Okado et al., 1987). Sparse data exist in the literature on the spinomesencephalic projections in birds. Spinal projections were demonstrated to the intercollicular nucleus (Necker, 1989) and to the tectum (Karten, 1963). Spinothalamic projections were already observed in degeneration studies (Karten, 1963; Karten and Rezvin, 1966). Later tracing studies revealed lumbar spinothalamic projections to the nucleus dorsointermedius ventralis anterior (DIVA) and to a lesser extent to the nucleus intercalatus thalami, nucleus subrotundus, and the stratum cellulare externum and internum (Schneider and Necker, 1989). whereas cervical spinothalamic projections reach the nucleus dorsolateralis posterior (Schneider and Necker, 1989). The cells of origin of spinothalamic projections were demonstrated in several studies (Wild, 1983; Necker, 1989; Schneider and Necker, 1989). Schneider and Necker (1989) labeled only contralateral lumbar neurons in the intermediate grey in experiments with WGA-HRP applications in DIVA.

Necker (1989) labeled neurons at the base of the dorsal horn, close to the midline, and at the lateral aspect of the neck of the dorsal horn, in dorsolateral thalamic HRP application experiments, and only in those cases with large applications, labeled neurons were observed in the contralateral lumbar laminae V-VI. On the other hand, HRP applications into the nucleus ovoidalis labeled contralateral neurons in the laminae V-VI and VIII.

In mammals, the presence of extensive ventral quadrant pathways is well known (see Mehler, 1969; Willis and Coggeshall, 1991; Tracey, 1995). It includes spinoreticular, spinomesencephalic and spinothalamic projections. Retrograde tracer studies (Kevetter et al., 1982; Kevetter and Willis, 1984; Menétrey et al., 1984; Lima, 1990) showed three main groups of spinoreticular neurons: 1) those projecting to the lateral reticular nucleus; 2) a group projecting to the medial nuclei of the pontomedullary reticular formation, and 3) neurons that innervate the dorsal reticular nucleus. Spinal projections in mammals to the medial reticular formation arise bilaterally, but with a contralateral predominance, in neurons located mainly in laminae V, VII and VIII and, to a lesser extent, in lamina X and at the neck of the dorsal horn, throughout the spinal cord, but more densely at higher spinal segments, where also large ventral spinal neurons give rise to spinoreticular projections. The dorsal reticular nucleus receives projections from lamina I and X as well.

Although some spinomesencephalic projections in mammals ascend in the dorsolateral funiculus, the bulk of the spinal projections to the mesencephalon ascends via the ventrolateral quadrant intermingled with spinoreticular and spinothalamic axons. A rough rostrocaudal somatotopic arrangement was observed in the spinal projections to the mesencephalon. Lumbosacral projections reach more caudal mesencephalic targets than those that arise at cervical levels (Willis and Coggeshall, 1991). Spinomesencephalic projections innervate, among other structures, the nucleus cuneiformis, the parabrachial nucleus, the periaqueductal grey, the intercollicular nucleus and the deep layers of the superior colliculus. The cells of origin of spinomesencephalic projections are located throughout the spinal cord, but in higher concentration at higher spinal levels, in laminae I, III-V, VII, and X, and in the lateral cervical and lateral spinal nuclei (see Willis and Coggeshall, 1991).

The spinothalamic tract in mammals arises from neurons throughout the spinal cord. However, together with the lateral cervical nucleus neurons that also project to the thalamus, the highest concentration of spinothalamic neurons is present in cervical segments (Willis and Coggeshall, 1991). In addition to the spinothalamic projections via the dorsolateral funiculus (dorsolateral spinothalamic tract) arising from laminae I, II and III neurons to the ventral posterolateral and the submedius nuclei and the intralaminar nuclear complex, in the brain of marsupials, carnivores, rodents and primates two distinct components of spinothalamic projections via the ventral quadrant exist, 1) a mainly contralateral projection from laminae IV-VI neurons to the ventral posterolateral and the central lateral nuclei, and 2) a projection from laminae VII-X neurons to the contralateral posterior nuclear complex and the central lateral nucleus (Willis and Coggeshall, 1991).

Kevetter and Willis (1984) reviewed the then available data on the collateralization of the ascending projections in the ventral quadrant of the mammalian spinal cord as well as the formation of direct spinothalamic projections throughout phylogeny. They showed spinal collateral projections to the medial (intralaminar) and lateral (ventrobasal) thalamus considered as paleospinothalamic and neospinothalamic tracts, respectively (Mehler, 1969). Additionally, ascending collateral projections to the spinal cord and the medullary reticular formation, thalamus and reticular formation, intralaminar thalamus, ventrobasal thalamus and reticular formation were reported in different mammalian species (Kevetter and Willis, 1984). Moreover, later studies demonstrated that spinal neurons projecting to the mesencephalic periaqueductal gray send axon collaterals to the medullary reticular formation (Pechura and Liu, 1986) or the ventrobasal complex of the thalamus (Harmann et al., 1988). In our amphibian material we could observe thin fibers that innervate the medullary reticular formation as collateral branches of thicker ascending fibers located peripherally in the rhombencephalon that ascend to more rostral regions. However, retrograde double labeling studies are needed to confirm whether these fibers reach more rostral mesencephalic or diencephalic targets. As far as we know, so far no double-labeling studies demonstrated the collateralization of ascending spinal projections in non-mammalian vertebrates.

Based on previous anterograde degeneration (Hayle, 1973a; Ebbesson, 1969; Ebbesson and Hodde 1981) and tracing (Neary and Wilczynski, 1977, 1979) studies, direct spinothalamic projections, althought present in cartilaginous fishes (Ebbesson and Hodde, 1981), were viewed by Kevetter and Willis (1984) as an advanced evolutionary character only present in the brains of amniotes. The spinothalamic projection in some elasmobranchs was viewed as a secondary obtained character. However, later studies in agnathans (Ronan and Northcutt, 1990), teleosts (Murakami and Ito, 1985; Ito et al., 1986) and amphibians (A. Muñoz et al., 1994a; the present study) confirmed the existence of direct spinothalamic projections suggesting a common pattern throughout all the vertebrate classes.

In reptiles, the existence of a paleospinothalamic pathway to the dorsolateral and the intermediodorsal thalamic nuclei, the probable homologue of the intralaminar and posterior thalamus of mammals, and a neospinothalamic pathway to the ventrolateral thalamus, equivalent to the mammalian ventrobasal complex were shown (Ebbesson, 1967, 1978; Riss et al., 1972; Pedersen, 1973; Northcutt and Pritz, 1978; Ebbesson and Goodman, 1981; Hoogland, 1981) that arise from different spinal neuronal populations (Hoogland 1981). Our amphibian material (A. Muñoz et al., 1994a; the present study) revealed spinal projections to the ventral thalamus and, more sparsely, to different dorsal thalamic nuclei allowing comparison to the spinothalamic projections demonstrated in the brains of reptiles and mammals. In reptiles, spinal fibers that innervate dorsal and ventral thalamic nuclei enter the diencephalon dorsomedially or ventrolaterally, respectively (Ebbesson, 1967; Pedersen, 1973; Hoogland, 1981). Spinothalamic fibers to the ventral thalamus in amphibians enter the diencephalon ventrolaterally (A. Muñoz et al., 1994a; the present study). However, due to the low density of spinal fibers present in the dorsal thalamus, the way of entrance of these fibers could not be determined. However, since some fibers were observed at the

pretectal gray at the mesodiencephalic junction it seems likely that they enter the diencephalon dorsally. Additionally, different spinal neuronal populations giving rise to dorsal and ventral thalamic projections could not be distinguished since tracer applications restricted to dorsal thalamic nuclei failed to label neurons at the spinal cord (Neary and Wilczynski, 1977; 1979, own unpublished observations), likely due to the low density of dorsal spinothalamic projection. Studies in amphibians and reptiles, therefore strongly suggest the presence of ventral thalamic nuclei reminiscent of the mammalian ventrobasal complex. Their ascending projections are only incompletely known, and form the subject of future studies.

### ACKNOWLEDGMENTS

This research was supported by a grant of the Spanish Government to A González (DGICYT PB 93-0083) and by a NATO Collaborative Research Grant (930542) to H.J. ten Donkelaar. The authors thank Mrs. Roelie de Boer-van Huizen for excellent technical assistance, Mrs. Riet Schraven and Ms. Inge Eijkhout for secretarial assistance. ÷

:

Species	Target at obex level	Octavolateral target	Cerebellar projection	PDCS	 Rcís.
t	(DCN)				
Agnathans Lamaceur					
- Sea lamproy	+	?	+	?	19, 23
(Petromyzon marinus)				2	2 23
<ul> <li>Suver tamprey</li> <li>(Ichthuomaga unicutnit)</li> </ul>	, <b>+</b>	r	Ŧ		
(Schligengin Dicespe) Hagfisher Redfin bestick	, 	,	2	7	23
(Epiatretus stouti)	Ŧ	•			
Gasthestomes					
- Spotted doglish	7	?	?	?	9
(Scyliorhinus canicula)				2	7
- Nurse shark	+ ••	+	+	,	•
Bony fishes	,				
- Bichir	+	+	•	4	23
(Polypierus palmas)	ــــــــــــــــــــــــــــــــــــ	+		?	23
- Oat (Leoisosteus osseus)	•	·			
- Himé saimon	.+	•	•	, ?	20
(Oncorinynchus nerka)			-	,	23
- Lunglish (Protocient species)	+	•	Ŧ	•	~
Amphibians				•	
Urodeles			•	1	74
- Lungless salamanders	+	+	4	ł	24
- Axoloti	+	+	•	•	18
(Ambystoma mexicanun	n)				17
- Ribbed newt	+	+	+	+	• •
(Pieuroaeies waiii) Anurans					
<ul> <li>Rana esculenta</li> </ul>	+	+	+	?	1, 25
- R. temporaria	+	+	+	?	9, 10
- 10 10110101				_	
- R. catesbeiana	+	+	-+	?	4, 5
– R. perezi	+	+	+	+	16, 17
– Xenapus laevis	+	+	+	+	16, 17, 26
Reptiles					
Turtles	<u>ـ</u>	ـ	+	7	21
- Letanos inpunter	•	•			-
- Pelomedusa subrufa	+	+	+	?	21
– Podocnemis unifilis	+	+	+	,	21
- Pseudemys scripta	+	+	+	?	4, 14
elegans					
<u>i izaros</u> - Lacerto vieldis	<b>_</b>	<b>_</b>	+	,	11
	•	•	•	•	
- Tupinambis teguizin	+	+	+	?	3
Snakes					
- Constrictor constrictor	+	+	+	?	4
- Thanmophis sirtalis	+	+	•	?	12
Crocodilians					
- Caiman crocodilus	+	+.	+	+	6, 22
Birde		•			
- Pigeon	+	•	•	+	8, 13, 27, 28
(Columba livia) Mammals	+	*	-	÷	29

Table 1 Dorsal funicular pathways in vertebrates (based in part on Ronan and Northcutt, 1990)

Symbols used: + present; - not reported; ? unknown or questionable. References: 1 - Antal et al., 1980; 2 - Dubuc et al., 1993; 3 - Ebbesson, 1967; 4 - Ebbesson, 1969; 5 - Ebbesson, 1976; 6 -Ebbesson and Goodman 1981; 7 - Ebbesson and Hodde, 1981; 8 - Funke, 1988; 9, 10 - Hayle, 1973a,b; 11 - Jacobs, 1968; 12 -Jacobs and Sis, 1980; 13 - Karten, 1963; 14 - Kanzle, 1982; 15 - Künzle and Woodson, 1963; 16 - A. Muñoz et al., 1995b; 17 -A. Muñoz et al., present study; 18 - Nieuwenhuys and Corneliaz, 1971; 19 - Northcutt and Ebbesson, 1980; 20 - Oka et a., 1986; 21 - Pedersen, 1973; 22 - Pritz and Stritzel, 1994; 23 - Ronan and Northcutt, 1990; 24 - Roth and Wake, 1985; 25 -Szelely et al., 1980; 26 - ten Donkeisar and de Boer-van Huizen, 1991; 27 - van den Akker 1970; 28 - Wild, 1985; 29 - Willis and Coggeshall, 1991.

,

## Table 2 Dorsolateral funicular pathways in vertebrates

Species	Spinocervical tract	Target at spinomedullary junction (LCN)	Projections to			Refs.
			nucleus of the solitary tract	lateral reticular zone	para- brachiat	
Agnethans		(		20110	arca	
Lampreys						
- Sea lamprey	•	•	?	?	?	18, 19
(rearomyzon marinus)		•				
<ul> <li>Suver tamprey</li> <li>(Ichthormoron unicurpic</li> </ul>	•	7	?	?	• •	19
Hawishes	,					
- Pacific haefish	•	2	•	•	-	
(Epiatretus siouti)		•	r	2	7	19
Gnathostomes						
Cartilaginous fishes						
<ul> <li>Spotted dogfish</li> </ul>	•	•	•	•	-	7
(Scyliorhinus canicula)						
- Nurse shark		?	?	?	?	3
(Ginglymostoma cirratun	n)					-
Bony Jisher						
- Sca robin	+	?	?	?	?	4
(Prianolus carolinus)						
– Sebasiiscus marmoratus	7	+	?	?	?	10
- Rudd	•		•			70
(Scardinius erythrophthal	mus)				-	1,0
Amphibians						
- Tiger salamander	+	?	?	?	2	o
(Ambystoma tigrinum)						
- Ribbed newt	+	+	+	+	+	15
(Pleurodeles walt)						
+ Clawed load	+	+	+	+	+	12-15
(Xenopus idevis)						
- Large green frog	+	+	+ •	+	+	12, 13, 15
Rentiles						
- Red-cared turtle	· .	•				
(Preudemys seriosa elegan		ł	+	•	?	2, 11
- Tegu lizard	=, _	L				
(Tupinambis tentirin)	Ŧ	Ŧ	+	7	•	1
Birds						
- Pigeon	+	+	+	•	•	
(Columba livia)	·	•	Ŧ	r	7	5, 6, 16, 17, 20
Mammals	+	+	+	+	<b>_</b>	<b>1</b> 1
			•	r	Ŧ	41

Symbols used: + positive evidence; \* suggestive evidence; - not reported; ? unknown or questionable. References: 1 - Ebbesson, 1967; 2 - Ebbesson, 1969; 3 - Ebbesson and Hodde, 1981; 4 - Finger, 1981; 5 - Funke, 1988; 6 -Funke and Necker 1986; 7, 8 - Hayle, 1973a,b; 9 - Herrick, 1930; 10 - Ito et al., 1986; 11 - Künzle and Woodson, 1982; 12-14, A. Muñoz et al., 1995a,b, 1996; 15 - A. Muñoz et al., present study; 16, 17 - Necker 1989, 1991; 18 - Northcutt and Ebbesson, 1980; 19 - Ronan and Northcutt, 1990; 20 - van den Akker, 1970; 21 - Willis and Coggeshall, 1991.

Species :	Spinoreticular component	Spinocerebellar component	Projection to torus or PAG	Spinotectal component	Spinothalamic component	Reis.
Agnathans						
Lampreys						20.22
<ul> <li>Sca and silver lampreys</li> <li>Hawlisher</li> </ul>	+	?	•	•	·	20, 22
- Pacific hagfish	+	-		+		22
(Epiatretus siousi)						
Gnathostomes						
Cartilaginous fishes			•			7 8
(Scyliorhinus canicula)	•	+	ľ	Ŧ	•	7, 0
- Nurse shark	+	+	?	+	+	5
(Ginglymostoma cirratu	ni)					
Bony fishes		+	1			77
(Polypienus palmas)	+	+	:	•	•	44
- Gar	+	+	?	-		22
(Lepisosteus osseus)						
- Himé salmon	+	+	+	•	•	21
- Sebatiliseut marmoranus	· +	+	-		+	17
- ocoasiscus municontain.	· . ·	r.	Ŧ	•		••
- Lungfish	+	+	7	+		22
(Protopierus species)						
Amphibians				•		
	+	<b>_</b>	2	,		10
(Anibystoma mexicanur	n)	·	÷	•	-	
- Tiger salamander	+	+	+	+	+	9
(A. tigrinum)						
<ul> <li>Ribbed newt</li> <li>(Plaurodalas waid)</li> </ul>	+	+	+	+	+	15, 16
Anurans			:			
- Rana temporaria	+	+	?	• .	•	7, 8
						•
- R. catesbeiana	+	+	+	•	?	3
- R. perezi	+	+	+	+	+	15, 16
- Xenopus laevis	+	+	+	+	+	16
Reptiles						
Turtles			•			
- rseudemys scripta elega	ins +	+	+	+	+	2, 12, 13
Lizards						
- Tupinambis teguizin	+	+	+	+	+	1
- Varanus exanthematicu	s +	+	+	+	+	10, 24
Snakes						
- Constructor constructor	+	+	+	+	+	2
Crocodilians						
- Caiman crocodilus	+	+	+	+	+	4
Birds						
- Pigcon	+	+	•. +	+	+	6, 11, 18, 23
(Columos livia) Mammali	+	<u>ـ</u>	يد	+	Ŧ	14 25
1 4 Mar 47 144 M				T	Ŧ	14. 67

#### Table 3 Ventral quadrant pathways in vertebrates (based in part on Ronan and Northcutt, 1990)

Symbols used: + positive evidence; \* suggestive evidence; • not reported; ? unknown or questionable. References: 1, 2, 3 - Ebbesson, 1967, 1969, 1976; 4 - Ebbesson and Goodman, 1981; 5 - Ebbesson and Hodde, 1981; 6 - Funke and Necker 1986; 7, 8 - Hayle, 1973a,b; 9 - Herrick, 1930; 10 - Hoogland, 1981; 11 - Karten, 1963; 12 - Künzle and Schnyder, 1983; 13 - Künzle and Woodson, 1982; 14 - Mehler, 1969; 15 - A. Muñoz et al., 1994a; 16 - A. Muñoz et al., present study; 17 - Murakami and Ito, 1985; 18 - Necker, 1989; 19 - Nieuwanhuys and Cornelisz, 1971; 20 - Northcutt and Ebbesson, 1980; 21 - Oka et al 1986; 22 - Ronan and Northcutt, 1990; 23 - Schneider and Necker, 1989; 24 - ten Donkelaar et al., 1987; 25 -Willis and Coggeshall, 1995.

ı,



Figure 1: Schematic drawing of a series of transverse sections through the brain stem and spinal cord of a young adult *Xenopus laevis* showing the labeling in the dorsal and dorsolateral funiculi after *in vitro* application of 3kD BDA into the lumbar spinal cord between the 9th and 10th dorsal roots.


Figure 2: Schematic drawing of a series of transverse sections through the brain stem and spinal cord of a young adult *Xenopus laevis* showing the labeling in the dorsal and dorsolateral funiculi after *in vitro* application of 3kD BDA into the cervical spinal cord, just rostral to the third dorsal root.



Figure 3: Schematic drawing of a series of transverse sections through the diencephalon (A-D), the mesencephalon (E-H), the rhombencephalon (I-S), and spinal cord (T-V) of *Rana perezi* showing the labeling of the ventral quadrant system after *in vivo* 10kD BDA application to the cervical spinal cord.

.



Figure 4: Photomicrographs illustrating the labeling observed in various BDA experiments in anurans. A, Ipsilateral labeling in the dorsal and dorsolateral funiculi as well as of some fibers of the descending tract of the trigeminal nerve, *Xenopus laevis*, thoracic *in vitro* 3kD BDA application; B, ipsilateral labeling of dorsolateral funicular fibers in the subcerebellar regions, *X. laevis*, cervical *in vitro* 3kD BDA application; C, ipsilateral labeling of dorsolateral funicular fibers at the level of the eighth cranial nerve root arborizing in the lateral reticular zone, *X. laevis*, cervical *in vitro* 3kD BDA application; D, ipsilateral labeling of dorsolateral funicular fibers at the level of the eighth cranial nerve root arborizing in the lateral reticular zone, *X. laevis*, cervical *in vitro* 3kD BDA application; D, ipsilateral labeling of dorsolateral funicular fibers at the level of the eighth cranial nerve root arborizing in the lateral reticular zone, *Rana perezi*, cervical *in vivo* 10kD BDA application; E, F, contralateral labeling of ventral quadrant fibers in the tectum mesencephali and torus semicircularis, respectively; *R. perezi*, cervical *in vivo* 10kD BDA application. Scale bars indicate 100 µm.



Figure 5: Schematic drawings illustrating the distribution of retrogradely labeled neurons in the spinal cord of *Xenopus laevis* following *in vitro* 3 kD BDA applications to the ventral thalamus (A), the torus semicircularis (B), and the inferior reticular nucleus (C). Examples of labeled neurons are shown in Fig. 6.

,



Figure 6: Examples of retrogradely labeled neurons contralaterally (A, B, D) and ipsilaterally (C, E, F) in the cervical spinal cord of *Xenopus laevis* after *in vitro* 3kD BDA applications to the ventral thalamus (A), inferior reticular nucleus (B, E) and torus semicircularis (C, D, F, G). Scale bars indicate 100 µm.



Figure 7: Schematic drawing of a series of transverse sections through the brain stem and spinal cord of *Pleurodeles waltl* showing the distribution of 10kD BDA labeled brachial dorsal root fibers. The levels of the sections are indicated along a dorsal view of the central nervous system of *P. waltl*. Insets show labeling in the ipsilateral dorsal column and lateral cervical nuclei (O), and in the cerebellum (P). Scale bars indicate 100  $\mu$ m.

13 4



Figure 8: Schematic drawing of a series of transverse sections through the brain stem and spinal cord of *Pleurodeles waltl* showing the labeling of dorsal and dorsolateral funicular components after an *in vivo* 10kD BDA application to the cervical spinal cod. The levels of the sections are indicated along a dorsal view of the central nervous system of *P. waltl*.

ŝ



Figure 9: Schematic drawing of a series of transverse sections through the diencephalon (A-C), mesencephalon (D, E), rhombencephalon (F-L), and spinal cord (M-O) of *Pleurodeles waltl* showing the labeling of the ventral quadrant system after *in vivo* 10kD BDA application to the cervical spinal cord.



Figure 10: Photomicrographs illustrating the labeling observed in various BDA experiments in *Pleurodeles* waltl. A, Ipsilateral labeling in the dorsal and dorsolateral funiculi in the caudal rhombencephalon after a cervical BDA application; B, ipsilateral labeling at the level of the nucleus reticularis medius, cervical BDA application; C, D and E show contralaterally labeled ventral quadrant fibers innervating the reticular formation (C), the tectum mesencephali (D) and the torus semicircularis (E). In F, a neuron retrogradely labeled from the ventral thalamus in the dorsal horn of the cervical spinal cord is shown. Scale bars indicate 100 µm.

#### LIST OF ABBREVIATIONS

A	anterior thalamic nucleus
Ad	anterodorsal tegmental nucleus
Av	anteroventral tegmental nucleus
С	central thalamic nucleus
с	central spinal field
cb	cerebellum
d	dorsal spinal field
dr (3-10)	dorsal root 3-10
	dorsal column nucleus
df df	dorsal funiculus
ы ПЦ	dorsal humothalamic nucleus
dh l	dorsal hom
जा तार	dorsolateral funiculus
un En	posterior entopeduncular pucleus
Lp Le	pusterior entopeduneurar nucleus
15	Interest spin at field
I I	lateral spinal field
La T	lateral malamic nucleus, anterior division
Lam	laminar nucleus of the torus semicircularis
LCN	lateral cervical nucleus
lm	lateral motor field
Lpd	lateral thalamic nucleus, posterodorsal
_	division
Lpv	lateral thalamic nucleus, posteroventral
	division
lrz	lateral reticular zone
Mag	magnocellular nucleus of the torus
	semicircularis
Mg	magnocellular preoptic nucleus
nV	nervus trigeminus
nVⅢ	nervus octavus
NPv	nucleus of the periventricular organ
Р	posterior thalamic nucleus
Pr	principal nucleus of the torus semicircularis
ptg	pretectal grey
ptrg	pretoral grey
Ri	nucleus reticularis inferior
Rm	nucleus reticularis medius
Rs	nucleus reticularis superior
SC	suprachiasmatic nucleus
tm	tectum mesencephali
tor	torus semicircularis
TP	posterior tuberculum
trVds	descending tract of the trigeminal nerve
VH	ventral hypothalamic nucleus
vh	ventral horn
VId	ventrolateral thalamic nucleus dorsal part
VLv	ventrolateral thalamic nucleus, ventral nart
vl	ventrolateral spinal field
VM	ventromedial thalamic nucleus
vm	ventromedial spinal field
vth	ventral thalamus
797	zona anteroventralis of the dorsal thalamus
2001 2001	zona posterodorealis of the doreal thalamus
ци Vde	nucleus of the descending treat of the
v us	trigeminel nerve
	u igenniai nerve

#### LITERATURE CITED

Apkarian, A.V., R.T. Stevens, and C.J. Hodge (1985) Funicular location of ascending axons of lamina I cells in the cat spinal cord. Brain Res. 334: 160-164.

Adams, J.C. (1981) Heavy metal intensification of DABbased HRP reaction product. J. Histochem. Cytochem. 29: 775.

Allison, J.D., and W. Wilczynski (1991) Thalamic and midbrain auditory projections to the preoptic area and ventral hypothalamus in the green treefrog (*Hyla cinerea*). Brain Behav. Evol. 38: 322-331.

Angaut-Petit, D. (1975a) The dorsal column system. I. Existence of long ascending postsynaptic fibres in the cat's fasciculus gracilis. Exp. Brain Res. 22: 457-470.

Angaut-Petit, D. (1975b) The dorsal column system. II. Functional properties and bulbar relay of the postsynaptic fibres of the cat's fasciculus gracilis. Exp. Brain Res. 22: 471-493.

Antal, M., I. Tornai, and G. Székely (1980) Longitudinal extent of dorsal root fibers in the spinal cord and brainstem of the frog. Neuroscience 5: 1311-1322.

Bangma, G.C., and H.J. ten Donkelaar (1982) Afferent connections of the cerebellum in various types of reptiles. J. Comp. Neurol. 207: 255-273.

Barnard, J.W. (1936) A phylogenetic study of the visceral afferent areas associated with the facial, glossopharyngeal and vagus nerves, and their fiber connctions. The efferent facial nucleus. J. Comp. Neurol. 65: 503-602.

Bennett, G.J., N. Nishikawa, G.-W. Lu, M.J. Hoffert, and R. Dubner (1984) The morphology of dorsal column postsynaptic spinomedullary neurons in the cat. J. Comp. Neurol. 224: 568-578.

Bernard, J.F., R. Dallel, P. Raboisson, L. Villanueva, and D. Le Bars (1995) Organization of efferent projections from the spinal cervical enlargement to the parabrachial area and the periaqueductal gray: A PHA-L study in the rat. J. Comp. Neurol. 353: 480-505.

Björkeland, M. and J. Boivie (1984) The termination of spinomesencephalic fibers in the cat. An experimental anatomical study. Anat. Embryol. 170: 265-277.

Blomqvist, A., and K.J. Berkley (1992) A re-examination of the spino-reticulo-diencephalic pathway in the cat. Brain Res. 579: 17-31.

Boivie, J. (1983) Anatomic and physiologic features of the spino-cervico-thalamic pathway. In G. Macchi, A. Rustioni and R. Spreafico (eds.) Somatosensory Integration in the Thalamus. Elsevier, Amsterdam, pp. 63-106.

Cochran, S.L., P. Kasik, and W. Precht (1987) Pharmacological aspects of excitatory synaptic transmission to second-order neurons in the frog. Synapse 1: 102-103.

Comer, C., and P. Grobstein (1981a) Tactually elicited acquisition prey behavior in the frog, *Rana pipiens*, and a comparision with visually elicited behavior. J. Comp. Physiol. 142: 141-150.

Comer, C., and P. Grobstein (1981b) Involvement of midbrain structures in tactually and visually elicited prey acquisition behavior in the frog. J. Comp. Physiol. 142: 151-160.

Comer, C., and P. Grobstein (1981c) Organization of sensory inputs to the midbrain of the frog, *Rana pipiens*. J. Comp. Physiol. 142: 161-168.

Dampney, R.A.L., J. Czachurski, K. Dembowsky, A.K. Googchild, and H. Seller (1987) Afferent connections and

spinal projections of the pressor region in the rostral ventrolateral medulla of the cat. J. Auton. Nerv. Syst 20: 73-86.

Delius, J.D., and K. Bennetto (1972) Cutaneous sensory projections to the avian forebrain., Brain Res. 37: 205-221.

Dubuc, R., F. Bongianni, Y. Ohta, and S. Grillner (1993) Anatomical and physiological study of brainstem nuclei relaying dorsal column inputs in lampreys. J. Comp. Neurol. 327: 260-270.

Ebbesson, S.O.E. (1967) Ascending axon degeneration following hemisection of the spinal cord in the tegu lizard (*Tupinambis nigropunctatus*). Brain Res. 5: 178-206.

Ebbesson, S.O.E. (1969) Brainstem afferents from the spinal cord in a sample of reptilian and amphibian species. Ann. N.Y. Acad. Sci. 167: 80-101.

Ebbesson, S.O.E. (1976) Morphology of the spinal cord. In R. Llinás and W. Precht (eds): Frog Neurobiology. New York: Springer Verlag, pp. 679-706.

Ebbesson, S.O.E. (1978) Somatosensory pathways in lizards: The identification of the medial lemniscus and related structures. In N. Greenberg and P.D. MacLean (eds.) Behavior and Neurology of Lizards. Natl. Inst. Mental Health, Rockville, pp. 91-104.

Ebbesson, S.O.E., and D.C. Goodman (1981) Organization of ascending spinal projections in *Caiman crocodilus*. Cell Tiss. Res. 215: 383-395.

Ebbesson, S.O.E., and K.C. Hodde (1981) Ascending spinal systems in the nurse shark *Ginglymostoma* cirratum. Cell Tiss. Res. 216: 313-331.

Ebbesson, S.O.E., J.A. Jane, and D.M. Schroeder (1972) A general overview of major interspecific variations in thalamic organization. Brain Behav. Evol. 6: 92-130.

Feil, K., and H. Herbert (1995) Topographic organization of spinal and trigeminal somatosensory pathways to the rat parabrachial and Kölliker-Fuse nuclei. J. Comp. Neurol. 353: 506-528.

Finger, T.E. (1981) Fish that taste with their feet: Spinal sensory pathways in the sea robin, *Prionotus carolinus*. Biol. Bull. 161: 343.

Finkenstädt, T., S.O.E. Ebbesson, and J.P. Ewert (1983) Projections to the midbrain tectum in *Salamandra* salamandra L. Cell Tiss. Res. 234: 39-55.

Forehand, C.J., and P.B. Farel (1982) Spinal cord development in anuran larvae. II. Ascending and descending pathways. J. Comp. Neurol. 209: 395-408.

Fritzsch, B. (1988) The lateral-line and inner-ear afferents in larval and adult urodeles. Brain Behav. Evol. 31: 325-348.

Funke, K. (1988) Spinal projections to the dorsal column nuclei in pigeons. Neurosci. Lett. 91: 295-300.

Funke, K. (1989a) Somatosensory areas in the telencephalon of the pigeon. I. Response characteristics. Exp. Brain Res. 76: 603-619.

Funke, K. (1989b) Somatosensory areas in the telencephalon of the pigeon. II. Spinal pathways and different connections. Exp. Brain Res. 76: 620-638.

Funke, K., and R. Necker (1986) Cells of origin of ascending pathways in the spinal cord of the pigeon. Neurosci. Lett. 71: 25-30.

Giesler, G.J., R.L. Nahin, and A.M. Madsen (1984) Postsynaptic dorsal column pathway of the rat. I. Anatomical studies. J. Neurophysiol. 51: 260-275.

Goldby, F., and L.R. Robinson (1962) The central connections of dorsal spinal nerve roots and the

ascending tracts in the spinal cord of Lacerta viridis. J. Anat. (Lond.) 96: 153-170.

González, A., and M. Muñoz (1987) Some connections of the area octavolateralis of *Pleurodeles waltlii*. A study with horseradish peroxidase under in vitro conditions. Brain Res. 423: 338-342.

González, A., H.J. ten Donkelaar, and R. de Boer-van Huizen (1984) Cerebellar connections in *Xenopus laevis*. An HRP study. Anat. Embryol. 169: 167-176.

González, A., A. Muñoz, and M. Muñoz (1993) Trigeminal primary afferent projections to the spinal cord of the frog, *Rana ridibunda*. J. Morphol. 217: 137-146.

González, A., and W.J.A.J. Smeets (1994) Catecholamine systems in the CNS of amphibians. In W.J.A.J. Smeets and A. Reiner (eds.) Phylogeny and Development of Catecholamine Systems in the CNS of Vertebrates. Cambridge University Press, Cambridge, pp. 77-102.

Grover, B.G., and U. Grüsser-Cornehls (1984) Cerebellar afferents in the frogs, *Rana esculenta* and *Rana temporaria*. Cell Tiss. Res. 237: 259-267.

Gruberg, E.R., and W.A. Harris (1981) The serotoninergic somatosensory projection to the tectum in normal and eyeless salamanders. J. Morphol. 170: 55-69.

Gruberg, E.R., and S.P. Solish (1978) The relationship of a monoamine fiber system to a somatosensory tectal projection in the salamander *Ambystoma tigrinum*. J. Morphol. 157: 137-150.

Harmann, P.A., S.M. Carlton, and W.D. Willis (1988) Collaterals of spinothalamic tract cells to the periaqueductal gray: A fluorescent double-labeling study in the rat. Brain Res. 441: 87-97.

Hayle, T.H. (1973a) A comparative study of spinal projections to the brain (except cerebellum) in three classes of poikilothermic vertebrates. J. Comp. Neurol. 149: 463-476.

Hayle, T.H. (1973b) A comparative study of spinocerebellar systems in three classes of poikilothermic vertebrates. J. Comp. Neurol. 149: 477-496.

Hazlett, J.C., R. Dom, and G.F. Martin (1972) Spinobulbar, spino-thalamic and medial lemniscal connections in the American opossum, *Didelphis marsupialis* virginiana. J. Comp. Neurol. 146: 95-118.

Herbert, H., M.M. Moga, and C.B. Saper (1990) Connections of the parabrachial nucleus with the nucleus of the solitary tract and the medullary reticular formation in the cat. Brain Res. 511: 540-580.

Herrick, C.J. (1914) The medulla oblongata of larval Amblystoma. J. Comp. Neurol 24: 343-427.

Herrick, C.J. (1930) The medulla oblongata of Necturus. J. Comp. Neurol 50: 1-96.

Herrick, C.J. (1948) The Brain of the Tiger Salamander, Amblystoma Tigrinum.

Herrick, C.J., and G.H. Bishop (1958) A comparative survey of the spinal lemniscus systems. In H.H. Jasper, C.D. Proctor, R.S. Knighton, W.C. Norway and R.T. Costello (eds.) Reticular Formation of the Brain. Little, Brown and Co., Boston, pp. 353-360.

Hoogland, P.V. (1981) Spinothalamic projections in a lizard, Varanus exanthematicus: An HRP study. J. Comp. Neurol. 198: 7-12.

Hylden, J.L.K., H. Hayashi, R. Dubner, and G.J. Bennett (1986) Physiology and morphology of the lamina I spinomesencephalic projection. J. Comp. Neurol. 247: 505-515.

Ito, H., T. Murakami, T. Fukuoka, and R. Kishida (1986) Thalamic fiber connections in a teleost (Sebastiscus *marmoratus*): Visual, somatosensory, octavolateral and cerebellar relay region to the telencephalon. J. Comp. Neurol. 250: 215-227.

Jacobs, V.L. (1968) An experimental study of the course and termination of the spino-cerebellar systems in a lizard (*Lacerta viridis*). Brain Res. 11: 154-176.

Jacobs, V.L., and R.F. Sis (1980) Ascending projections of the dorsal column in a garter snake (*Thamnophis sirtalis*): A degeneration study. Anat. Rec. 196: 37-50.

Jhaveri, S., and E. Frank (1983) Central projections of the brachial nerve in bullfrogs: Muscle and cutaneous afferents project to different regions of the spinal cord. J. Comp. Neurol. 221: 304-312.

Joseph, B.S., and D.G. Whitlock (1968a) Central projections of selected spinal dorsal roots in anuran amphibians. Anat. Rec. 160: 279-288.

Joseph, B.S., and D.G. Whitlock (1968b) Central projections of brachial and lumbar dorsal roots in reptiles. J. Comp. Neurol. 132: 469-484.

Karten, H.J. (1963) Ascending pathways from the spinal cord in the pigeon (*Columba livia*). Proc. XVIth Int. Congr. Zool., 2: 23.

Karten, H.J., and A.M. Rezvin (1966) The afferent connections of the nucleus rotundus in the pigeon. Brain Res. 2: 368-377.

Kevetter, G.A., and W.D. Willis (1984) Collateralization of the spinothalamic tract: New methodology to support or deny phylogenetic theories. Brain Res. Rev. 7: 1-14.

Kevetter, G.A., L.H. Haber, R.P. Yezierski, J.M. Chung, R.F. Martin, and W.D. Willis (1982) Cells of origin of the spinoreticular tract in the monkey, J. Comp. Neurol. 207: 61-74.

Kitamura, T., J. Yamada, H. Sato, and K. Yamashita (1993) Cells of origin of the spinoparabrachial fibers in the rat: A study with Fast Blue and WGA-HRP. J. Comp. Neurol. 328: 449-461.

Künzle, H. (1982) Dorsal root projections to the cerebellum in turtle. Exp. Brain Res. 45: 464-466.

Künzle, H. (1983) Spinocerebellar projections in the turtle. Observations on their origin and terminal organization. Exp. Brain Res. 53: 129-141.

Künzle, H., and H. Schnyder (1983) Do retinal and spinal projections overlap within the turtle thalamus? Neuroscience 10: 161-168.

Künzle, H., and W. Woodson (1982) Mesodiencephalic and other target regions of the ascending spinal projections in the turtle: *Pseudemys scripta elegans*. J. Comp. Neurol. 212: 349-364.

Künzle, H., and W. Woodson (1983) Primary afferent projections to the spinal cord and the dorsal column nuclear complex in the turtle *Pseudemys*. Anat. Embryol. 166: 229-245.

Kuru, M. (1956) The spino-bulbar tracts and the pelvic sensory vagus. Further contributions to the theory of the sensory dual innervation of the viscera. J. Comp. Neurol. 104: 207-231.

Kusuma, A. (1979) The Organization of the Spinal Cord in Reptiles with Different Locomotor Patterns. Thesis, University of Nijmegen (The Netherlands).

Kusuma, A., and H.J. ten Donkelaar (1980) Dorsal root projections in various types of reptiles. Brain Behav. Evol. 17: 291-309.

Larsell, O. (1923) The cerebellum of the frog. J. Comp. Neurol. 39: 89-112.

Lima, D. (1990) A spinomedullary projection terminating in the dorsal reticular nucleus in the rat. Neuroscience 34: 577-591. Sistemas somatosensoriales en anfibios PROYECCIONES ESPINALES ASCENDENTES 3.2

Luksch, H., W. Walkowiak, A. Muñoz, and H.J. ten Donkelaar (1996) The use of *in vitro* preparations of the isolated amphibian CNS in neuroanatomy and electrophysiology. J. Neurosci. Meth., submitted.

McMahon, S.B., and P.D. Wall (1983) A system of rat spinal cord lamina I cells projecting through the contralateral dorsolateral funiculus. J. Comp. Neurol. 214: 217-223.

Mehler, W.R. (1957) The mammalian 'pain tract' in phylogeny. Anat. Rec. 127: 332.

Mehler. W.R. (1969) Some neurological species differences - A posteriori. - Ann. N.Y. Acad. Sci. 167: 424-467.

Menétrey, D., and A.I. Basbaum (1987) Spinal and trigeminal projections to the nucleus of the solitary tract: A possible substrate for somatovisceral and viscerovisceral reflex activation. J. Comp. Neurol. 255: 439-450.

Menétrey, D., J. De Pommery, and J.M. Besson (1984) Electrophysiological characteristics of lumbar spinal cord neurons backfired from lateral reticular nucleus in the rat. J. Neurophysiol. 52: 595-611.

Morin, J. (1955) A new spinal pathway for cutaneous impulses. Amer. J. Physiol. 183: 245-252.

Muñoz, A., M. Muñoz, A. González, and H.J. ten Donkelaar (1994a) Spinothalamic projections in amphibians as revealed with anterograde tracing techniques. Neurosci. Lett. 171: 81-84.

Muñoz, A., M. Muñoz, A. González, and H.J. ten Donkelaar (1994b) The dorsal column-medial lemniscal projection of anuran amphibians. Eur. J. Morphol. 32: 283-287.

Muñoz, A., M. Muñoz, A. González, and H.J. ten Donkelaar (1995a) Ascending spinal projections in the dorsolateral funiculus of anuran amphibians. Eur. J. Neurosci., Suppl. 8: 151.

Muñoz, A., M. Muñoz, A. González, and H.J. ten Donkelaar (1995b) The anuran dorsal column nucleus: organization, immunohistochemical characterization and fiber connections in *Rana perezi* and *Xenopus laevis*. J. Comp. Neurol. 363: 1-25.

Muñoz, A., M. Muñoz, A. González, and H.J. ten Donkelaar (1996) Evidence for an anuran homologue of the mammalian spinocervicothalamic system: An *in vitro* tract-tracing study in *Xenopus laevis*. Eur. J. Neurosci., submitted.

Muñoz, M., A. Muñoz, A. González, C. Machín, and C. Rúa (1991) Ascending connections from the spinal cord in the anuran amphibian Rana ridibunda. Eur. J. Neurosci., Suppl. 4: 24.

Muñoz, M., A. Muñoz, O. Marín, and A. González (1994) Primary afferents and second order projections of the trigeminal system in a frog (*Rana ridibunda*). Eur. J. Morphol. 32: 288-292.

Murakami, T., and H. Ito (1985) Long ascending projections of the spinal dorsal horn in a teleost, *Sebastiscus marmoratus*. Brain Res. 346: 168-170.

Neary, T.J. (1988) Forebrain auditory pathways in ranid frogs. In B. Fritzsch, M.J. Ryan, W. Wilczynski, T.E. Hetherington and W. Walkowiak (eds.) The Evolution of the Amphibian Auditory System. Wiley, New York, pp. 233-252.

Neary, T.J. (1995) Afferent projections to the hypothalamus in ranid frogs. Brain Behav. Evol. 46: 1-13.

Neary, T.J., and R.G. Northcutt (1983) Nuclear organization of the bullfrog diencephalon. J. Comp. Neurol. 213: 262-278.

Neary, T.J., and W. Wilczynski (1977) Ascending thalamic projections from the obex region in ranid frogs. Brain Res. 138: 529-533.

Neary, T.J., and W. Wilczynski (1979) Anterior and posterior thalamic afferents in the bullfrog *Rana* catesbeiana. Soc. Neurosci. Abstr. 5: 144.

Neary, T.J., and W. Wilczynski (1986) Auditory pathways to the hypothalamus in ranid frogs. Neurosci. Lett. 71: 142-146.

Necker, R. (1989) Cells of origin of spinothalamic, spinotectal, spinoreticular, and spinocerebellar pathways in the pigeon as studied by the retrograde transport of horseradish peroxidase. J. Hirnforsch. 30: 33-43.

Necker, R. (1990) Cells of origin of ascending and descending as well as branching fibers in the cervical spinal cord of the pigeon. Neurosci. Lett. 119: 1-4.

Necker, R. (1991) Cells of origin of avian postsynaptic dorsal column pathways. Neurosci. Lett. 126: 91-93.

Nieuwenhuys, R., and M. Cornelisz (1971) Ascending projections from the spinal cord in the axolotl (Ambystoma mexicanum). Anat. Rec. 169: 388.

Nijensohn, D.E., and F.W.L. Kerr (1975) The ascending projections of the dorsolateral funiculus of the spinal cord in the primate. J. Comp. Neurol. 161: 459-470.

Nikundiwe, A.M., and R. Nieuwenhuys (1983) The cell masses in the brainstem of the South African clawed frog *Xenopus laevis*: A topographical and topological analysis. J. Comp. Neurol. 213: 199-219.

Nikundiwe, A.M., R. de Boer van Huizen, and H.J. ten Donkelaar (1982) Dorsal root projections in the clawed toad (*Xenopus laevis*) as demonstrated by anterograde labeling with horseradish peroxidase. Neuroscience 7: 2089-2103.

Northcutt, R.G., and S.O.E. Ebbesson (1980) Ascending spinal pathways in the sea lamprey. Soc. Neurosci. Abstr. 6: 628.

Northcutt, R.G., and M.B. Pritz (1978) A spinothalamic pathway to the dorsal ventricular ridge in the spectacled caiman *Caiman crocodilus*. Anat. Rec. 190: 618-619.

Oka, Y., M. Satou, and K. Ueda (1986) Ascending pathways from the spinal cord in the Himé salmon (landlocked red salmon, *Oncorhynchus nerka*) J. Comp. Neurol. 254: 104-112.

Okado, N., R. Ito, and S. Homma (1987) The terminal distribution pattern of spinocerebellar fibers. An anterograde labelling study in the posthatching chick. Anat. Embryol. 176: 175-182.

Opdam, P., and R. Nieuwenhuys (1976) Topological analysis of the brain stem of the axolotl Ambystoma mexicanum. J. Comp. Neurol. 165: 285-306.

Opdam, P., M. Kemali, and R. Nieuwenhuys (1976) Topological analysis of the brain stem of the frogs *Rana* esculenta and *Rana catesbeiana*. J. Comp. Neurol. 165: 307-332.

Pechura, C., and R. Liu (1986) Spinal neurons wich project to the periaqueductal gray in the medullary reticular formation via axon collaterals: A double-label fluorescence study in the rat. Brain Res. 374: 357-361.

Pedersen, R. (1973) Ascending spinal connections in three species of side-necked turtle (Podocnemis unifilis, Pelusios subniger, Pelomedusa subrufa). Anat. Rec. 175: 409.

Potter, H.D. (1965) Mesencephalic auditory region of the bullfrog. J. Neurophysiol. 28: 1132-1154.

Pritz, M.B., and M.E. Stritzel (1989) Reptilian somatosensory midbrain: Identification based on input from the spinal cord and dorsal column nucleus. Brain Behav. Evol. 33: 1-14.

Pritz, M.B., and M.E. Stritzel (1994) Anatomical identification of a postsynaptic dorsal column system in a reptile, *Caiman crocodilus*. Brain Behav. Evol. 43: 233-243.

Puelles, L., C. Robles, M. Martínez-de-la-Torre, and S. Martínez (1994) New subdivision schema of the avian torus semicircularis: Neurochemical maps in the chick. J. Comp. Neurol. 340: 98-125.

Rettig, G. (1984) Neuroanatomische Untersuchungen der Visuellen Projektionen bei Salamandern (Ordnung Caudata). Thesis, University of Bremen (Germany).

Rettig, G. (1988) Conecctions of the tectum opticum in two urodeles (*Salamandra salamandra* and *Bolitoglossa subpalmata*) with a special reference to the nucleus isthmi. J. Hirnforsch. 29: 5-16.

Riss, W., R.A. Pedersen, J.S. Jakway, and C.B. Ware (1972) Levels of function and their representation in the vertebrate thalamus. Brain Behav. Evol. 6: 26-41.

RoBards, M.J., D.W. Watkins, and R.B. Masterton (1976) An anatomical study of some somesthetic afferents to the intercollicular terminal zone of the midbrain of the opossum. J. Comp. Neurol. 170: 499-524.

Ronan, M., and R.G. Northcutt (1990) Projections ascending from the spinal cord to the brain in petromyzontid and myxinoid agnathans. J. Comp. Neurol. 291: 491-508.

Ross, C.A., D.A. Ruggiero, T.H. Joh, D.H. Park, and D.J. Reis (1984) Rostral ventrolateral medulla: Selective projections to the thoracic autonomic cell column from the region containing C1 adrenaline neurons. J. Comp. Neurol. 228: 168-185.

Roth, G., and D.B. Wake (1985) The structure of the brainstem and cervical spinal cord in lungless salamanders (Family Plethodontidae) and its relation to feeding. J. Comp. Neurol. 241: 99-110.

Rustioni, A. (1973) Non-primary afferents to the nucleus gracilis from the lumbar cord of the cat. Brain Res. 51: 81-95.

Rustioni, A. and A.B. Kaufman (1977) Identification of cells of origin of non-primary afferents to the dorsal column nuclei of the cat. Exp. Brain Res. 27: 1-14.

Saper, C.B. (1995) The spinoparabrachial pathway: Shedding new light on and old path. J. Comp. Neurol. 353: 477-479.

Schneider, A., and R. Necker (1989) Spinothalamic projection in the pigeon. Brain Res. 484: 139-149.

Slugg, R.M., and A.R. Light (1994) Spinal cord and trigeminal projections to the pontine parabrachial region in the rat as demonstrated with Phaseolus vulgaris-leucoagglutinin. J. Comp. Neurol. 339: 49-61.

Sternberg, L.A. (1979) Immunocytochemistry. Willey, New York

Székely, G., M. Antal, and T. Görcs (1980) Direct dorsal root projection onto the cerebellum in the frog. Neurosci. Lett. 19: 161-165.

ten Donkelaar, H.J., and R. de Boer-van Huizen (1991) Observations on the development of ascending spinal pathways in the clawed toad, *Xenopus laevis*. Anat. Embryol. 183: 589-603.

ten Donkelaar, H.J., R. de Boer-van Huizen, F.T.M. Schouten, and S.J.H. Eggen (1981) Cells of origin of

descending pathways to the spinal cord in the clawed toad (Xenopus laevis). Neuroscience 6: 2297-2312.

ten Donkelaar, H.J., G.C. Bangma, H.A. Barbas-Henry, R. de Boer-van Huizen, and J.G. Wolters (1987) The Brain Stem in a Lizard, Varanus exanthematicus. Adv. Anat. Embryol. Cell Biol. Vol. 107.

Tracey, D.J. (1995) Ascending and descending pathways in the spinal cord. In G. Paxinos (ed.) The Rat Nervous System. Second Ed. Academic Press, San Diego, pp. 67-80.

Tucker, D.C., C.B. Saper, D.A. Ruggiero, and D.J. Reis (1987) Organization of central adrenergic pathways: I. Relationships of ventrolateral medullary projections to the hypothalamus and spinal cord. J. Comp. Neurol. 259: 591-603.

Urbán, L., and G. Székely (1982) The dorsal column nuclei of the frog. Neuroscience 7: 1187-1196.

van den Akker, L.M. (1970) An Anatomical Outline of the Spinal Cord of the Pigeon. Thesis, University of Leiden (The Netherlands).

Vielvoye, G.J. (1977) Spinocerebellar Tracts in the Leghorn (*Gallus domesticus*). Thesis, University of Leiden (The Netherlands).

Wicht, H., and W. Himstedt (1988) Topologic and connectional analysis of the dorsal thalamus of *Triturus alpestris* (Amphibia, Urodela, Salamandridae). J. Comp. Neurol. 267: 545-561.

Wilczynski, W. (1981) Afferents to the midbrain auditory center in the bullfrog Rana catesbeiana. J. Comp. Neurol. 198: 421-433.

Wilczynski, W., and R.R. Capranica (1984) The auditory system of anuran amphibians. Prog. Neurobiol. 22: 1-38. Wilczynski, W., and T.J. Neary (1986) Terminal field organization in the bullfrom torus saminizaularis. Soc

organization in the bullfrog torus semicircularis. Soc. Neurosci. Abstr. 12: 105. Wild, J.M. (1983) Connectional analysis of the

somatosensory system of the pigeon. Soc. Neurosci. Abstr. 9: 244.

Wild, J.M. (1985) The avian somatosensory system. I. Primary spinal afferent input to the spinal cord and brainstem in the pigeon (*Columba livia*). J. Comp. Neurol. 240: 377-395.

Willis, W.D., and R.E. Coggeshall (1991) Sensory Mechanisms of the Spinal Cord. 2nd Ed. Plenum Press, New York.

Yezierski, R.P. (1988) Spinomesencephalic tract: Projections from the lumbosacral spinal cord of the rat, cat and monkey. J. Comp. Neurol. 267: 131-146. Zemlan, F.P., C.M. Leonard, L.M. Kow, and D.W. Pfaff (1978) Ascending tracts of the lateral columns of the rat spinal cord: A study using the silver impregnation and horseradish peroxidase techniques. Exp. Neurol. 62: 298-334.

## Spinothalamic projections in amphibians as revealed with anterograde tracing techniques

Spinal ascending pathways in amphibians: cells of origin and main targets

**COMENTARIOS** 

3.3

En vertebrados terrestres se ha considerado la existencia de dos sistemas de proyecciones espinales ascendentes: 1) El sistema columna dorsal-lemnisco medial formado por proyecciones espinales primarias y no primarias que, a través del funículo dorsal, alcanzan los núcleos de la columna dorsal, los cuales originan el lemnisco medial que asciende hasta el tálamo. 2) El sistema ventrolateral formado por proyecciones secundarias que ascienden inicialmente a través del cuadrante ventral, para alcanzar la formación reticular, el mesencéfalo y el tálamo (Willis y Coggeshall, 1991). Además, en mamíferos (Willis y Coggeshall, 1991) se considera un tercer sistema, denominado espino-cervico-talámico, constituído por las proyecciones que ascienden en el funículo dorsolateral como el tracto espinocervical que termina en el núcleo cervical lateral, el cual a su vez proyecta contralateralmente al mesencéfalo y al tálamo a través del lemnisco medial. Otras proyecciones que ascienden a través del funículo dorsolateral, alcanzan diversas dianas en el tronco cerebral.

Los datos sobre las proyecciones espinales ascendentes en anfibios, disponibles en la bibliografía, proceden de estudios basados en tinciones argénticas en urodelos (Herrick, 1914, 1930; Herrick y Bishop, 1958) y en técnicas degenerativas tanto en anuros (Ebbesson, 1969, 1976; Hayle, 1973) como en urodelos (Nieuwenhuys y Cornelisz, 1971; Wicht y Himstedt, 1988). En los citados trabajos se ha propuesto la existencia de proyecciones espínales a través de los funículos dorsal, dorsolateral y ventrolateral. Sin embargo, en ocasiones únicamente se han presentado como dos sistemas ascendentes. Además, existen en ellos diferencias en cuanto a la nomenclatura, así como a las proyecciones propuestas en cada uno de dichos sistemas. Recientemente se ha descrito en anuros el sistema espino-cervico-talámico (capítulo 5 de la presente memoria), lo que sugiere que, en vertebrados anamniotas, las proyecciones espinales ascendentes están igualmente organizadas en tres sistemas.

La presencia de vías espinotalámicas en el funículo ventrolateral, parece ser un carácter común y compartido en el cerebro de las distintas clases de amniotas. Así, se han demostrado ampliamente aferencias talámicas desde neuronas espinales en mamíferos (Willis y Coggeshall, 1991), aves (Scheneider y Necker, 1989), y reptiles (Ebbesson, 1967; 69; Künzle y Woodson, 1982). En anamniotas únicamente se describieron proyecciones espinotalámicas en una especie de tiburón galeomorfo (Ginglymostoma cirratum) (Ebbesson y Hodde, 1981), y fueron consideradas como un carácter no homólogo a las vías espinotalámicas de amniotas, que evolucionó independientemente (Kevetter y Willis, 1984; Ronan y Northcutt, 1990). En anfibios, hasta el momento, no se habían descrito proyecciones espinotalámicas directas, quizás debido a las limitaciones propias de las técnicas degenerativas (Ebbesson, 1969, 1976; Nieuwenhuys y Cornelisz, 1971; Hayle, 1973; Wicht y Himstedt, 1988).

En el presente capítulo se incluyen dos artículos en los que se ha estudiado la organización de las proyecciones ascendentes espinales en los anuros *Rana perezi y Xenopus laevis*, y en el urodelo *Pleurodeles waltl*, mediante la utilización de técnicas de trazado anterógrado y retrógrado, tanto *in vivo* como *in vitro*, utilizando los trazadores peroxidasa de rábano (HRP), leucoaglutinina de *Phaseolus vulgaris* (PHA-L) o dextrano amina combinada con biotina (BDA). En el primer artículo se presentan evidencias de la existencia de proyecciones espinotalámicas en anfibios, mediante trazado anterógrado. En el segundo artículo se demuestra la organización de las proyecciones espinales ascendentes en tres sistemas fundamentales, funículo dorsal, funículo dorsolateral, y cuadrante vental (funículos ventral y ventrolateral); caracterizándose los principales centros a los que proyectan, así como las células que originan algunos de ellos, y estableciéndose una comparación, desde un punto de vista evolutivo, con los datos disponibles de otros grupos de vertebrados.

#### Provecciones incluídas en el funículo dorsal (DE)

En los experimentos con aplicaciones de diversos trazadores en el asta dorsal, en distintos niveles espinales en Rana perezi y en Xenopus laevis, observamos un patrón de marcaje de proyecciones ascendentes que terminan en el núcleo de la columna dorsal (ver capítulo 5 de la presente memoria), similar al descrito para las ramas ascendentes de las aferencias primarias espinales (Joseph y Whitlock, 1968a, Antal y cols., 1980; Nikundiwe y cols., 1982, M. Muñoz y cols., 1991; A. Muñoz y cols., 1995b), y a las proyecciones ascendentes no primarias que forman el sistema postsináptico de la columna dorsal (PDCS) (ten Donkelaar y de Boer van Huizen, 1991; A. Muñoz y cols., 1995b). Algunas fibras inervan en dicho nivel el polo caudal del núcleo del tracto solitario y la formación reticular. Coincidiendo con la descripción de las aferencias primarias espinales (Joseph y Whitlock, 1968a, Antal y cols., 1980; Nikundiwe y cols., 1982, M. Muñoz y cols., 1991; A. Muñoz y cols., 1995b), un componente de fibras continúa rostralmente, dando fibras terminales en la formación reticular rombencefálica, núcleo del tracto

descendente del nervio trigémino, área octavolateral y capa granular del cerebelo. Sin embargo, hay que señalar que algunas de estas fibras podría corresponder igualmente, a proyecciones no primarias originadas en la médula espinal.

En urodelos existen muy pocos datos en la bibliografía en cuanto a la organización de las aferencias primarias espinales. En nuestros experimentos, con aplicaciones de BDA en raíces dorsales cervicales y lumbares en Pleurodeles wall, hemos observado un patrón de organización similar a nuestros estudios en anuros. Las aferencias primarias espinales, al entrar en la médula espinal, establecen un componente medial cuyas fibras ascienden y descienden en el DF y un componente lateral, equivalente al tracto de Lissauer, que discurre en el funículo dorsolateral. Las fibras del componente medial ascienden en el DF somatotópicamente organizadas, de manera que las aferencias lumbares y cervicales discurren medial y lateralmente en el DF respectivamentre, y alcanzan de la misma forma el núcleo de la columna dorsal (ver capítulo 6 de la presente memoria). Las aferencias lumbares más rostrales se observan en niveles ligeramente anteriores al núcleo de la columna dorsal, mientras que las cervicales continuan ascendiendo a lo largo del rombencéfalo, dorsalmente al tracto descendente del nervio trigémino, para terminar en la capa granular del cerebelo. Dichos resultados coinciden con la distribución de las aferencias braquiales descrita por Roth y Wake (1992) en salamandras de la familia Plethodontidae, si bien dichos autores no observaron proyecciones directas al cerebelo.

En nuestros experimentos en *Pleurodeles* waltl con aplicaciones de BDA en el asta dorsal espinal, en niveles cervicales, se observaron en el DF dos componentes de fibras ascendentes, lateral y medial, que en niveles rombencefálicos están separados por una zona ocupada por aferencias del oido interno (Fritzsch, 1988). El lateral coincide con nuestra descripción de las aferencias primarias, si bien podría igualmente contener axones no primarios del PDCS, presente en otros vertebrados (A. Muñoz y cols., 1995b). El medial, formado por fibras gruesas densamente empaquetadas, se extiende rostralmente hasta niveles medios rombencefálicos, donde termina en el lóbulo de la línea lateral. Dicho componente está formado por aferencias de la segunda raíz del complejo de los nervios glosofaríngeo y vago, cuyas ramas descendentes alcanzan niveles espinales, donde ingresan en el DF (Roth y Wake, 1985).

La presencia de aferencias primarias espinales que ascienden en el DF, para terminar en la región del óbex, distintos centros rombencefálicos y en el cerebelo, constituye un carácter común en el cerebro de los vertebrados, aunque existen algunas diferencias respecto a los distintos lugares de destino descritos en los diferentes trabajos (ver Tabla 1 del artículo 2), que quizás reflejen variaciones en cuanto a las aproximaciones experimentales empleadas para su estudio. Así, se ha observado un patrón de organización similar en agnatos (Northcutt y Ebbesson, 1980; Ronan y Northcutt, 1990; Dubuc y cols., 1993), condrictios (Hayle, 1973a; Ebbesson y Hodde, 1981), teleósteos (Oka y cols., 1986; Ronan y Northcutt, 1990), anfibios (Joseph y Whitlock, 1968a, Antal y cols., 1980; Nikundiwe y cols., 1982, M. Muñoz y cols., 1991; A. Muñoz y cols., 1995b), reptiles (Ebbesson, 1967, 1969; Jacobs, 1968; Joseph y Whitlock, 1968b; Pedersen, 1973; Jacobs y Sis, 1980; Kusuma y ten Donkelaar, 1980; Ebbesson y

Goodman, 1981; Künzle, 1982; Künzle y Woodson, 1983), aves (Karten, 1963; van den Akker, 1970; Wild, 1985) y mamíferos (Willis y Coggeshall, 1991). La presencia de proyecciones espinales no primarias del PDCS en el DF, se ha demostrado hasta el momento en todos los vertebrados terestres (Rustioni, 1973; Angaut-Petit, 1975a,b; Rustioni y Kaufman, 1977; Bennett y cols., 1984; Giesler y cols., 1984; Funke, 1988; ten Donkelaar y de Boervan Huizen, 1991; Pritz y Stritzel, 1994; A. Muñoz y cols., 1995b). Sin embargo, su existencia en agnatos, elasmobranquios y teleósteos no ha sido investigada hasta el momento.

# Provecciones incluídas en el funículo dorsolateral (DLF).

En el presente estudio se ha demostrado la existencia en anuros de un sistema de proyecciones, originado en todos los niveles espinales, que asciende a través del funículo dorsolateral. Hemos observado que rostralmente a la aplicación de los diversos trazadores en distintos niveles espinales, dicho sistema inerva los campos espinales dorsal y lateral, y con mayor densidad el núcleo cervical lateral en el nivel cervical superior. Esta última proyección se ha propuesto como el equivalente en anfibios del tracto espinocervical (ver capítulo 5 de la presente memoria). Algunas de las fibras que alcanzan el núcleo cervical lateral, a través del DLF, corresponden a aferencias primarias braquiales que ascienden por el tracto de Lissauer (Joseph y Whitlock, 1968a, Antal y cols., 1980; Nikundiwe y cols., 1982, M. Muñoz y cols., 1991; A. Muñoz y cols., 1995b). A lo largo del rombencéfalo las fibras del DLF se desplazan ventrolateralmente y ascienden inmediatamente ventrales al tracto descendente del nervio trigémino, organizando proyecciones que alcanzan el polo caudal del núcleo del tracto solitario, y mayoritariamente la formación reticular lateral, y rostralmente el área parabraquial en la región subcerebelosa. Un menor número de fibras penetran en la capa granular del cerebelo, y otras continúan rostralmente para terminar en el polo caudal del núcleo posterodorsal tegmental mesencefálico. En algunas ocasiones se observaron fibras cruzando en la comisura del velo medular anterior. En todos los casos se observó una inervación más densa desde niveles cervicales y torácicos que desde niveles espinales lumbares.

En Pleurodeles waltl, en experimentos con aplicaciones de trazadores a nivel cervical, se observó un patrón de marcaje similar al de anuros. Las fibras ascendentes en el DLF organizan, en niveles cervicales superiores de transición con el rombencéfalo, un campo denso de terminales en la región ventrolateral del asta dorsal. Dicho campo terminal ya fue sugerido en Ambystoma tigrinum por Herrick (1930). En dicha localización se ha considerado la posible existencia del equivalente en urodelos del núcleo cervical lateral de mamíferos, debido a sus proyecciones al torus semicircularis y al tálamo ventral (ver capítulo 6 de la presente memoria). En experimentos con aplicaciones de BDA en la segunda raíz dorsal espinal se observó que algunas aferencias braquiales ascienden en el tracto de Lissauer, a través del DLF, hasta niveles del núcleo glosofaríngeo y alcanzan a nivel del óbex el núcleo cervical lateral (ver capítulo 6 de la presente memoria). Las fibras no primarias del DLF ascienden a través del rombencéfalo en la sustancia blanca inmediatamente ventral al tracto descendente del nervio trigémino hasta niveles más rostrales, dando a lo largo de todo su recorrido fibras varicosas terminales que penetran hasta la gris periventricular entre los niveles de las raíces del complejo glosofaríngeo-vago y la del trigémino. Rostralmente inervan densamente la región subcerebelosa y en menor medida el tegmento mesencefálico caudal.

La existencia del tracto espinocervical en el DLF, descrito inicialmente en mamíferos (ver Willis y Coggeshall, 1991), es un hecho más común en el cerebro de todos vertebrados de lo que se suponía (ver tabla 2 del artículo 2). Su presencia se ha demostrado recientemente en anfibios (ver capítulo 5 de la presente memoria). Además existen evidencias que sugieren que dicho sistema existe en agnatos (Northcutt y Ebbesson, 1980; Ronan y Northcutt, 1990), condrictios (Hayle, 1973a,b; Ebbesson y Hodde, 1981), teleósteos (Hayle, 1973a,b; Finger, 1981; Ito y cols., 1986), reptiles (Ebbesson, 1967; Künzle y Woodson, 1982), así como en aves (van den Akker; 1970; Funke y Necker, 1986; Funke, 1988; Necker, 1991). Otras proyecciones a través del DLF descritas en anfibios en el presente trabajo incluyen las espinosolitarias, espinoreticulares y espinoparabraquiales que son igualmente comunes en todas las clases de vertebrados (ver Tabla 2 del artículo 2). Por último, las proyecciones espinales que a través del DLF alcanzan el tegmento mesencefálico caudal, tanto en anuros como en urodelos, podrían representar el equivalente de las proyecciones espinales en mamíferos que terminan en la sustancia gris periventricular y en el núcleo cuneiforme (Nijensohn and Kerr, 1975; Zemlan et al., 1984; Björkeland and Boivie, 1984; Hylden et al., 1986; Yezierski, 1988; Bernard et al., 1995).

### Provecciones incluídas en el cuadrante ventral ventrolateral (funículos ventral y ventrolateral) (VQ).

En nuestros experimentos con aplicaciones de diversos trazadores en el asta dorsal de distintos niveles espinales en Rana perezi y Xenopus laevis hemos observado que las proyecciones espinales ascendentes presentes en el VQ, son bilaterales aunque con predominio contralateral. Dichas proyecciones se originan en neuronas espinales, cuyos axones se dirigen ventromedialmente en el mismo nivel en el que se encuentra el soma neuronal, ingresan en el VO y ascienden hasta distintos centros supraespinales. El presente estudio de trazado neuronal, demuestra un patrón más amplio en cuanto al número de centros rombencefálicos y mesencefálicos inervados por el VQ que los resultados descritos en trabajos previos, obtenidos mediante técnicas degenerativas (Ebbesson, 1969, 1976; Hayle, 1973a). En niveles rombencefálicos numerosas fibras terminan en los núcleos inferior, medio y superior de la formación reticular medial, mientras que la formación reticular lateral recibe únicamente algunas fibras. Los núcleos motores de los nervios glosofaríngeo y vago, y los del rafe y descendente del nervio trigémino están igualmente inervados. Más rostralmente, las fibras del VO inervan la región situada entre los núcleos motores de los nervios trigémino y facial, y de manera más dispersa la sustancia gris central rombencefálica y los núcleos dorsal y ventral del nervio octavo. Algunas fibras terminan en la región subcerebelosa y en la capa granular del cerebelo. En niveles ístmicos se observaron fibras marcadas en regiones ventromediales, ventrolaterales y mediales al núcleo de istmo. En el mesencéfalo caudal el VQ inerva los núcleos posterodorsal y posteroventral tegmentales, aunque la mayor parte de las fibras presentes en este

nivel, se dirigen dorsalmente para inervar los núcleos laminar, magnocelular, principal y en menor medida el comisural del torus semicircularis, y el techo óptico.

La presencia en el VO de aferencias espinales (presente estudio), trigeminales (M. Muñoz y cols., 1994, A. Muñoz y cols., 1995b), y de los núcleos de la columna dorsal (Wilczynski, 1981; Wilczynski y Neary, 1986; A. Muñoz y cols., 1994b, 1995b) y cervical lateral (A. Muñoz y cols, 1995b, artículo 3 del capítulo 5 de la presente memoria) permiten la llegada de información somatosensorial al torus semicircularis, y por tanto la existencia de un mapa de representación de la superficie corporal en dicho centro, descrito por Comer and Grobstein (1981c), que interviene en la elaboración de respuestas frente a estímulos tactiles (Comer and Grobstein, 1981a,b). En niveles mesencefálicos rostrales hemos observado fibras en la sustancia gris pretectal y pretoral, en la comisura posterior, así como en las regiones anterodorsal y anteroventral mesencefálicas, donde se han descrito el núcleo rojo y el núcleo del fascículo longitudinal medial.

En estudios morfológicos, basados en técnicas degenerativas (Ebbesson, 1969; Hayle, 1973), no se han descrito hasta el momento proyecciones espinotalámicas directas. Sin embargo, Vesselkin y cols. (1971), en un estudio electrofisiológico en *Rana temporaria* mediante estimulación del nervio ciático, demostraron el procesamiento bilateral de información somatosensorial en toda la extensión rostrocaudal del tálamo dorsal.

En los dos artículos del presente capítulo mediante trazado anterógrado se demuestra la existencia de proyecciones espinotalámicas directas, a través del sistema VQ, que alcanzan el tálamo dorsal y ventral. En el tálamo dorsal se observó un escaso número de fibras marcadas en los núcleos posterior, central y en menor medida en el anterior. Sin embargo, la mayor densidad de fibras termina en la parte dorsomedial del tubérculo posterior, y en los núcleos ventrolateral (división ventral) y ventromedial del tálamo ventral.

En urodelos Herrick (1936, Ambystoma tigrinum) sugirió la inervación espinal de la zona medial del tálamo dorsal. Sin embargo, estudios basados en técnicas de degeneración anterógrada (Nieuwenhuys y Cornelisz, 1971) o en técnicas de trazado con HRP (Wicht y Himstedt, 1988) no han confirmado las consideraciones de Herrick. En el presente estudio, en Pleurodeles waltl, se ha demostrado, al igual que en anuros, la presencia de un sistema de fibras ascendentes en el VQ que terminan en la formación reticular rombencefálica, área octavolateral y cerebelo. En el mesencéfalo se observaron fibras terminales marcadas en la sustancia blanca adyacente al torus semicircularis, áreas tegmentales, y en menor número en el techo óptico. En el diencéfalo, la mayoría de las fibras termina en el tálamo ventral, si bien se observó un menor número en las zonas anteroventral y porterodorsal del tálamo dorsal.

No existen datos en la bibliografía sobre la localización y morfología de las neuronas que originan las proyecciones espinales ascendentes en anfibios. En el artículo segundo de este capítulo se presentan resultados correspondientes a experimentos con aplicaciones de BDA en el tálamo ventral, torus semicircularis y núcleo reticular inferior. En todos los casos se observó un mayor número de neuronas contralaterales que ipsilaterales, retrógradamente marcadas en la médula espinal, de acuerdo con los datos de trazado anterógrado anteriormente descritos, procedentes de experimentos con aplicaciones de distintos trazadores en la médula espinal.

En los casos en que se realizaron aplicaciones de BDA en el tálamo se observaron neuronas de diferentes morfologías en niveles cervicales y torácicos, mayoritariamente en el campo dorsal, y en menor número en el lateral y ventral. Las aplicaciones de BDA en el torus semicircularis marcaron células preferentemente en niveles cervicales y en menor grado en segmentos torácicos y lumbares, en las zonas profundas del campo dorsal, y en los campos lateral, ventromedial, ventrolateral y lateral motor. En experimentos con aplicaciones de BDA en el núcleo reticular inferior se obtuvieron neuronas marcadas hasta niveles lumbares superiores, aunque la mayor densidad neuronal se observó en segmentos cervicales, en los campos dorsal (superficial y profundo), lateral, ventromedial y ventrolateral.

En experimentos en *Pleurodeles walti* con aplicaciones de trazadores retrógrados en el tálamo ventral y en el torus semicircularis, resultaron marcadas neuronas, en su mayoría contralaterales preferentemente en el asta dorsal, y en menor número en la zona intermedia y el asta ventral en niveles espinales cervicales.

La presencia de un sistema de proyecciones ascendentes espinales en el VQ, que en ocasiones se ha denominado lemnisco espinal debido a los trabajos de Herrick (1948) en *Ambystoma tigrinum*, es un caracter común en el cerebro de los vertebrados, si bien existen diferencias en cuanto a los distintos tractos incluídos en dicho sistema en cada grupo (ver tabla 3 del artículo 2). Sin embargo, dichas diferencias, al menos en parte, podrían ser debidas a las distintas metodologías aplicadas en cada estudio. La presencia en el VQ del tracto espinoreticular es común en todos los vertebrados. Sin embargo, el tracto espinocerebeloso ventral parece estar únicamente presente en gnatóstomos. Las proyecciones espinomesencefálicas son propias de todas las clases de vertebrados, en particular terminando en la región intercolicular. Keveter y Willis (1984) basándose en resultados negativos en anamniotas, procedentes de estudios previos de degeneración anterógrada (Hayle, 1973a; Ebbesson, 1969), y de trazado neuronal (Neary and Wilczynski, 1977, 1979), describieron la presencia de proyecciones espinotalámicas como un carácter exclusivo de amniotas, con la excepción de algunos condrictios (Ebbesson y Hodde, 1981) en los que su presencia se consideró un carácter que evolucionó independientemente (Keveter y Willis, 1984). Sin embargo, trabajos posteriores en teleósteos (Murakami e Ito, 1985), anfibios (presente capítulo), y posiblemente en agnatos (Ronan y Nothcutt, 1990), han demostrado proyecciones espinotalámicas en anamniotas, por lo que la presencia de dichas proyecciones directas debe considerarse como un carácter común para todos los vertebrados.

# CAPÍTULO 4

Ontogenia del sistema Columna Dorsal-Lemnisco Medial 132

- **4.1.** Early development of dorsal column-medial lemniscal projections in the clawed toad, <u>Xenopus laevis</u>
- 4.2.- Comentarios

Early development of dorsal column-medial lemniscal projections in the clawed toad <u>Xenopus</u> <u>laevis</u>

4.1

### A. Muñoz<sup>b</sup>, R. de Boer-Van Huizen<sup>a</sup>, I. Bergevoet-Vernooy<sup>a</sup> and H.J. ten Donkelaar a

a Department of Anatomy and Embryology, Universyty of Nijmegen, Nijmegen (The Netherlands)
b Department of Cell Biology, Universidad Complutense de Madrid, Madrid (Spain)

Based on: Delvelopmental Brain Research, 74 (1993) 291-294

Key words: Development; Somatosensory system; Dorsal column nucleus; Medial lemniscus; Tract tracing; Anurans.

#### ABSTRACT

In Xenopus laevis fluorescent dextran amines were applied to study the development of the dorsal column-medial lemniscal projection: rhodamine dextran amine was applied at the mesodiencephalic border to retrogradely label the cells of origin of the medial lemniscus in the dorsal column nucleus (DCN); fluorescein dextran amine to the spinal cord to anterogradely label the primary afferent projections to the DCN. The first mesodiencephalic projections were found at stage 51, i.e. almost immediately after spinal afferent fibers had reached the DCN.

In terrestrial vertebrates, basically two systems of ascending spinal projections are found: (1) a primary afferent ascending spinal projection via the dorsal funiculus to the dorsal column nucleus (DCN), giving rise to the medial lemniscal pathway to the thalamus; (2) a secondary afferent projection via the lateral funiculus, i.e. the spinal lemniscus, to the reticular formation, mesencephalon and thalamus<sup>25</sup>. Despite rather extensive somatotopically arranged spinal afferents to the anuran dorsal column nucleus<sup>1,9,13,19</sup>, only a relatively sparse lemniscal channel projecting to the torus semicircularis in the midbrain, and to thalamic levels, appears to be present<sup>11,24</sup>. In frogs somatosensory information reaches the striatum<sup>22</sup> and the medial pallium<sup>10,14,21</sup> via various thalamic nuclei.

Developmental studies<sup>4,18</sup> indicate that the dorsal column-medial lemniscal projection arises rather early in development. In Xenopus laevis, HRP applications at the mesodiencephalic border (see Fig. 1) showed that in late Xenopus laevis tadpoles, neurons dorsal and lateral to the solitary tract were labeled, contralateral to the application side, in keeping with data in adult ranid frogs<sup>11</sup>. This perisolitary band includes a dorsal column nucleus and the caudal part of the descending trigeminal nucleus. In such HRP experiments (for details on techniques see ref. 18) the presence of DCN-thalamic projections at least as early as stage 56 could be shown. Similar observations were made in bullfrog (Rana catesbeiana) tadpoles<sup>4</sup>: retrograde labeling of DCN cells could be obtained by stage VIII/IX<sup>17</sup>, comparable to Nieuwkoop and Faber's<sup>12</sup> stage 54 in Xenopus laevis. These data suggest that the projection of dorsal root ganglion (DRG) cells to the DCN precedes the projection of these second-order cells to the thalamus. Data from the opossum<sup>6-8</sup> are consistent with these observations. In the rat, however, a temporal overlap has been reported between the arrival of primary afferent projections to the gracile nucleus<sup>23</sup>, and the development of thalamic projections from the DCN<sup>2</sup>.

In the present study fluorescent dextran amines, sensitive tracers particularly useful in developmental studies<sup>5</sup>, were used to analyze the early development of the DCN-medial lemniscal projection in Xenopus laevis. (Tetramethyl)rhodamine dextran amine (RDA; D-1817; Molecular Probes Inc., Eugene, OR, USA) was applied at the mesodiencephalic border or at the torus level in the rostral midbrain to retrogradely label the cells of origin of the medial lemniscus. Contralaterally, fluorescein dextran amine (FDA; D-1820) was applied to the dorsal funiculus or to DRG at cervical levels to characterize that part of the perisolitary band receiving primary spinal afferents. In about 35 Xenopus laevis tadpoles ranging from stage 47, i.e. at the time of the appearance of the limb buds, until stage  $66^{12}$ (metamorphosis complete), RDA and FDA, recrystallized from distilled water onto sharp tungsten needles, were applied under tricaine methanesulphonate (0.1 mg/ml tap water; MS 222; Sandoz). After survival times of 2-3 days, animals were reanesthetised with an overdose of MS222, and perfused through the heart with 0.1 M phosphate buffer (pH 7.4) followed by a fixative containing 4% paraformaldehyde in phosphate buffer. The brain and (rostral) spinal cord were dissected out, embedded in polyacrylamide<sup>18</sup>, left overnight in 15% saccharose in 0.1 M phosphate buffer, and cut on a freezing

microtome at 40  $\mu$ m, either transversally or horizontally. They were mounted in glycerin-gelatin. The pattern of labeling in representative experiments is shown in Fig. 2.

Following RDA application at the mesodiencephalic border or rostral tegmentum mesencephali (including the torus semicircularis), retrogradely labeled cells were observed in the dorsal column nucleus as early as stage 51 (see Fig. 2A). The localization of this neuron population coincides with the place of arborization of ascending collaterals of spinal primary afferents (PAFs; Fig. 2A, arrow). FDA-labeled PAFs were found ascending through the dorsal funiculus and along Lissauer's tract. At the level of the dorsal column nucleus the PAFs give off collaterals that pass ventrally into the grey matter where the RDA labeled cells were found (Fig. 2A). The dendrites of these cells extend laterally and dorsally to the area of FDA labeling. A few laterally situated neurons extend their dendrites dorsomedially. The axons of the DCN neurons could be traced ventromedially to cross the midline towards the contralateral ventral ascending pathway in the brainstem. In stage 51 tadpoles, only a few labeled cells were found. The number of labeled DCNneurons increased from stage 51 to 55 (e.g. Fig. 2B). From stage 55 on a more or less adult pattern of rostrally projecting DCN cells could be observed (see Fig. 2C,D).

By applying FDA to the dorsal funiculus, in general all fibers passing rostrally at the level of application, mostly the third spinal segment (brachial enlargement), were labeled. In all of the stages studied i.e. from stage 47 until 66, labeled PAFs could be observed (Fig. 2), but a progressive increase of the projections in the youngest stages studied (47-52) was observed. In a previous study<sup>18</sup> it was shown that around stage 48 projections from DRG cells reach the level of the developing DCN, but only from nonlimb bud-innervating DRG. Limb bud innervating ganglia give rise to ascending collaterals definitively later in development. At stage 52 a few ascending branches extended into the brain stem<sup>20</sup>. By stage 53 all spinal ganglia including lumbar DRG send ascending collaterals to the brain stem<sup>18</sup>. In addition to the anterogradely labeled PAFs, a few retrogradely labeled neurons were found ipsilaterally in the DCN, at least as early as stage 51. The number of caudally projecting DCN cells increased in later stages.

The present data show that the anuran dorsal column-medial lemniscal system arises earlier than previously thought<sup>4,18</sup>. The first mesodiencephalic projections from the dorsal column nucleus were found at stage 51, i.e. not much later than projections of DRG cells reach the area of the DCN. It should be noted, however, that the first spinal projections reaching the developing dorsal column nucleus arise from thoracic DRG cells<sup>3,15,16,18</sup>. During development, primary sensory neurons innervate peripheral targets and then form central connections appropriate to these targets<sup>15,16,20</sup>. The projection of thoracic DRG cells to the DCN precedes the projection of these second-order cells to the thalamus. It seems likely that the projections of cervical and lumbar DRG cells to the DCN also precede the projection of related second-order cells giving rise to the medial lemniscus. Current research is focussed on the application of the sensitive tracer, biotylinated dextran amine, to dorsal root ganglia, in combination

:

with the application of this tracer at the mesodiencephalic border. In this way it can be established which DRG cells innervate particularly DCN neurons.

The authors would like to thank Mr. Ron Engels and Mr. Henny Eikholt for breeding and taking care of the animals, Mr. Jelle Eigenstein for help with the confocal laser microscope at the Department of Experimental Zoology, Mr. Cor de Bruyn for help with the photomicrographs, and Mrs. Cocky Udo for typing the manuscript.



Figure 1: Reconstruction of the caudal brain stem and rostral spinal cord in a late *Xenopus laevis* tadpole (stage 60). HRP was applied at the mesodiencephalic border. Small dots indicate the labeled cells (with dendrites), contralateral to the application side of the tracer; triangles indicate a few ipsilaterally found neurons; DRG 3-5 are shown. Photomicrographs show examples of the labeling of the perisolitary band, modified by the DAB-staining techniques<sup>18</sup> (magnification x 100). Arrows indicate labeled medial lemniscal fibers.



Figure 2: Transverse sections at the obex level of *Xenopus laevis*, showing RDA labeled neurons in the developing dorsal column nucleus. A: in a stage 51 tadpole (magnification x400). B: in a stage 53 tadpole (magnification x550). Superposed pictures A and B were taken with a Zeiss fluorescence microscope with appropriate filter combinations. C,D: in a stage 61 tadpole (magnification x215), confocal microscopy (Bio-Rad Lasersharp MRC-500). Arrows indicate the FDA labeled spinal primary afferents.

1 Antal, M., Tornai, 1. and Székely, G., Longitudinal extent of dorsal root fibres in the spinal cord and brain stem of the frog, Neuroscience, 5 (1980) 1311-1322.

2 Asanuma, C., Ohkawa, R., Stanfield, B.B. and Cowan, W.M., Observations on the development of certain ascending inputs to the thalamus in rats. 1. Postnatal development, Dev. Brain Res., 41 (1988) 159-170.

3 Clarke, J.D.W., Tonge, D. and Holder, N., Stage dependent restoration of sensory dorsal columns following spinal cord transection in anuran tadpoles, Proc. R. Soc. Lond. (Biol.), 227 (1986) 67-82.

4 Forehand, C.J. and Farel, P.B., Spinal cord development in anuran larvae. II. Ascending and descending pathways, J. Comp. Neurol., 209 (1982) 395-408.

5 Glover, J.C., Petursdottir, G. and Jansen, J.K.S., Fluorescent dextran-amines used as axonal tracers in the nervous system of the chicken embryo, J. Neurosci. Methods, 18 (1986) 243-254.

6 Johnson, J.I., Hamilton, T.C., Hsung, J.C. and Ulinski, P.S., Gracile nucleus absent in adult opossums after leg removal in infancy, Brain Res., 38 (1972) 421-424.

7 Martin, G.F., Culberson, J.L. and Hazlett, J.C., Observations on the early development of ascending spinal pathways: studies using the North American opossum, Anat. Embryol., 166 (1983) 191-207.

8 Martin, G.F., Cabana, T., Hazlett, J.C., Ho, R. and Waltzer, R. Development of brain stem and cerebellar projections to the diencephalon with notes on thalamocortical projections: studies in the North American opossum, J. Comp. Neurol., 260 (1987) 186-200.

9 Muñoz, M., Muñoz, A., González, A., Machin, C. and Rua, C. Ascending projections from the spinal cord in the anuran amphibian *Rana ridibunda*, Eur. J. Neurossi, Suppl. 4 (1991) 24.

10 Neary, T.J., The pallium of anuran amphibians. In E.G. Jones and A. Peters (Eds.), Cerebral Cortex, Vol. 8A, Plenum, New York, 1990, pp. 107-138.

11 Neary, T.J. and Wilczynski, W., Ascending thalamic projections from the obex region in ranid frogs, Brain Res., 138 (1977) 529-533.

12 Nieuwkoop, P.D. and Faber, J., Normal Table of *Xenopus laevis* (Daudin), 2nd edn., Elsevier, Amsterdam, 1967.

13 Nikundiwe, A.M., de Boer-van Huizen, R. and ten Donkelaar H.J., Dorsal root projections in the clawed toad (*Xenopus laevis*) as demonstrated by anterograde labeling with horseradish peroxidase, Neuroscience, 7 (1982) 2089-2103.

14 Northcutt, R.G. and Ronan, M., Afferent and efferent connections of the bullfrog medial pallium, Brain Behalv. Erol., 40 (1992) 1-16.

15 Smith, C.L. and Frank, E., Specificity of sensory projections to the spinal cord during development in bullfrogs, J. Comp. Neurol., 269 (1988a) 96-108.

16 Smith, C.L. and Frank, E., Peripheral specification of sensory connections in the spinal cord, Brain Behav. Evol., 31 (1988b) 227-242.

17 Taylor, A.C. and Koliros, J.J., Stages in the normal development of *Rana pipiens* larvae, Anat. Rec., 94 (1946) 7-23.

18 ten Donkelaar, H.J. and de Boer-van Huizen, R., Observations on the development of ascending spinal pathways in the clawed toad, Xenopus laevis, Anat. Embryol., 183 (1991) 589-603.

19 Urbán, L. and Székely, G., The dorsal column nuclei of the frog, Neuroscience, 7 (1982) 1187-1196.

20 van Mier P. and ten Donkelaar, H.J., The development of primary afferents to the lumbar spinal cord in *Xenopus laevis*, Neurosci. Lett., 84 (1988) 35-40.

21 Vesselkin, N.P., Agayan, A.L. and Nomokonova, L.M., A study of thalamo-telencephalic

afferent system in frogs, Brain Behav. Evol., 4 (1971) 295-306.

22 Vesselkin, N.P., Ermakova, T.V., Kenigfest, N.B. and Goikovic M., The striatal connections in frog *Rana* temporaria: an HRP study, J. Hirnforsch., 21 (1980) 381-392.

23 Wessels, W.J.T., Feirabend, H.K.P. and Marani, E., Development of projections of primary afferent fibers from the hindlimb to the gracile nucleus: a WGA-HRP study in the rat, Dev. Brain Res., 63 (1991) 265-279.

24 Wilczynski, W. and Neary, T.J., Terminal field organization in the bullfrog torus semicircularis, Soc. Neurosci. Abstr., 12 (1986) 105.

25 Willis, W.D. and Coggeshall, R.E., Sensory Mechanisms of the Spinal Cord. 2nd edn. Plenum. New York. 1991.

۰.

Early development of dorsal column-medial lemniscal projections in the clawed toad <u>Xenopus</u> <u>laevis</u>

:

# **COMENTARIOS**

- 4.2

En vertebrados terrestres el sistema columna dorsal-lemnisco medial está formado por las proyecciones aferentes espinales, primarias y no primarias, que alcanzan los núcleos de la columna dorsal a través del funículo dorsal, y por las proyecciones eferentes de éstos que constituyen el lemnisco medial, el cual asciende contralateralmente hasta el mesencéfalo y el tálamo (Willis y Coggeshall, 1991).

En anfibios existe un sistema de aferencias espinales primarias, somatotópicamente organizado, que alcanza el núcleo de la columna dorsal (DCN) (Antal y cols., 1980; Nikundiwe y cols., 1982; Urbán y Székely, 1982). Sin embargo, parece existir un canal lemniscal poco desarrollado, que proyecta al torus semicircularis en el mesencéfalo, y a niveles talámicos (Neary y Wilczynski, 1977; Wilczynski y Neary, 1986).

Estudios sobre el desarrollo larvario de las proyecciones espinales ascendentes (Forehand y Farel, 1982; ten Donkelaar y de Boer-van Huizen, 1991) indican que la proyección columna dorsal-lemnisco medial aparece en estadios tempranos del desarrollo. ten Donkelaar y de Boer-van Huizen (1991) demostraron que las aferencias primarias espinales alcanzan la región del óbex entre los estadios 48 y 53 (Nieuwkoop y Faber, 1967) en Xenopus laevis. Forehand y Farel (1982) en experimentos con aplicaciones talámicas de HRP en distintos estadios del desarrollo en Rana catesbeiana, obtuvieron marcaje retrógrado de neuronas del DCN en animales del estadio VIII/IX (Taylor y Kollros, 1946), comparable al estadio 54 de Nieuwkoop y Faber (1967) en Xenopus laevis. Estos datos sugieren que el desarrollo de la proyección de las neuronas ganglionares de las raíces dorsales al DCN, precede al de la proyección talámica de este núcleo.

En mamíferos, los datos obtenidos en la zarigüella (Johnson y cols., 1972; Martin y cols., 1983, 1987) coinciden con dichas observaciones. En la rata, sin embargo, se ha descrito un solapamiento temporal entre la llegada de aferencias primarias al núcleo gracilis, (Wessels y cols., 1991) y el desarrollo de las proyecciones talámicas desde los núcleos de la columna dorsal (Asanuma y cols., 1988).

En el presente capítulo se ha estudiado el desarrollo de las proyecciones columna dorsallemnisco medial en *Xenopus laevis*, mediante la aplicación en animales entre los estadios 47 y 66 del desarrollo, de dextrano aminas combinadas con rodamina (RDA) en el torus semicircularis y en la zona mesodiencefálica, con objeto de marcar en el DCN retrógradamente las células de origen del lemnisco medial. En los mismos experimentos se aplicó contralateralmente dextrano aminas combinadas con fluoresceína (FDA) en la médula espinal o en los ganglios de la raíces dorsales, con objeto de marcar las proyecciones aferentes al DCN.

Las aplicaciones de RDA en la zona mesodiencefálica o en el torus semicircularis marcaron neuronas en el DCN desde el estadio 51, en el que observa un escaso número de células, que se incrementa progresivamente entre los estadios 51 y 55. A partir del estadio 55 se establece un patrón, más o menos semejante al del adulto, de neuronas del DCN de proyección rostral. La localización de esta población neuronal, coincide con el sitio de arborización de las fibras primarias espinales, marcadas anterógradamente con FDA desde los ganglios dorsales o la médula espinal, que ascienden a través del funículo dorsal y del tracto de Lissauer en el funículo dorsolateral. Asimismo se observó que las dendritas de las neuronas del DCN, marcadas con RDA, se extienden dorsal y lateralmente hacia el área ocupada por las fibras marcadas con FDA. En todos los estadios estudiados se pudieron observar aferencias primarias espinales marcadas, pero se produce un incremento progresivo del número de fibras entre los estadios 47 y 52.

En un estudio previo ten Donkelaar y de Boer-van Huizen (1991) demostraron que hacia el estadio 48, las proyecciones desde neuronas de los ganglios de la raíces dorsales alcanzan el DCN, pero únicamente desde los ganglios torácicos que no inervan los primordios de las extremidades. Los ganglios que inervan las extremidades originan ramas ascendentes en estadios más avanzados. En el estadio 52 algunas fibras alcanzan el tronco cerebral (van Mier y ten Donkelaar, 1988), y en el estadio 53 todos los ganglios espinales, incluyendo los lumbares, envían colaterales ascendentes al tronco cerebral (ten Donkelaar y de Boer-van Huizen, 1991).

El presente estudio demuestra que el sistema columna dorsal-lemnisco medial en anuros, se origina en estadios anteriores a lo que se había descrito previamente (Forehand y Farel, 1982; ten Donkelaar y de Boer-van Huizen, 1991). Nuestros resultados indican que la proyección de las neuronas ganglionares torácicas al DCN, precede al desarrollo de la proyección talámica de dicho núcleo. Las primeras proyecciones mesodiencefálicas desde el DCN se originan en el estadio 51, poco después de que este núcleo reciba las primeras aferencias espinales, desde células ganglionares torácicas (estadio 47) (Clarke y cols., 1986; Smith y Frank, 1988a, b; ten Donkelaar y de Boer-van Huizen, 1991). Durante la ontogenia, las neuronas sensoriales primarias inervan primero las dianas periféricas y posteriormente establecen conexiones centrales apropiadas a dichas dianas (Smith y Frank, 1988a, b; van Mier and ten Donkelaar, 1988), ten Donkelaar y de Boer-van Huizen (1991) observaron que los axones de las neuronas de los ganglios cervicales y lumbares, alcanzan el área del DCN en los estadios 52-53. Nuestros resultados demuestran que durante los estadios 51 a 55 se produce un incremento en el número de células del DCN cuyos axones llegan al borde mesodiencefálico. Por lo tanto, parece probable que el desarrollo de las proyecciones primarias al DCN desde los ganglios cervicales y lumbares (estadios 52-53), preceda igualmente al aumento de la proyección lemniscal desde las neuronas del DCN, relacionadas con díchas aferencias, pudiendo existir así, en la ontogenia de las proyecciones somatosensoriales, un patrón de determinación periférico-central del sistema columna dorsal-lemnisco medial.



